

ORIGINAL ARTICLE

The occurrence of three species of the genus *Oscheius* Andrassy, 1976 (Nematoda: Rhabditida) in Iran

Ahmad Valizadeh¹, Shila Goldasteh^{1*}, Zahra Rafiei-Karahroodi¹, Majid Pedram²¹ Department of Entomology, Arak Branch, Islamic Azad University, Arak, Iran² Department of Plant Pathology, Faculty of Agriculture, Tarbiat Modares University, Tehran, Iran

Vol. 57, No. 3: 248–255, 2017

DOI: 10.1515/jppr-2017-0035

Received: December 31, 2016

Accepted: August 28, 2017

*Corresponding address:
shilagoldasteh@yahoo.com

Abstract

Due to importance and effectiveness of some entomopathogenic or insect parasitic nematodes in controlling of agricultural pests, or pests of non-agricultural plants, a study was conducted in order to identify the species of this group of nematodes in city of Tehran. As the result, three species belonging to the genus *Oscheius* were recovered in association with bark samples having the bark beetle galleries. Morphological and molecular data were provided for two recently recovered species of the genus, namely *O. necromenus* and *O. onirici*. Molecular data were also provided for a recently recovered isolate of *O. tipulae*. All three species were recovered in association with bark samples collected from dead trees in the city of Tehran. Morphological characters and morphometric data of the two aforementioned species are in accordance with the data given in their original descriptions. One recovered individual from a small bark sample characterized by its short body length was sequenced for its 28S and internal transcribed spacer (ITS) rDNA loci, and the results of BLAST search using the newly obtained partial sequences revealed that it belonged to *O. tipulae*. Molecular phylogenetic studies revealed recently sequenced Iranian populations of *O. onirici* and *O. tipulae* forming a clade with other isolates/populations of these species in ITS tree with maximal Bayesian posterior probability (BPP), and presently sequenced isolates of *O. tipulae* and *O. necromenus* form a clade with other isolates of these species in 28S tree. The two species *O. onirici* and *O. necromenus* were reported in Iran for the first time.

Key words: Bayesian posterior probability, ITS, molecular analysis, *Oscheius necromenus*, *Oscheius onirici*, *Oscheius tipulae*

Introduction

The genus *Oscheius* was established by Andrassy (1976) with *O. insectivorus* (Körner, 1954) Andrassy, 1976 as its type species. The genus is in the family Rhabditidae Örley, 1880 (Andrassy 2005). According to Sudhaus (1976) and Sudhaus and Hooper (1994), species of the genus form two separate subclades. The 42 known species of the genus are clustered in two *Insectivora* and *Dolichura* groups (Tabassum *et al.* 2016). According to Liu *et al.* (2012), different types of host association are known for the species of the genus in the *Insectivora* group. In a recent study (Sudhaus 2016), host associations of *Dolichura* group members with insects were discussed. Some types

of associations with insects have also been reported for *Dolichura* group members, like the relation of *O. pherosophi* (Smart and Nguyen 1994). Sudhaus (2011) with cadavers of the bombardier beetle (Smart and Nguyen 1994). Torrini *et al.* (2015) reported and described the first entomopathogenic nematodes (EPN) belonging to the *Dolichura* group. This finding was further studied/criticized by Campos-Herrera *et al.* (2015).

The genus has also some unique biological characters, making it a suitable candidate for genetic studies and a third model nematode species (Sommer 2000; Dichtel-Danjoy and Félix 2004 a, b).

Several nematodes associated with the bark of dead or dying trees having beetle galleries or recovered from organic material in Iran have been recently described (Pedram *et al.* 2011, 2012; Atighi *et al.* 2012; Aliramaji *et al.* 2014a, b, 2015; Ghaemi *et al.* 2015; Miraeiz *et al.* 2015; Alvani *et al.* 2016). In a short report, Hassani-Kakhaki *et al.* (2012) reported the species *O. tipulae* (Lam and Webster 1971) in Iran. However, no morphological and/or morphometric data were provided for the recovered population. Occurrence of the species *O. rugaensis* (Zhang *et al.* 2012) in Iran was reported by Darsouei *et al.* (2014). Just recently, Torrini *et al.* (2016) used molecular data of an Iranian isolate of *O. tipulae* in their phylogenetic analysis of several isolates of the species from several geographic regions. Recently, a study by Shahabi *et al.* (2016) was carried out on some free-living nematodes of northern Iran. During our recent samplings, some specimens belonging to three species of the genus *Oscheius* were recovered from the city of Tehran. The present study aims to perform morphological and molecular studies on recovered populations of two species *O. necromenus* (Sudhaus and Schulte 1989) and *O. onirici* Torrini, Mazza, Carletti, Benvenuti, Roversi, Fanelli, De Luca, Troccoli & Tarasco, 2015 and to characterize the single individual of *O. tipulae* with its molecular data. The two former species were reported in Iran for the first time and molecular data were provided for the recovered isolate of *O. tipulae*.

Materials and Methods

Sampling, extracting, mounting and taxonomy

Wood and bark samples were collected in the city of Tehran. Nematodes were extracted from bark samples having bark beetle galleries using the tray method (Whitehead and Hemming 1965). Nematodes were heat-killed by adding boiling 4% formalin solution, transferred to anhydrous glycerin according to De Grisse (1969), mounted on permanent slides and examined using a Nikon Eclipse E600 light microscope. Photographs were taken using an Olympus DP72 digital camera attached to an Olympus BX51 microscope powered with differential interference contrast (DIC).

Morphological characters and morphometric data, given in original descriptions and the data given by Tabassum *et al.* (2016), were used for morphological identification.

DNA extracting, PCR and sequencing

DNA was extracted from single individuals using the proteinase K method of Soleymanzadeh *et al.* (2016)

or according to Pedram *et al.* (2015). In the first method, a single nematode individual was transferred to an Eppendorf tube containing 1 μ l proteinase K (CinnaGen, Tehran) (10 mg \cdot ml⁻¹) and 49 μ l of extraction buffer [worm lysis buffer, *cf.* Williams *et al.* (1992); containing: 50 mM KCl, 10 mM Tris-Cl pH 8.3, 2.5 mM MgCl₂, 0.45% NP40, and 0.45% Tween 20, frozen at -80°C (20 min), followed by incubation at 65°C (2 h) and then at 95°C (10 min). DNA samples were stored at -20°C until used as polymerase chain reaction (PCR) templates.

PCR was carried out in a total volume of 30 μ l (19.2 μ l distilled water, 3 μ l 10 \times PCR buffer, 0.6 μ l 10 mM dNTP mixture, 1.2 μ l 50 mM MgCl₂, 1.2 μ l of each primer (10 pmol \cdot μ l⁻¹), 0.6 μ l of Taq DNA polymerase (5 unit \cdot μ l⁻¹, CinnaGen, Tehran, Iran) and 3 μ l of DNA template). The thermal cycling program for amplifying three genomic fragments [18S rDNA, 28S rDNA D2/D3 and internal transcribed spacer (ITS)] was as follows: denaturation at 95°C for 4 min, followed by 32 cycles of denaturation at 94°C for 30 s, annealing at 55°C for 40 s, and extension at 72°C for 80 s. A final extension was performed at 72°C for 10 min. Primers for amplification of 28S rDNA D2/D3 were forward primer D2A (5'-ACAA GTACCGTGAGGGAAAGT-3') and reverse primer D3B (5'-TGCGAAGGAACCAGCTACTA-3') (Nunn 1992). A combination of the primers TW81 (5'-GTTTCCGTAGGTGAACCTGC-3') (forward) and AB28 (5'-ATATGCTTAAGTTCAGCGGGT-3') (reverse) (Joyce *et al.* 1994) and 5.8SM5 (5'-GG CGCAATGTGCATTCGA-3') (Zheng *et al.* 2000) were used for amplifying the ITS fragment. The PCR products were sequenced in both directions using the same primers with an ABI 3730XL sequencer (Applied Biosystems) at Macrogen (Seoul, South Korea). Newly obtained sequences of the studied species were deposited in GenBank (accession numbers: KY366261 and KY366262 for partial ITS sequences of *O. onirici* and *O. tipulae* and KY366263 and KY366264 for partial 28S rDNA D2/D3 sequence of *O. tipulae* and *O. necromenus* respectively).

Phylogenetic analysis

The newly obtained ITS and 28S rDNA D2/D3 sequences of the recovered populations were compared with those of other nematode species available in GenBank using the BLAST homology search program. The ITS rDNA sequences of 20 species/isolates of the genus *Oscheius* and one outgroup species (accession numbers and full species names in Fig. 1) and 27 sequences of species/isolates of 28S rDNA D2/D3 belonging to the genus *Oscheius*, some unidentified rhabditid taxa and one outgroup species (accession numbers and full species names in Fig. 2) were aligned using




Fig. 1. Bayesian tree inferred under the GTR + G model using ITS sequence of *Oscheius* spp. Posterior probability values exceeding 50% are given on appropriate clades




Fig. 2. Bayesian tree inferred under the HKY + G model using 28S rDNA D2/D3 fragments of *Oscheius* spp. Posterior probability values exceeding 50% are given on appropriate clades

MUSCLE (Edgar 2004) as implemented in MEGA6 (Tamura *et al.* 2013). The online version of Gblocks 0.91b (Castresana 2000) was used to eliminate ambiguous parts of the alignment, with all three options for a less stringent selection (http://molevol.cmima.csic.es/castresana/Gblocks_server.html). The most appropriate model of nucleotide substitution was selected using the Akaike information criterion in MrModeltest 2 (Nylander 2004). The general time reversible model including gamma distribution for rates across sites (GTR + G) was selected for 28S and a Hasegawa-Kishino-Yano model including gamma distribution for rates across sites (HKY + G) was selected for ITS dataset. Bayesian inference (BI) was performed using MrBayes v3.1.2 (Ronquist and Huelsenbeck 2003) running the chains for one million generations. After discarding burn-in samples, the remaining samples were retained for further analyses. The Markov chain Monte Carlo (MCMC) method within a Bayesian framework was used to estimate the posterior probabilities of the phylogenetic trees (Larget and Simon 1999) using the 50% majority rule. Burn-in was determined upon stabilization of log likelihood using TRACER v1.5 (Drummond and Rambaut 2007). For both ITS and 28S datasets, the species *Steinernema carpocapsae* (Weiser, 1955) Wouts *et al.* (1982) was used as outgroup taxon. The resultant tree files were visualised using Dendroscope V.3.2.8 (Huson and Scornavacca 2012) and re-drawn in Core-IDRAW software version 16.

Results and Discussion

Oscheius necromenus (Sudhaus and Schulte 1989)

This species was recovered from bark samples of dead grapevine trees in the Yaftabad region, Tehran. Morphological characters and morphometric data of this population are in full agreement with the data given in the original description of the species and the morphometric data ranges given by Tabassum *et al.* (2016) (Fig. 3, Table 1). This species is new for Iran's nematode fauna.

Oscheius onirici Torrini, Mazza, Carletti, Benvenuti, Roversi, Fanelli, De Luca, Troccoli & Tarasco, 2015

The presently studied population of this species was in association with galleries on bark samples of *Pinus eldarica* trees in the Lavizan region, Tehran. No morphological or morphometric differences were observed between the presently studied population and the original data given by Torrini *et al.* (2015) (Fig. 4,

Table 1. Morphometric data of two species *Oscheius onirici* and *O. necromenus* recovered in the present study. All measurements are in μm and in the form mean \pm SD (range)

Measurements	<i>O. necromenus</i>	<i>O. onirici</i>
N	10 ♀♀	8 ♀♀
L	1189 \pm 193 (958–1580)	728.5 \pm 102.0 (506–842)
a	17.2 \pm 3.2 (13.6–24.8)	30.4 \pm 14.3 (16.2–49.5)
b	6.8 \pm 1.1 (5.1–8.5)	5.8 \pm 0.9 (4.4–6.7)
c	11.3 \pm 2.7 (8.0–15.5)	9.9 \pm 1.2 (8.1–11.5)
c'	4.7 \pm 1.5 (2.4–7.0)	5.3 \pm 1.2 (3.9–6.7)
V	49.5 \pm 3.5 (41.4–523)	49.0 \pm 1.1 (47.6–5032)
Stoma	15 \pm 2 (13–18)	15.0 \pm 1.2 (13–17)
Pharynx	177.0 \pm 17.5 (143–193)	125 \pm 12 (113–152)
Head-vulva	587.5 \pm 104.5 (467–817)	373.0 \pm 29.5 (330–423)
Head-anus	1081 \pm 199 (853–1478)	682 \pm 48 (608–755)
Body width	72 \pm 21 (49–116)	28.5 \pm 12.5 (15–48)
Anal body width	25.0 \pm 7.8 (16–42)	14.5 \pm 2.7 (12–20)
Tail	108 \pm 13.5 (91–127)	78.0 \pm 10.3 (61–88)

N – number of individuals, L – body length, a – body length/body width, b – body length/length of pharynx, c – body length/tail length, c' – tail length/body width at anus level, V – anterior end to vulva \times 100/body length

Table 1). This is the first report of this species occurring in Iran.

Oscheius tipulae (Lam and Webster 1971)

Only one specimen of this species was recovered (a damaged female, not suitable for rearing on Nutrient Agar plates) from a bark sample. Its small body size prompted us to sequence it for both 28S rDNA D2/D3 and ITS fragments. The results confirmed that it belongs to *O. tipulae*.

Molecular phylogenetic relationships

Sequencings of ITS rDNA of *O. onirici* and *O. tipulae* yielded single fragments of 495 and 754 nt, respectively. The size of 28S rDNA D2/D3 fragments of *O. necromenus* and *O. tipulae* was 610 and 599 nt.




Fig. 3. Iranian population of *Oscheius necromenus*; A–D, F – female, E – male: A – anterior region, B – vagina, C – pharyngeal bulb, D – tail, E – cloacal region, F – juveniles inside the body of the female (scale bars: A, C, D = 10 μ m, the rest = 20 μ m)



Fig. 4. Iranian population of *Oscheius onirici* – female: A – anterior region, B – pharynx, C – vulval region, D – tail (scale bars = 10 μ m)

Blast search using newly obtained ITS and 28S rDNA sequences of the sequenced isolate of *O. tipulae* revealed that the ITS sequence has 99–100% identity with the available ITS sequences of the species in the database. Blast search using 28S rDNA D2/D3 fragment yielded the same result. Blast search using partial ITS sequence of Iranian isolate of *O. onirici* revealed that the sequence has 100% identity with the ITS sequences of the species provided in its original description and sequences of other isolates deposited in GenBank. The 28S rDNA D2/D3 sequence of *O. necromenus* had 100% identity with an isolate of the species deposited in the database (accession number: KT884891). For reconstructing the ITS tree, a total number of 20 sequences of species/isolates of the genus *Oscheius* and one outgroup taxon (*Steinernema carpocapsae*, GQ421615) were used. Figure 4 presents the tree inferred using the aforementioned dataset. In this tree, the Iranian isolate of *O. tipulae* formed a clade with other isolates of the species with maximal BPP. The Iranian isolate of *O. onirici* also formed a fully supported monophyletic group with other isolates of the species. In 28S tree, sequences of 27 species/isolates of *Oscheius* spp. including one outgroup species (*S. carpocapsae*, KJ950293) were used. The Iranian isolate of the species that was sequenced in the present study formed a clade with other isolates of the species and some other unidentified species/isolates (most probably belonging to *O. tipulae*). The Iranian isolate of the species *O. necromenus* was recovered and sequenced while the present study also formed a clade with a sequence labelled as *O. cf. necromenus* (accession number: KT884891) number and one unidentified isolate, probably belonging to this species.

In this study three species belonging to the genus *Oscheius* were recovered from the city of Tehran. From the recovered species, the species *O. necromenus* belongs to the *Insectivora* group, and two other species belong to the *Dolichura* group. As Campos-Herrera *et al.* (2015) cited, there are true entomopathogenic forms in the former group, making them suitable bio-control agents against soil-dwelling pests, harmful for agricultural crops and natural ecosystems. The entomopathogenicity of the species in the *Dolichura* group however needs further studies for confirmation (Campos-Herrera *et al.* 2015). Recovery of three species of the genus from Tehran suggests that it is possible to have higher species diversity of the genus *Oscheius* in Iran, and needs further research and extensive samplings. The potential of some tentative endemic entomopathogenic forms in controlling endemic agricultural pests is another issue, and could be tested using endemic entomopathogenic strains.

In the present study, two species *O. onirici* and *O. necromenus* were characterized with their morphological and molecular data, however, the species

O. tipulae was successfully characterized with molecular sequences of two genomic fragments. The usefulness of molecular data to distinguish this species from its closely related forms has already been documented by Félix *et al.* (2001).

Acknowledgements

We appreciate the kind help of Prof. Marie-Anne Félix (Institute of Biology of the Ecole Normale Supérieure) and Dr. Raquel Campos-Herrera (Universidade do Algarve, Portugal).

References

- Aliramaji F., Pourjam E., Atighi M.R., Pedram M. 2014a. *Devibursaphelenchus kheirii* sp. n. (Nematoda: Ektaphelenchinae) from Iran with remarks on *Devibursaphelenchus* Kakuliya, 1967. *Nematology* 16 (9): 1069–1078. DOI: <https://doi.org/10.1163/15685411-00002833>
- Aliramaji F., Pourjam E., Atighi M.R., Roshan-Bakhsh A., Ye W., Pedram M. 2014b. Description of *Ektaphelenchoides poinari* sp. n. (Nematoda: Ektaphelenchinae) from Iran with compendium of the valid species of the genus *Ektaphelenchoides* Baujard, 1984. *Russian Journal of Nematology* 22: 11–22.
- Aliramaji F., Pourjam E., Karegar A., Atighi M.R., Pedram M. 2015. Description of *Ektaphelenchoides caspiensis* n. sp. (Nematoda: Ektaphelenchinae) from Iran. *Nematology* 17 (2): 169–177. DOI: <https://doi.org/10.1163/15685411-00002862>
- Alvani S., Mahdikhani-Moghadam E., Giblin-Davis R.M., Pedram M. 2016. Description of *Ektaphelenchus berbericus* n. sp. (Rhabditida: Ektaphelenchinae) from eastern Iran. *Nematology* 18 (9): 1063–1077. DOI: <https://doi.org/10.1163/15685411-00003015>
- Andrásy I. 1976. *Evolution as a Basis for the Systematization of Nematodes*. Pitman Publishing, Budapest, London, San Francisco, 288 pp.
- Andrásy I. 2005. *Free Living Nematodes of Hungary*. Vol. 1: Nematoda errantia. Hungarian Natural History Museum, Budapest, 495 pp.
- Atighi M.R., Pourjam E., Pedram M., Ye W., Robbins R.T. 2012. Molecular and morphological characterization of *Ektaphelenchoides kelardashtensis* sp. n. (Nematoda: Ektaphelenchinae) from northern Iran. *Russian Journal of Nematology* 21: 23–30.
- Campos-Herrera R., Půža V., Jaffuel G., Blanco-Pérez R., Čepulyte-Rakauskienė R., Turlings T. 2015. Unraveling the intraguild competition between *Oscheius* spp. nematodes and entomopathogenic nematodes: implications for their natural distribution in Swiss agricultural soils. *Journal of Invertebrate Pathology* 132: 216–227. DOI: <https://doi.org/10.1016/j.jip.2015.10.007>
- Castresana J. 2000. Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis. *Molecular Biology and Evolution* 17 (4): 540–552. DOI: <https://doi.org/10.1093/oxfordjournals.molbev.a026334>
- Darsouei R., Karimi J., Shokoohi E. 2014. *Oscheius rugaoensis* and *Pristionchus maupasi*, two new records of entomophilic nematodes from Iran. *Russian Journal of Nematology* 22 (2): 141–55.
- De Grisse A.T. 1969. Redescription ou modifications de quelques techniques utilisées dans l'étude des nematodes phytoparasitaires. [Redescription or modification of some techniques used in the study of phytoparasitic nematodes]. *Mededelingen Faculteit Landbouwwetenschappen Rijksuniversiteit Gent* 34: 351–369. (in French)

- Drummond A.J., Rambaut A. 2007. BEAST: Bayesian evolutionary analysis by sampling trees. *BMC Evolutionary Biology* 7, 214 pp. DOI: <https://doi.org/10.1186/1471-2148-7-214>
- Dichtel-Danjoy M.L., Félix M.A. 2004a. Phenotypic neighborhood and micro-evolvability. *Trends in Genetics* 20 (5): 268–276. DOI: <https://doi.org/10.1016/j.tig.2004.03.010>
- Dichtel-Danjoy M.L., Félix M.A. 2004b. The two steps of vulval induction in *Oscheius tipulae* CEW1 recruit common regulators including a MEK kinase. *Developmental Biology* 265 (1): 113–126. DOI: <https://doi.org/10.1016/j.ydbio.2003.09.010>
- Edgar R.C. 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Research* 32 (5): 1792–1797. DOI: <https://doi.org/10.1093/nar/gkh340>
- Félix M.A., Vierstraete A., Vanfleteren J. 2001. Three biological species closely related to *Rhabditis (Oscheius) pseudodolichura* Körner in Osche, 1952. *Journal of Nematology* 33 (2–3): 104–109. PMID: PMC2638136
- Ghaemi R., Pourjam E., Álvarez-Ortega S., Pedram M., Atighi M.R. 2015. A curious nematode, *Diploscaeteroides persicus* n. sp. (Nematoda: Rhabditida: Rhabditomorpha) from Iran with note on status of male genital papillae. *Nematology* 17 (9): 1045–1056. DOI: <https://doi.org/10.1163/15685411-00002922>
- Hassani-Kakhki M., Karimi J., Hosseini M., Heydari Sh. 2012. Efficacy of entomopathogenic nematodes against potato tuber moth, *Phthorimaea operculella* Zeller (Lep.: Gelechiidae). 20th Iranian Plant Protection Congress, Shiraz University, Shiraz, Iran, 25–28 August 2012, 987 pp.
- Huson D.H., Scornavacca C. 2012. Dendroscope 3: an interactive tool for rooted phylogenetic trees and networks. *Systematic Biology* 61 (6): 1061–1067. DOI: <https://doi.org/10.1093/sysbio/sys062>
- Joyce S.A., Reid A., Driver F., Curran J. 1994. Application of polymerase chain reaction (PCR) methods to identification of entomopathogenic nematodes. p. 178–187. In: “Biotechnology: Genetics of Entomopathogenic Nematode–Bacterium Complexes” (A.M. Burnell, R.U. Ehlers, J.P. Masson, eds.). Proceedings of Symposium and Workshop, St. Patrick's College, Maynooth, Co. Kildare, Ireland, Luxembourg, European Commission, DG XII.
- Lam A.B.Q., Webster J.M. 1971. Morphology and biology of *Panagrolaimus tipulae* n. sp. (Panagrolaimidae) and *Rhabditis (Rhabditella) tipulae* n. sp. (Rhabditidae), from leatherjacket larvae, *Tipula paludosa* (Diptera: Tipulidae). *Nematologica* 17 (2): 201–212. DOI: <https://doi.org/10.1163/187529271x00044>
- Larget B., Simon D.L. 1999. Markov chain Monte Carlo algorithms for the Bayesian analysis of phylogenetic trees. *Molecular Biology and Evolution* 16 (6): 750–759. DOI: <https://doi.org/10.1093/oxfordjournals.molbev.a026160>
- Liu Q.Z., Mráčĕk Z., Zhang L.J., Půža V., Dong L.M. 2012. Re-description of *Oscheius chongmingensis* (Nematoda: Rhabditidae) and its entomopathogenicity. *Nematology* 14 (2): 139–149. DOI: <https://doi.org/10.1163/138855411x580777>
- Miraeiz E., Heydari R., Álvarez-Ortega S., Pedram M., Atighi M.R. 2015. Molecular and morphological characterization of *Veleshkinema iranicum* n. gen., n. sp. (Nematoda: Hexatylinea, Sphaerularioidea) from Iran. *Zootaxa* 4000 (5): 531–546.
- Nunn G.B. 1992. Nematode molecular evolution. An investigation of evolutionary patterns among nematodes based on DNA sequences. Ph.D. thesis, Nottingham, University of Nottingham, UK.
- Nylander J.A. 2004. MrModeltest v2. Evolutionary Biology Centre, Uppsala University. Available on: <https://github.com/nylander/MrModeltest2>. Accessed [March 15, 2015]
- Pedram M., Pourjam E., Ye W., Atighi M.R., Robbins R.T., Rysse A. 2011. Description of *Bursaphelenchus mazandaranense* sp. n. (Nematoda: Parasitaphelenchidae) from Iran. *Russian Journal of Nematology* 19: 121–129.
- Pedram M., Pourjam E., Atighi M.R., Ye W., Houshmand A. 2012. Description of *Ektaphelenchoides sylvestris* sp. nov. (Nematoda: Ektaphelenchinae) from Iran. *Annales Zoologici* 62 (2): 325–329. DOI: <https://doi.org/10.3161/000345412X652864>
- Pedram M., Pourjam E., Atighi M.R., Panahandeh Y. 2015. Further studies on soil nematode fauna in north western Iran with the description of one new species. *Journal of Nematology* 47 (2): 148–152. PMID: PMC4492290
- Shahabi S., Kheiri A., Rakhshandehroo F., Jamali S. 2016. Occurrence and distribution of nematodes in rice fields in Guilan province, Iran and the first record of *Mylonchulus polonicus* (Stefanski, 1915) Cobb, 1917 (Nematoda: Mononchina). *Journal of Plant Protection Research* 56 (4): 420–429. DOI: <https://doi.org/10.1515/jppr-2016-0062>
- Ronquist F., Huelsenbeck J.P. 2003. MrBAYES 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19 (12): 1572–1574. DOI: <https://doi.org/10.1093/bioinformatics/btg180>
- Smart G.C. Jr, Nguyen K.B. 1994. *Rhabditis (Oscheius) pheropsophi* n. sp. (Rhabditida: Rhabditidae). *Journal of Nematology* 26 (1): 19–24. PMID: PMC2619479
- Soleymanzadeh M., Pedram M., Pourjam E., Álvarez-Ortega S. 2016. Description of *Lelenchus brevislitus* n. sp. (Nematoda: Tylenchidae), an example of a cryptic species from Iran and its phylogenetic relationships with other species in the family. *Nematology* 18 (8): 987–998. DOI: <https://doi.org/10.1163/15685411-00003010>
- Sommer R.J. 2000. Comparative genetics: A third model nematode species. *Current Biology* 10 (23): 879–881. DOI: [https://doi.org/10.1016/S0960-9822\(00\)00815-0](https://doi.org/10.1016/S0960-9822(00)00815-0)
- Sudhaus W. 1976. Vergleichende Untersuchungen zur Phylogenie, Systematik, Ökologie, Biologie und Ethologie der Rhabditidae (Nematoda). [Comparative studies on the phylogeny, systematics, ecology, biology and ethology of the Rhabditidae (Nematoda)]. *Zoologica* 43: 1–228. (in German)
- Sudhaus W. 2011. Phylogenetic systemization and catalogue of paraphyletic “Rhabditidae” (Secernentea: Nematoda). *Journal of Nematode Morphology and Systematics* 14: 113–178.
- Sudhaus W. 2016. The guild of saprobiontic nematodes associated with ants. *Ecologica Montenegrina* 7: 600–613.
- Sudhaus W., Hooper D.J. 1994. *Rhabditis (Oscheius) guentheri* sp. n., an unusual species with reduced posterior ovary, with observations on the *Dolichura* and *Insectivora* groups (Nematoda: Rhabditidae). *Nematologica* 40 (1): 508–533. DOI: <https://doi.org/10.1163/003525994x00391>
- Sudhaus W., Schulte F. 1989. *Rhabditis (Rhabditis) necromena* sp. n. (Nematoda: Rhabditidae) from South Australian Diplopoda with notes on its siblings *R. myriophila* Poinar, 1986 and *R. caulleryi* Maupas, 1919. *Nematologica* 35 (1): 15–24. DOI: <https://doi.org/10.1163/002825989x00025>
- Tabassum K.A., Shahina F., Nasira K., Erum Y.I. 2016. Description of six new species of *Oscheius* Andrassy, 1976 (Nematoda: Rhabditida) from Pakistan with a key and diagnostic compendium to species of the genus. *Pakistan Journal of Nematology* 34 (2): 109–161. DOI: <https://doi.org/10.18681/pjn.v34.i02.p109>
- Tamura K., Stecher G., Peterson D., Filipowski A., Kumar S. 2013. MEGA6: molecular evolutionary genetics analysis version 6.0. *Molecular Biology and Evolution* 30 (12): 2725–2729. DOI: <https://doi.org/10.1093/molbev/mst197>
- Torrini G., Mazza G., Carletti B., Benvenuti C., Roversi P.F., Fanelli E., De Luca F., Troccoli A., Tarasco E. 2015. *Oscheius onirici* sp. n. (Nematoda: Rhabditidae): a new entomopathogenic nematode from an Italian cave. *Zootaxa* 3937 (3): 533–548. DOI: <https://doi.org/10.11646/zootaxa.3937.3.6>
- Torrini G., Mazza G., Strangi A., Barabaschi D., Landi S., Mori E., Menchetti M., Sposimo P., Giuliani C., Zoccola A., Lazzaro L., Ferretti G., Foggi B., Roversi P.F. 2016. *Oscheius*

- tipulae* in Italy: evidence of an alien isolate in the Integral Natural Reserve of Montecristo Island (Tuscany). *Journal of Nematology* 48 (1): 8–13. DOI: <https://doi.org/10.21307/jofnem-2017-003>
- Whitehead A.G., Hemming J.R. 1965. A comparison of some quantitative methods of extracting small vermiform nematodes from soil. *Annals of Applied Biology* 55 (1): 25–38. DOI: <https://doi.org/10.1111/j.1744-7348.1965.tb07864.x>
- Williams B.D., Schrank B., Huynh C., Shownkeen R., Waterston R.H. 1992. A genetic mapping system in *Caenorhabditis elegans* based on polymorphic sequence-tagged sites. *Genetics* 131 (3): 609–624. PMID: PMC1205034
- Wouts W., Mráček M.Z., Gerdin S., Bedding R.A. 1982. *Neoplectana* Steiner, 1929 a junior synonym of *Steinernema* Travassos, 1927 (Nematoda: Rhabditida). *Systematic Parasitology* 4 (2): 147–154. DOI: <https://doi.org/10.1007/bf00018998>
- Yaghoubi A., Pourjam E., Pedram M., Siddiqi M.R., Atighi M.R. 2014a. Molecular and morphological characterization of *Abursanema iranicum* n. gen., n. sp. (Nematoda: Hexatylna, Sphaerularioidea) from Iran. *Zootaxa* 3826 (2): 301–314. DOI: <https://doi.org/10.11646/zootaxa.3826.2.1>
- Yaghoubi A., Pourjam E., Atighi M.R., Pedram M. 2014b. Molecular and morphological characterization of *Ektaphelenchoides ruehmi* sp. n. (Nematoda: Ektaphelenchinae) from south-western Iran. *Russian Journal of Nematology* 22: 31–38.
- Zhang K.Y., Liu X.H., Tan J., Wang Y., Qiao L., Yedid G., Dai C.S., Qiu R.L., Yan X.W., Tan H.W., Su Z.Y., Lai R., Gao G.F. 2012. *Heterorhabditoides rugaensis* n. sp. (Rhabditida: Rhabditidae), a novel highly pathogenic entomopathogenic nematode member of Rhabditidae. *Journal of Nematology* 44: 348–360.
- Zheng J., Subbotin S.A., Waeyenberge L., Moens M. 2000. Molecular characterisation of Chinese *Heterodera glycines* and *H. avenae* populations based on RFLPs and sequences of rDNA-ITS regions. *Russian Journal of Nematology* 8: 109–113.