

ORIGINAL ARTICLE

Effects of wheat-canola intercropping on *Phelipanche aegyptiaca* parasitism

Zeynab Razavifar, Hassan Karimmojeni*, Fatemeh Ghorbani Sini

Department of Agronomy and Plant Breeding, College of Agriculture, Isfahan University of Technology, Isfahan, 84156-83111, Iran

Vol. 57, No. 3: 268–274, 2017

DOI: 10.1515/jppr-2017-0038

Received: April 23, 2017

Accepted: August 21, 2017

*Corresponding address:
kmojeni@cc.iut.ac.ir

Abstract

Parasitic weeds especially *Phelipanche aegyptiaca* decrease severely the production of canola. This study evaluated the effect of intercropping different wheat genotypes with canola on *Phelipanche aegyptiaca* growth. Ten wild wheat genotypes with different ploidy levels including TRI11712, TRI19322, TRI18664, TRI19652, TRI565, TRI15593, TRI12911, TRI11554, TRI17606, TRI7259P and seven cultivated bread wheats, namely: Falat, Chamran, Alamut, Baiat, Kavir, Sepahan, Alvand in addition to a canola cultivar called Zarfam were studied. The results revealed that intercropping of canola with wheat could significantly reduce broomrape growth depending on the type of wheat genotype. A significant genetic variation of allelopathic activity in wheat was observed, indicating the contribution of multiple genes conferring the allelopathic trait. TRI565 and TRI12911, TRI15593, TRI18664, TRI19652, TRI17606, TRI19322, and TRI7259 genotypes showed strong inhibitory effects and can be considered as potential allelopathic genotypes to suppress broomrape. The inhibitory potential of wild wheat genotypes was stronger than cultivated wheat genotypes. Alamut, Baiat, Alvand, Sepahan, and TRI11712 possessed strong stimulatory effects on broomrape germination. Such genotypes may be valuable as trap crops for depleting the Egyptian broomrape seed bank.

Key words: allelopathy, intercropping, parasitic plants, *Phelipanche aegyptiaca*, *Striga*, wheat-canola

Introduction

Canola (*Brassica napus* L.) is an annual or biennial herbaceous plant in the Brassicaceae family. This species has such a long history of cultivation and diversification that its origin center is not known. It may originate from a garden hybrid of *B. olearacea* var. *capitata* and *B. rapa* var. *rapa* (Quiros and Paterson 2004).

Since parasitic plants are not able to synthesize adequate nutrients for their development, they require a host plant to survive. Parasitic plants cause severe damage to economically important plants (Mokhtar *et al.* 2009; Blagojević *et al.* 2014).

Broomrape species (*Orobanche* spp. syn. *Phelipanche* spp.) are obligate root parasitic plants which are devoid of chlorophyll (Gauthier *et al.* 2012). Some are noxious parasitic weeds in important crops, including *O. crenata*, *O. cumana*, *O. minor*, *P. aegyptiaca* and *P. ramosa* (Amri 2013).

Phelipanche aegyptiaca poses a serious threat to host plants such as canola, carrot, lettuce, tomato, capeweed and vetch (Babaei *et al.* 2010). Parasitic weeds especially *P. aegyptiaca* decrease severely the production of canola.

Several approaches have been applied to control broomrapes, but none has enjoyed unequivocal success (Gauthier *et al.* 2012). Since the greatest damage caused by these parasites appears prior to their shoot emergence and flowering, the majority of yield loss may occur before the infection diagnosis (Babaei *et al.* 2010). In addition, these weeds produce thousands of minute seeds which are highly persistent in the soil and can easily be distributed to new areas (Acsoy *et al.* 2013). Integrated weed management is the only efficient method for broomrape control (Chittapur *et al.* 2001).

Crop rotation may have direct and indirect effects on parasitic weeds in infested areas. Although trap and catch-crops in rotation may decrease the parasitic seed bank in the soil, other rotation crops may have allelopathic impacts on the seeds of parasitic weeds. Trap cropping is used to protect the main crop from the impact of parasitic weeds (Rubiales and Fernández-Aparicio 2012). It is necessary to identify potential trap crops for each species (Fernández-Aparicio *et al.* 2009).

Some crops have been known to have allelopathic effects. Bread wheat (*Triticum aestivum* L.), durum wheat (*T. durum* L.), barley (*Hordeum vulgare* L.), oats (*Avena sativa* L.), maize (*Zea mays* L.), beet (*Beta vulgaris* L.), lupine (*Lupinus lutes* L.) (Rice 1984; Oueslati 2002) have been examined for differential allelopathy in sustainable weed management strategies.

Intercropping systems may demonstrate advantages for weed suppression over sole cropping in two ways. First, higher crop yield and less weed growth could be achieved if intercropping is more efficient than sole cropping in usurping resources from weeds or suppressing weed growth via allelopathy. Weed suppression in intercropping via more effective use of environmental resources by component crops has been reported in many studies (Liebman and Dyck 1993; Oleszek 1994; Mashingaizde *et al.* 2000; Wanic *et al.* 2004; Poggio 2005; Eskandari and Kazemi 2011). Second, intercropping may provide greater crop yield without weed suppression below levels observed in component sole cropping if intercropping uses resources which are not exploitable by weeds or convert resources to harvestable material more effectively than sole cropping systems (Liebman and Dyck 1993; Eskandari and Kazemi 2011).

Parker and Riches (1993) reported that the effects of intercropping on the control of *Striga* is due to the fact that the intercropped non-host legumes may act as trap crops, and stimulate suicidal *Striga* germination. Abebe *et al.* (2005) evaluated the effect of ten potential trap crops on the *Orobanchae* soil seed bank and tomato yield. They reported that maize and snap bean showed better performance in stimulating the germination of *Orobanchae* seed. Maize and snap bean also complement each other under intercropping and tomato yield increased due to the decrease of the *Orobanchae* seed bank. According to Babaei *et al.* (2010), the soil seed bank of broomrape could be decreased by using trap crops such as sesame, brown Indian hemp, common flax and black-eyed pea in rotation. Ghotbi *et al.* (2012) determined the allelopathic potential of different crops for their abilities to stimulate *P. aegyptiaca* seed germination. They reported corn, oat, beet, sugar beet, triticale, castor-oil plant, millet, fiber flax, pepper, cotton and sorghum as potential trap crops for the weed *P. aegyptiaca*. Baumann *et al.* (2000) suggested that leek could be intercropped with celery (*Apium*

graveolens L.) to improve weed suppression relative to a leek monoculture, whose open canopy structure allows weeds to proliferate. They reported that, for an intercrop of leek and celery, light interception and soil cover were increased compared with a leek monoculture, and yield reduction due to weed competition was decreased. Oswald *et al.* (2002) examined the effect of eight different intercrops on *Striga* populations and crop yield. They found that intercropping is an effective component of an integrated *Striga* control program. In the long run, *Striga* populations can only be reduced if intercropping is combined with hand weeding of mature *Striga* plants to avoid the replenishment of the *Striga* seed bank in the soil.

The objective of this study was to examine the effect of intercropping of different wheat genotypes with canola on reducing *P. aegyptiaca* damage.

Materials and Methods

Plant materials

The present study was performed on ten wild wheat genotypes with different ploidy levels (TRI11712, TRI19322, TRI18664, TRI19652, TRI565, TRI15593, TRI12911, TRI11554, TRI17606, TRI7259P) and seven cultivated bread wheats, including: Falat, Chamran, Alamut, Baiat, Kavir, Sepahan, Alvand in addition to a canola cultivar called Zarfam. Details of the genetic materials are presented in Table 1.

Table 1. Plant materials (wild and cultivated wheats) used in the study

No.	Genotype	Chromosome No.	Code
1	Alvand	42	TRI.1.C.Alv
2	Alamut	42	TRI.2.C.Alm
3	Baiat	42	TRI.3.C.Bi
4	Chamran	42	TRI.4.C.Ch
5	Sepahan	42	TRI.5.C.S
6	Falat	42	TRI.6.C.F
7	Kavir	42	TRI.7.C.C
8	TRI7259	56	TRI.8.W.7259
9	TRI11712	42	TRI.9.W.11712
10	TRI11554	42	TRI.10.W.11554
11	TRI18664	42	TRI.11.W.18664
12	TRI19322	42	TRI.12.W.19322
13	TRI12911	28	TRI.13.W.12911
14	TRI17606	28	TRI.14.W.17606
15	TRI15593	28	TRI.15.W.15593
16	TRI19652	28	TRI.16.W.19652
17	TRI565	14	TRI.17.W.565

Growth conditions and measurements

Broomrape (*P. aegyptiaca*) seeds were collected from naturally infested canola fields. Seeds of broomrape were treated with GA3 (Merck) to break the dormancy. Each pot was contaminated at the rate of 25 mg of broomrape seeds per pot. Pots (20-cm-diameter) were filled 2/3 with soil-broomrape mixture while the remaining 1/3 was filled with clay and vermicompost. In the experimental group, 15 seeds of wheat and 10 seeds of canola were planted per pot and were later thinned down to 2 wheat seedlings and 1 canola seedling per pot at two-leaf stage for wheat and at four-leaf stage for canola. In the control group, only canola seeds were planted. Pots were watered every other day as needed. The plants were grown at a day/night cycle of 16/8 h, at 25/20°C, and at a light intensity of 12,000 LUX. Plants were fertilized with NPK fertilizer [20–20–20 + TE (trace element), 1 g · l⁻¹] several times to avoid any nutrient deficit during the growth period. The number of broomrape stems and tubercles, the length and dry weight of the underground parts of the broomrape, and the dry weight of aerial parts were recorded at the ripening stage of canola.

Statistical analysis

The experiment was conducted in the form of a completely randomized design with 3 replications and 18 treatments. All collected data were subjected to analysis of variance (ANOVA) by SAS Ver.9.1 and mean comparisons were performed using LSD. For grouping the genotypes all data were analyzed using cluster analysis with the JMP statistic program after data standardization by the Ward method.

Results

Results of ANOVA indicated that wheat-canola intercropping significantly ($\alpha = 0.01$) affected measured traits of broomrape (Table 2).

Effect of wheat-canola intercropping on the growth of broomrape

Number of broomrape stems

The control treatment produced 4.66 broomrape stems, while the highest number of broomrape stems at the soil surface was found in Alvand-canola intercropping, which was 7.33 (Table 3). Alvand, Sepahan, Alamut, Baiat, and TRI11712 were categorized in the same statistical group in terms of this trait. The least number of stems was observed in Chamran-canola intercropping (3.66), which was classified in the same group as Falat, TRI18664 and TRI15593. The Chamran genotype had

the greatest inhibitory effect on this trait. No broomrape stems were found at the soil surface of TRI12911- and TRI 565-canola intercropping. Hence, these two genotypes had strong inhibitory effects.

Number of broomrape tubercles

Control treatment produced 2.66 broomrape stems, while the maximum number of broomrape tubercles was observed in Baiat-canola intercropping (4.66). This genotype was classified in the same statistical group as: Alamut, Sepahan, Alvand, TRI11712 and TRI11554 in terms of the number of tubercles. Broomrape produces more tubercles in the presence of some genotypes which release more chemicals with the ability to stimulate broomrape germination. The greater number of broomrape tubercles means further growth of broomrape and more attachments of this species to near-by canola. The minimum number of tubercles was found in TRI19322-canola intercropping. TRI19322, Falat, TRI17606, TRI7259, TRI15593, and TRI18664 were classified in the same statistical group. No tubercles were detected in TRI565- and TRI12911-intercropping (Table 3). Thus, broomrape germination was completely inhibited by these two genotypes.

Length of broomrape underground parts

The length of broomrape underground parts was (5.53 cm) in sole culture of canola (control treatment). The highest length of broomrape underground parts (7.26 cm) was found when Sepahan genotype was intercropped with canola, and the lowest broomrape root length (1.43 cm) occurred in TRI7259-canola intercropping (Table 3).

Dry weight of broomrape underground parts

While intercropping Baiat genotype with canola led to the highest dry weight of broomrape underground parts (0.673 g · pot⁻¹), TRI15593 caused the lowest amount (0.113 g · pot⁻¹). TRI15593, TRI7259, and TRI19652 were placed in the same statistical group in terms of this trait (Table 3). The dry weight of broomrape underground parts was (0.346 g · pot⁻¹) in sole culture of canola.

Dry weight of broomrape aerial parts

The highest and the lowest dry weights of broomrape aerial parts were produced by Alvand- and Chamran-canola intercropping, which were 0.626 and 0.21 (g · pot⁻¹), respectively. Chamran, TRI7259, TRI15593, and TRI19322 genotypes were classified in the same statistical group in terms of this trait (Table 3).

Orthogonal contrasts

According to orthogonal contrasts results (Table 3), there were no significant differences between wild and

Table 2. Analysis of variance results for effects of wheat-canola intercropping on broomrape traits. Mean square values are presented for each variable

Source of variation	Degree of freedom	Stem No.	Tubercle No.	Length of underground part	Dry weight of underground part	Dry weight of aerial part
Treatment	15	7.088**	3.22**	7.86**	0.0749**	0.0476**
Comparison of wild wheats	6	–	–	–	–	–
Comparison of cultivated wheats	7	–	–	–	–	–
Cultivated wheats compared to wild wheats	1	6.816 ns	5.08*	26.821**	0.000054 ns	0.0138 ns
Error	32	0.645	0.458	0.048	0.0025	0.00075
CV%	–	16.91	22.41	4.716	14.33	7.62

*significant at 5% probability level, **significant at the 1% probability level, ns – not significant, CV – coefficient of variance

Table 3. Effect of wheat-canola intercropping on broomrape traits

Wild and cultivated wheats	Genotype	Stem No.	Tubercle No.	Length of underground part [cm · pot ⁻¹]	Dry weight of underground part [cm · pot ⁻¹]	Dry weight of aerial part [g · pot ⁻¹]
Cultivated wheats	TRI.1.C.Alv	7.33 a	4 ab	6 c	0.536 b	0.626 a
	TRI.2.C.AIm	6.66 a	4.33 ab	6.13 c	0.41 cd	0.52 b
	TRI.3.C.Bi	6 ab	4.66 a	6.3 c	0.673 a	0.453 c
	TRI.4.C.Ch	3.66 cde	3.33 bcd	5.26 d	0.486 bc	0.21 i
	TRI.5.C.S	7 a	4.33 ab	7.26 a	0.470 bc	0.556 b
	TRI.6.C.F	3 de	2.33 def	6.83 b	0.32 ef	0.293 f
	TRI.7.C.C	4.66 bc	2.66 cde	4.5 e	0.283 ef	0.28 fg
Cultivated wheats mean		5.47 a	4.66 a	6.08 a	0.452 a	0.4 a
Wild wheats	TRI.8.W.7259	2.33 e	2 ef	1.43 i	0.123 h	0.233 hi
	TRI.9.W.11712	6.66 a	4 ab	4.06 f	0.536 b	0.383 de
	TRI.10.W.11554	4.33 cd	3.66 abc	4.63 e	0.243 fg	0.26 fgh
	TRI.11.W.18664	3.33 cde	2.33 def	3.56 g	0.236 fg	0.386 de
	TRI.12.W.19322	4.33 cd	1.33 f	3.96 f	0.346 de	0.246 ghi
	TRI.13.W.12911	–	–	–	–	–
	TRI.14.W.17606	4.33 cd	1.66 ef	3.03 h	0.333 de	0.303 f
	TRI.15.W.15593	3.33 cde	2.33 def	3.2 h	0.113 h	0.243 ghi
	TRI.16.W.19652	4.33 cd	2.66 cde	3 h	0.186 gh	0.423 cd
	TRI.17.W.565	–	–	–	–	–
Wild wheats mean		4.12 a	2.49 b	3.35 b	0.45 a	0.339 a
Control	Can.18.ZA	4.66 bc	2.66 cde	5.53 d	0.346 de	0.353 e

In each column, means which have similar letters do not have significant difference based on LSD test

cultivated wheat genotypes in terms of the number of broomrape stems at the soil surface, the dry weight of broomrape underground parts, and the dry weight of broomrape aerial parts. However, significant differences were detected between them in terms of the number

of broomrape tubercles and the length of broomrape underground parts at 1% and 5% probability levels, respectively. In general, the inhibitory potential of wild wheat genotypes was stronger than cultivated wheat genotypes (Table 3).




Fig. 1. Cluster analysis of wheat genotypes based on their allelopathic effects on measured traits of broomrape in wheat-canola intercropping

Classification of wheat genotypes

Cluster analysis divided the genotypes into four groups (Fig. 1): 1 – Alamut, Baiat, Alvand, Sepahan, and TRI11712 with no inhibitory effect on broomrape and with stimulatory effect on most measured traits, 2 – Falat variety with slight stimulatory effect on broomrape, and 3 – this group consisted of 2 subgroups. The 1 subgroup included Zarfam monoculture, Chamran, Kavir, and TRI11554 with no impact on broomrape; and the 2 subgroup contained wild varieties, TRI18664, TRI19652, TRI17606, and TRI19322 with intermediate inhibitory effects, 4 – TRI7259 with a significant inhibitory effect on broomrape traits. No broomrape and tubercles were found in TRI565- and TRI12911-canola intercropping. Hence, these two genotypes are not present in the diagram. According to the results of this study, TRI565 and TRI12911 were genotypes with the highest inhibitory effects against broomrape.

Discussion

The findings indicated that intercropping of canola with wheat could significantly reduce broomrape growth depending on the type of wheat genotype. The results of our research are consistent with Naeem *et al.* (2012), who evaluated intercropping of wheat with canola under varied spatial arrangements for their effects on weeds and reported that all treatments significantly reduced weed density and dry weight more than a sole crop of wheat. Similarly, Liebman and Dyck (1993) found that weed population density and biomass

production may be significantly decreased using crop rotation and intercropping strategies. In another research, Babaei *et al.* (2010) reported that sesame, common flax, and black-eyed pea led to high control of broomrape so that broomrape biomass was reduced by 86, 85.3, 57.2, and 74.4%, respectively, compared to the control (tomato monoculture).

The results confirmed that there was considerable genetic variation in broomrape germination and growth inhibitory activity of wheat germplasm. The results of the current study are broadly in agreement with those of Wu *et al.* (2000, 2001), who also found a significant genetic variation of weed growth inhibitory activity in wheat and reported that wheat allelopathic activity is genetically controlled, and a multi-genetic model is involved in wheat allelopathy.

TRI565 and TRI12911 were identified as the genotypes with strong inhibitory potential to suppress broomrape. Also, TRI15593, TRI18664, TRI19652, TRI17606, TRI19322, and TRI7259 genotypes showed relatively high inhibitory effects. These genotypes can be considered as potential allelopathic genotypes to suppress broomrape. Our results are in agreement with Fernández-Aparicio *et al.* (2007), who demonstrated that intercropping with cereals decreases infection by *O. crenata* in legumes, and allelochemicals released by cereal roots inhibited *O. crenata* seed germination. A limiting effect of intercropping on the number and biomass of weeds has been reported by a number of researchers (Carruthers *et al.* 1998; Poggio 2005; Amanullah *et al.* 2006; Banik *et al.* 2006; Gharineh and Moosavi 2010; Eskandari and Kazemi 2011).

Since Alamut, Baiat, Alvand, Sepahan, and TRI11712 possessed weak inhibitory effects and strong stimulatory

effects on broomrape germination, they can be applied as trap crops in rotation with broomrape hosts. Using trap crops to deplete soil seed banks of parasitic weeds has been reported in several studies (Carson *et al.* 1989; Linke *et al.* 1993; Parker and Riches 1993; Kliefeld *et al.* 1994; Carsky *et al.* 1994; Muscolo *et al.* 2001; Abebe *et al.* 2005; Babaei *et al.* 2010; Ghotbi *et al.* 2012). Most of broomrape traits were reduced more in response to wild wheats compared to cultivated wheats. Hence, it could be concluded that wild wheat genotypes possessed strong inhibitory effects against broomrape. Hence, the results of this study suggest using wild wheat genotypes to suppress weeds in intercropping strategies of canola production.

Considerable genetic variation was found among the studied wheat genotypes in terms of allelopathic potential, which may permit selecting more allelopathic cultivars to be applied against broomrape. Identification of the composition of chemicals released from these genotypes, especially wild wheat genotypes, may provide more opportunities for applying reliable intercropping and rotation systems and new strategies in suppressing this parasitic weed.

Acknowledgements

This work was supported by the research council of Isfahan University of Technology.

References

- Abebe G., Sahile G., Altawaha A.R.M. 2005. Evaluation of potential trap crops on *Orobanche* soil seed bank and tomato yield in the central rift valley of Ethiopia. *World Journal of Agricultural Sciences* 1 (2): 148–151.
- Acsoy E., Filiz Arsalan Z., Öztürk N. 2013. *Phelipanche aegyptiaca* (Pers.) Pomel: A new record as a parasitic weed on apricot root in Turkey. *African Journal of Agricultural Research* 8 (29): 4002–4006. DOI: 10.5897/AJAR2013.6639
- Amanullah M.M., Alagesan A., Vaiyapui K., Sathyamoorthi K., Pazhanivelan S. 2006. Effect of intercropping and organic manures on weed control and performance of cassava (*Manihot esculenta* Cranz). *Journal of Agronomy* 5 (4): 589–595. DOI: <https://doi.org/10.3923/ja.2006.589.594>
- Amri M. 2013. First report of the parasitic plant *Phelipanche ramosa* on berseem clover (*Trifolium alexandrinum*) in Tunisia. *Tunisian Journal of Plant Protection* 8: 127–132.
- Babaei S., Alizadeh H., Jahansouz R., Rahimian H., Minbashi Moeini M. 2010. Management of *Phelipanche aegyptiaca* Pomel. using trap crops in rotation with tomato (*Solanum lycopersicom* L.). *Australian Journal of Crop Science* 4 (6): 437–442.
- Banik P., Midya A., Sarkar B.K., Ghose S.S. 2006. Wheat and chickpea intercropping systems in an additive experiment: Advantages and weed smothering. *European Journal of Agronomy* 24 (4): 325–332. DOI: <https://doi.org/10.1016/j.eja.2005.10.010>
- Blagojević M., Konstantinović B., Samardžić N., Popov M., Konstantinović B. 2014. Biological characteristics of some parasitic flowering plants. *Herbologia* 14 (2): 71–80.
- Baumann D.T., Kropff M.J., Bastiaans L. 2000. Intercropping leeks to suppress weeds. *Weed Research* 40 (4): 361–376. DOI: <https://doi.org/10.1046/j.1365-3180.2000.00197.x>
- Carruthers K., Fe Q., Cloutier D., Smith D.L. 1998. Intercropping corn with soybean, lupin and forages: Weed control by intercrops combined with inter row cultivation. *European Journal of Agronomy* 8 (3–4): 225–238. DOI: [https://doi.org/10.1016/s1161-0301\(97\)00062-2](https://doi.org/10.1016/s1161-0301(97)00062-2)
- Carson A.G. 1989. Effect of intercropping sorghum and groundnuts on density of *Striga hermonthica* in the Gambia. *Tropical Pest Management* 35 (2): 130–132. DOI: <https://doi.org/10.1080/09670878909371340>
- Chittapur B.M., Hunshal C.S., Shenoy H. 2001. Allelopathy in parasitic weeds management: Role of catch and trap crops. *Allelopathy Journal* 8: 147–160.
- Carsky R.J., Singh L. and Ndikawa R. 1994. Suppression of *Striga hermonthica* on sorghum using a cowpea intercrop. *Experimental Agriculture* 30 (03): 349–358. DOI: <https://doi.org/10.1017/s0014479700024467>
- Eskandari H., Kazemi K. 2011. Weed control in maize-cowpea intercropping system related to environmental resources consumption. *Notulae Scientia Biologicae* 3 (1): 57–60. DOI: <http://dx.doi.org/10.15835/nsb315609>
- Fernández-Aparicio M., Sillero J.C., Rubiales D. 2007. Intercropping with cereals reduces infection by *Orobanche crenata* in legumes. *Crop Protection* 26 (8): 1166–1172. DOI: <https://doi.org/10.1016/j.cropro.2006.10.012>
- Fernández-Aparicio M., Flores F., Rubiales D. 2009. Recognition of root exudates by seeds of broomrape (*Orobanche* and *Phelipanche*) species. *Annals of Botany* 103 (3): 423–431. DOI: <https://doi.org/10.1093/aob/mcn236>
- Gauthier M., Véronési C., El.Halmouch Y., Leflon M., Jestin C., Labalette F., Simier P., Delourme R., Delavault P. 2012. Characterisation of resistance to branched broomrape, *Phelipanche ramosa*, in winter oilseed rape. *Crop Protection* 42: 56–63. DOI: <https://doi.org/10.1016/j.cropro.2012.07.002>
- Gharineh M.H., Moosavi S.A. 2010. Effects of intercropping (canola-faba bean) on density and diversity of weeds. *Notulae Scientia Biologicae* 2 (4): 109–112. DOI: <http://dx.doi.org/10.15835/nsb213612>
- Ghotbi M., Rouhi H.R., Amini Dehaghi M., Ghotbi M., Moghaddam Khamseh A.R. 2012. Mitigate *Phelipanche aegyptiaca* Pers. infestation considering natural environment conservation. *International Journal of AgriScience* 2 (1): 62–77.
- Kliefeld Y., Goldwasser Y., Herzlinger G., Joel D.M., Golan S., Kahana D. 1994. The effect of flax (*Linum usitatissimum* L.) and other crops as trap and catch crops for control of Egyptian broomrape (*Orobanche aegyptiaca* Pers.). *Weed Research* 34 (1): 37–44. DOI: <https://doi.org/10.1111/j.1365-3180.1994.tb01971.x>
- Liebman M., Dyck E. 1993. Crop rotation and intercropping strategies for weed management. *Ecological Applications* 3 (1): 92–122. DOI: <https://doi.org/10.2307/1941795>
- Linke K.H., AbdelMoneim A.M., Saxena M.C. 1993. Variation in resistance of some forage legume species to *Orobanche crenata* Forsk. *Field Crops Research* 32 (3–4): 277–285. DOI: [https://doi.org/10.1016/0378-4290\(93\)90037-n](https://doi.org/10.1016/0378-4290(93)90037-n)
- Mashingaizda A.B., Nyakanda C., Chivinge O.A., Washaireni A.M., Dube K.W. 2000. Influence of a maize pumpkin live mulch on weed dynamics and maize yield. *African Plant Protection* 6 (1): 57–63.
- Mokhtar M., Abdel K., Nehal S.E.M. 2009. Prospects of mycoherbicides for control of broomrapes (*Orobanche* spp.) in Egypt. *Journal Plant Protection Research* 49 (1): 64–74. DOI: <https://doi.org/10.2478/v10045-009-0009-1>
- Muscolo A., Panuccio M.R., Sidari M. 2001. The effect of phenols on respiratory enzymes in seed germination. Respiratory enzyme activities during germination of *Pinus laricio* seeds treated with phenols extracted from different forest soils. *Plant Growth Regulation* 35: 31–35.
- Naem M., Zahid A.C., Ahmad A., Abdul W., Kamaran M., Arif M. 2012. Weed dynamics in wheat-canola intercrop-

- ping systems. *Chilean Journal of Agricultural Research* 72 (3): 434–439. DOI: <https://doi.org/10.4067/s0718-58392012000300019>
- Oleszek K. 1994. Brassicacea as alternative plants for weed control in sustainable agriculture. *Fragmenta Agronomica* 4 (44): 5–15.
- Oswald A., Ransom J.K., Kroschel J., Sauerborn J. 2002. Intercropping controls *Striga* in maize based farming systems. *Crop Protection* 21 (6): 367–374. DOI: [https://doi.org/10.1016/s0261-2194\(01\)00104-1](https://doi.org/10.1016/s0261-2194(01)00104-1)
- Oueslati O. 2002. Allelopathy in two durum wheat (*Triticum durum* L.) varieties. *Agriculture Ecosystems & Environment* 96 (1–3): 161–163. DOI: [https://doi.org/10.1016/s0167-8809\(02\)00201-3](https://doi.org/10.1016/s0167-8809(02)00201-3)
- Parker C., Riches C.R. 1993. *Parasitic Weeds of the World: Biology and Control*. CAB International, Wallingford, UK, 332 pp.
- Poggio S.L. 2005. Structure of weed communities occurring in monoculture and intercropping of field pea and barley. *Agriculture Ecosystems & Environment* 109 (1–2): 48–58. DOI: <https://doi.org/10.1016/j.agee.2005.02.019>
- Quiros C.F., Paterson A.H. 2004. Genome mapping and analysis. p. 31–42. In: “Biotechnology in Agriculture and Forestry” (E.C. Pua, C.J. Douglas, eds). Vol. 54. Springer-Verlag, Berlin-Heidelberg-New York. DOI: https://doi.org/10.1007/978-3-662-06164-0_3
- Rice E.L. 1984. *Allelopathy*. 2nd ed. Academic Press, Orlando, FL, USA, 422 pp.
- Rubiales D., Fernández-Aparicio M. 2012. Innovation in parasitic weeds management in legume crops. A review. *Agronomy for Sustainable Development* 32 (2): 433–449. DOI: <https://doi.org/10.1007/s13593-011-0045-x>
- Wanic M., Kostrzevska M., Jastrzebska M. 2004. Role of intercrop sowing in weeds control for spring barley in cereal crops rotation. *Fragmenta Agronomica* 1 (81): 85–102.
- Wu H., Pratley J., Lemerle D., Haig T. 2000. Evaluation of seedling allelopathy in 453 wheat (*Triticum aestivum*) accessions against annual ryegrass (*Lolium rigidum*) by the equal compartment-agar method. *Australian Journal of Agricultural Research* 51 (7): 937–944. DOI: <https://doi.org/10.1071/AR00017>
- Wu H., Pratley J., Lemerle D., Haig T. 2001. Allelopathy in wheat (*Triticum aestivum*). *Annals of Applied Biology* 39 (1): 1–9. DOI: <https://doi.org/10.1111/j.1744-7348.2001.tb00124.x>