

ORIGINAL ARTICLE

Production, characterization and iron binding affinity of hydroxamate siderophores from rhizosphere associated fluorescent *Pseudomonas*

Monali Deori, Nellickal Subramanyan Jayamohan, Belur Satyan Kumudini*

Department of Biotechnology, School of Science, Jain University, Bangalore 18/3, 9th Main, 3rd Block, Jayanagar, Bangalore 560011, India

Vol. 58, No. 1: 36–43, 2018

DOI: 10.24425/119116

Received: September 5, 2017

Accepted: December 12, 2017

*Corresponding address:
kumudini.satyan.ju@gmail.com

Abstract

Fluorescent *Pseudomonas* (FP) is a major group of plant growth promoting rhizobacteria and a well-known synthesizer of siderophores, which imparts a selective advantage on rhizosphere competence and their biocontrol traits. The present study was aimed at examining the factors affecting the production of siderophores and their potential biocontrol traits. Sixteen FP isolates were shortlisted based on their siderophore-producing ability in chrome azural S medium. The isolates were checked for variations in siderophore production under varying incubation times, temperatures, pH, iron (Fe^{3+}) concentrations and mutagens. In addition, the iron binding affinity of siderophores, mycelial inhibition assay and plant growth promotion traits were assessed. Results showed that the siderophore production was highly influenced by the time of incubation, changes in pH, temperature and iron concentration. Chemical characterization showed that the produced siderophores were hydroxamates. Maximum siderophore production was observed at pH 7 whereas UV and EtBr exposure invariably suppressed siderophore production drastically in all isolates. All FPs from maize rhizosphere showed excellent siderophore production which could be due to the competence in strategy-II of the plant rhizosphere and significant growth inhibition on *Fusarium oxysporum*. Our results suggest the inclination of siderophores to iron, in terms of various criteria affecting production and the possible role of environmental mutations that affect the natural iron harvesting mechanism.

Key words: siderophore, hydroxamate, fluorescent *Pseudomonas*, succinate medium, iron-binding, mutation

Introduction

Iron is an essential element of all living microorganisms by serving as a cofactor for various enzymes, with the possible exceptions of *Lactobacilli* and *Borrelia burgdorferi* (Masaki *et al.* 2013; Sah and Singh 2015). Although plentiful, iron does not normally occur in its biologically relevant Fe^{2+} form under aerobic conditions since it is unstable in nature (Krewulak and Vogel 2008; Sah and Singh 2015). Iron availability for microbial assimilation in environments such as the rhizosphere, is limiting (Mishra *et al.* 2008). Microorganisms overcome iron scarcity through two mechanisms: i) iron acquisition by cognate receptors using

low molecular weight iron chelators (siderophores), ii) receptor-mediated iron acquisition from host proteins. Siderophores are non-protein, non-porphyrin compounds synthesised by microbes, where its synthesis is repressed when iron is abundant in the environment (Jenifer *et al.* 2013). The major difference between these two mechanisms is that the siderophores can be taken up by the bacterial cell as intact molecules, whereas prior to transport, iron must be extracted from the host carrier proteins such as transferrin or lactoferrin (Krewulak and Vogel 2008). The mode of action of siderophores in the biocontrol of plant pathogens

appears to be through the induction of systemic resistance, which depends on iron-regulated metabolites (Helmy *et al.* 2008).

More than 500 different siderophores have been identified. Essentially they have the same basic structure, which consists of a ligand that binds to iron and a peptide backbone that interacts with the receptor of bacterial membrane (Lee *et al.* 2011). They are diverse, highly specific and biosynthesized from amino acids by nonribosomal peptide synthetases (NRPS) or from dicarboxylic and diamine compounds in NRPS independent manner (Miethke and Marahiel 2007). Structurally, siderophores are classified as catacholates (phenolates), hydroxamates and carboxylates (Murugappan *et al.* 2011). Secondary siderophores are weakly expressed or not expressed under iron limited conditions (Miethke and Marahiel 2007). Most aerobic microorganisms produce at least one siderophore, and in some cases, a single bacterial strain can produce two or more (Helmy *et al.* 2008).

Extreme environmental conditions are quite common on the earth and the microbial diversity in such areas is of particular interest as microorganisms inhabiting these places are well adapted to the prevailing atmospheric and edaphic conditions (Mishra *et al.* 2008). The secondary alleochemicals produced by plant growth promoting rhizobacteria (PGPR), which include siderophores, antibiotics, mycolytic enzymes, volatile organic metabolites, systemic fungicides, etc. enable them to combat the adverse environmental circumstances (Saraf *et al.* 2014; Khan *et al.* 2016). In the present study an attempt was made to study the various parameters that affect siderophore production in rhizosphere isolated fluorescent *Pseudomonas* (FP) and their binding affinity towards iron under *in vitro* conditions.

Materials and Methods

Isolation, screening and identification of isolates for siderophore production

Plants with their root region soil were carefully removed, placed in sterile plastic bags and gently transferred to the laboratory. The soil adhering to roots was carefully removed, serially diluted and colonies were isolated using King's B medium with 8-HQ supplementation (120 ppm) as per Fallahzadeh-Mamaghani *et al.* (2009).

Pseudomonas colonies which were fluoresced under ultraviolet light ($\lambda_{\max} = 365 \text{ nm}$) were isolated, biochemically characterized, purified and stored at 4°C. Qualitative siderophore production of FPs was analysed using universal chrome azural S (CAS) medium (Alexander and Zuberer 1991).

In order to identify the genus of the shortlisted FPs, the genomic DNA was amplified using *Pseudomonas*-specific 16S primer pair Psmn 289-5'-GGTCGA GAGGATGATCAGT-3' and Psmn 1258 5'-TTAGC TCCACCTCGCGGC-3' (Widmer *et al.* 1998). The polymerase chain reaction (PCR) conditions were optimized according to Varsha and Kumudini (2016).

Chemical characterization of siderophores

CAS-positive isolates were tested for the quantitative siderophore production in succinate medium and quantified at 24 h intervals for 96 h (Gupta *et al.* 2008; Ali and Vidhale 2011). Then, extracted cell free culture supernatants were analysed for hydroxamates by spectrophotometry (Shimadzu UV-1800, Japan). Characterization of siderophore was carried out chemically as per Neilands assay (1981), confirmed by Tetrazolium test (1954) and Arnow's assay (1937).

Effect of incubation time, pH, temperature, Fe³⁺ concentration and mutagenic agents (UV and EtBr) on siderophore production

The culture density of the overnight FP cultures of the isolates was adjusted to $\sim 1 \times 10^8 \text{ CFU} \cdot \text{ml}^{-1}$ ($\text{OD}_{600} = 0.3$). 20 μl of this was inoculated into succinate medium and incubated at 37°C. Cultures were centrifuged at 10,000 rpm for 10 min. Cell free supernatant was diluted with 0.5 M phosphate buffer (pH 7) and absorbance was read at 400 nm.

To evaluate the ability of FPs to produce siderophores at varying pH, isolates were allowed to grow in succinic acid medium for 72 h at different pH (5–9). Siderophore production was analysed every 24 h. To study the influence of different incubation temperatures, isolates were inoculated into succinic acid medium and incubated at four different temperatures (16, 25, 37 and $50 \pm 2^\circ\text{C}$). To determine the effect of iron concentration in siderophore production, isolates were allowed to grow in succinate medium (37°C) supplemented with Fe³⁺ (ferric ammonium citrate, Merck, India) at different concentrations (1, 10 and 50 μM). Siderophore production was recorded at 24 h intervals.

To study the role of mutagens on siderophore production, 24 h old cultures of FPs in Luria Bertani medium (LB) were pelleted by centrifugation (3,000 rpm for 15 min). Fluorescent *Pseudomonas* were treated with three different concentrations of ethidium bromide (EtBr; 0.05, 0.1 and 1.5 $\text{g} \cdot \text{l}^{-1}$) as per Kamal *et al.* (2003) at a cell concentration of $\sim 1 \times 10^8 \text{ CFU} \cdot \text{ml}^{-1}$. They were incubated for 30 min and washed until the remnants of EtBr were completely removed and re-inoculated into succinate medium, incubated for 72 h and checked for siderophore production. Physical

mutagenesis was carried out using ultraviolet (UV) as per Gawel *et al.* (2012) with minor modifications. Bacterial cells were pelleted from overnight LB culture broth and exposed to UV-light ($\lambda_{\max} = 365$ nm) for 30 and 60 s. The pellets were re-inoculated into succinate medium and analysed for siderophore production.

Iron binding affinity of siderophores

72 h old FP cultures were centrifuged and the supernatant was transferred to a separating funnel. An equal volume of chloroform containing 3% (w/v) 8-hydroxy quinolone (8-HQ, Himedia, India) was added and the chloroform layer was separated. Three different concentrations (10, 25 and 50 μ M) of Fe^{3+} (ferric ammonium citrate – Merck, India) were added to the supernatant. Optical density was read ($\lambda_{\max} = 450$ nm) after of 5, 15, 30, 60, 90 and 120 min (Ali and Vidhale 2011).

In vitro antifungal activity and seed germination studies

All the FPs were tested for their *in vitro* antifungal activity by well diffusion assay (Zongzheng *et al.* 2009) against the fungal pathogen *Fusarium oxysporum* (MTCC-1755) and the zone of inhibition was recorded after 7 days.

Tomato seeds (PKM 1 variety) were soaked in 72 h old FP suspension (washed in saline) for 2 h, blot dried and plated on to blotter discs. They were incubated in a growth chamber for 15 days along with distilled water control. Root length and shoot length were measured and percentage germination was calculated.

Statistical analysis

All the experimental analyses were carried out in triplicate. Means were compared using ANOVA with Graph Pad Prism 6.01 version and IBM SPSS statistics 20.

Results and Discussion

Siderophores represent one important class of molecules which solubilize Fe^{3+} from the surrounding environment. In the present study, the siderophore producing ability of a set of shortlisted FPs was carried out. Additionally, the ability of FPs to respond to the common mutagens and their phytoaugmentation traits were analysed.

Isolation, shortlisting and molecular identification of FPs

Comprehensive screening of siderophore showed orange halos around bacterial colonies on CAS agar

Table 1. Shortlisted CAS positive fluorescent *Pseudomonas* isolates and their site of collection

<i>Pseudomonas</i> isolates	Site of collection
P88	<i>Nyctanthes arbour-tristis</i> , Assam
T125, T126	<i>Solanum lycopersicum</i> , Karnataka
M131, M132, M136, M138, M139, M140	<i>Zea mays</i> , Karnataka
R40, R44, R45, R61	<i>Oryza sativum</i> , Bihar
C43	<i>Capsicum annum</i> , Karnataka
J49	<i>Jasminum sambac</i> , Bihar
A55	<i>Aloe vera</i> , Bihar

plates, indicating siderophore production. Based on the qualitative analysis, 16 FPs were shortlisted for further studies (Table 1). It was noticeable that six out of 16 shortlisted isolates were from maize rhizosphere and were excellent producers of siderophores. Previously many authors reported siderophore-producing FPs as biocontrol agents to combat plant pathogens (Beneduzi *et al.* 2012; Gawel *et al.* 2012). Lemanceau *et al.* (2009) reported that in plants with the strategy-II iron-acquisition system, only indirect mechanisms may account for improved plant nutrition by pyoverdines due to their higher affinity towards iron, than phyto siderophores. Moreover, a plant's iron status affects the composition of rhizosphere micro-flora, which is probably due to secretion of phenolics, eventually resulting in elevated siderophore synthesis (Jin *et al.* 2010).

Partial amplification of 16S rRNA of the 16 listed isolates using *Pseudomonas* specific primers showed a 957 bp amplified product in all the isolates including control strains. The molecular weight of the amplified product, confirmed by 1 kb ladder, showed that all isolates belong to genus *Pseudomonas*.

Chemical characterization of siderophores

Succinate medium, following incubation, turns green to yellow as is characteristic of pyoverdins. The presence of siderophores is seen when, with the addition of ferric chloride, it turns red or purple. Spectrophotometric assay results showed a λ_{\max} at 408 nm, which was absent in a de-ferrated condition (Fig. 1). An absorption peak between 360 and 420 nm is characteristic of pyoverdins (Bhattacharya 2010), confirming the presence of pyoverdins. Ferric chloride assay did not show any peak at 495 nm and in 190–280 nm, indicating the absence of catecholates and carboxylates (Baakza *et al.* 2004). Positive results for the tetrazolium salt test confirmed the hydroxamate nature of siderophores. Siderotyping is a very useful analysis which correlates well-defined species and also detects




Fig. 1. Representative depiction of UV-visible spectrum of eluted siderophores by fluorescent *Pseudomonas* isolates. Peak at 408 nm indicates the presence of ferric Fe^{3+}

misclassified individual strains among fluorescent/non-fluorescent *Pseudomonas*. This explains the role of pyoverdine as a powerful taxonomic marker (Meyer *et al.* 2002). The latest study on the peptide side chains of pyoverdine siderophores from *Pseudomonas putida* showed that the structure is unique among all pyoverdins. This shows the strength of the peptide structure for phylogenetic analysis (Ye *et al.* 2014).

A time course study on siderophore production was carried out up to 96 h. Maximum siderophore production was observed after 72 h by M139 ($182.03 \pm 9.79 \mu\text{g} \cdot \text{ml}^{-1}$), followed by T125 ($165.71 \pm 5.26 \mu\text{g} \cdot \text{ml}^{-1}$) as shown in Figure 2. Siderophore production gradually increased till 72 h, followed by a steady decline in all the isolates. Therefore, further experiments were restricted to 72 h incubation.

Several studies on strain-to-strain variations in the production of siderophores have been reported




Fig. 2. A time course assay on production of siderophores by fluorescent *Pseudomonas* isolates in succinic acid medium. Data of the qualitative analyses are represented as mean \pm standard error where $n = 3$. Means were compared between treatments by Duncan multiple-ranges test. Different letters indicate significant differences among treatments ($p \leq 0.05$)

previously (Tailor and Joshi 2012). Nagata *et al.* (2013) reported that tomato plants supplemented with siderophore from *P. fluorescens* ATCC13525 showed increased iron absorption efficiency in comparison to direct iron supplementation. Although pyoverdine-mediated antagonism gives a competitive advantage to the FP (Lemanceau *et al.* 2009), this requires a specific outer membrane receptor, which is advantageous to the organism under iron stress (Nagata *et al.* 2013). Previous reports clearly indicate that *Pseudomonas* can produce siderophores with different chemical natures. This may be due to evolutionary pressures lodged on microbes to produce structurally different siderophores which cannot be transported by other microbial transport systems (Tripathi *et al.* 2005). Synthesis of more than one siderophore and the production of multiple siderophore uptake systems by a single bacterial species are common. The selective advantages conferred by the multiplicity of siderophore synthesis remains unclear. Recent evidence suggests that siderophores may have other physiological roles besides their involvement in iron acquisition (Adler *et al.* 2012).

The role of physical parameters in siderophore production

Studies on the effect of temperature showed reduced siderophore production at 16°C and 50°C . The optimum temperature for siderophore production was 37°C , but seven isolates (R40, C43, R44, R45, J49, A55 and R61) showed high siderophore production, 70.90 ± 0.77 , 73.90 ± 1.39 , 97.24 ± 4.57 , 59.81 ± 1.31 , 108.93 ± 0.76 , 96.54 ± 0.80 and 75.39 ± 1.06 , at 28°C after 72 h incubation (Fig. 3A). This result indicates the specificity of isolates to temperature. Other isolates had moderate siderophore production at room temperature indicating that they are more versatile in the context of siderophore production. Isolates P88, T125, M132, M136, M138 and M139 showed the highest siderophore production at $37 \pm 2^\circ\text{C}$ instead of room temperature (107.57 ± 4.65 , 165.71 ± 5.26 , 135.03 ± 4.68 , 147.70 ± 1.38 , 141.57 ± 2.50 and 182.03 ± 7.97 , respectively), which may be due to the adaptation of FPs to the environmental temperature. It was observed that with a change in pH, there was a marginal difference in siderophore production. The highest siderophore production was observed at pH 7. But production drastically declined as pH diverged towards acidic and alkaline ranges (Fig. 3B). The effect of Fe^{3+} on siderophore production showed that $1 \mu\text{M}$ iron concentration resulted in the highest siderophore production, 53.86 ± 0.58 (by isolate A55) followed by 53.68 ± 0.16 and 51.77 ± 0.26 (by the isolates M138 and M132, respectively) (Fig. 4). In the present study an increased siderophore production with reduced Fe^{3+} concentration was observed.




Fig. 3. Impact of (A) temperature and (B) different pH on hydroxamate siderophore production by fluorescent *Pseudomonas* isolates. Data presented are mean \pm standard error of three independent replicate experiments. Means were compared between treatments using two-way ANOVA. Same letters are not significantly different according to Tukey's HSD-test ($p \leq 0.0001$) between treatments. Capital letters indicate significant mean difference between variables and small letters indicate significance between time




Fig. 4. Production of hydroxamate siderophores by fluorescent *Pseudomonas* isolates in the presence of varying concentrations of Fe³⁺ in succinic acid medium. Data presented are mean \pm standard error of three independent replicate experiments. Means were compared between treatments using two-way ANOVA. Same letters are not significantly different according to Tukey's HSD-test ($p \leq 0.0001$) between treatments. Capital letters indicate significant mean differences between concentrations and small letters indicate significance between time

The specificity of pH in siderophore production indicates that the soil pH is a critical factor for iron acquisition mechanisms. In addition to pH, heavy metal concentrations in the bacterial environment also influence siderophore biosynthesis (Rajkumar *et al.* 2009). Biosynthesis and secretion of siderophore by microbial systems are known to be related to the requirement of iron for the metabolism of specific growth substrates. In the environment, the presence of aromatic compounds along with easily metabolizable co-substrates supports siderophore production (Gaonkar *et al.* 2012). Environmental factors modulating siderophore synthesis include pH, the ionic state of iron, the presence of other trace elements and an adequate supply of carbon, nitrogen and phosphorus

(Duffy and Défago 1999). Bacterial growth as well as siderophore production is stimulated by ammonium sulphate and amino acids. However, the optimum siderophore yield was obtained with urea (Sayed *et al.* 2005). Statistical analysis using two-way ANOVA in the analysed parameters revealed a significant difference in p value except in siderophore production at 16°C, where p value was 0.78.

Iron-binding affinity of siderophores

The binding affinity experiments showed that hydroxamates have an expeditious affinity with Fe³⁺, whereas there was a change in affinity with respect to the surrounding iron concentration. A drastic absorption of Fe³⁺ by siderophores was observed within 5 min, followed by non-significant increments in absorption (Fig. 5). Further increments in affinity were less, which were evident in all three concentrations of Fe³⁺, as free siderophore ligands left in the sample were extremely limited. Different concentrations of Fe³⁺ (i.e., 10, 25 and 50 µM) did not show much variation in affinity. Binding affinity with Fe³⁺ showed well advanced fondness with the crucial metal ion. The siderophore-iron binding is completed during the initial 5 min and the affinity progress against time is uniform even at higher iron intensity.

The role of mutation in siderophoregenesis

From the mutation study results with UV and EtBr (Figs. 6A and B) it can be concluded that the resulting mutation suppressed siderophore production to a great extent. UV at 60 s and EtBr at higher concentrations (0.5 and 1.0 µM) further suppressed hydroxamate biosynthesis. Ferric uptake regulator (*Fur*) is a widespread bacterial protein that regulates the expression of iron acquisition and storage systems in




Fig. 5. Time course assay of *in vitro* iron binding affinity of siderophores produced by fluorescent *Pseudomonas* isolates, with varying ferric ion (Fe^{3+}) concentration. Data represented are the mean \pm standard error of three replicates. Vertical bars indicate standard error. Two-way ANOVA analysis showed a non-significant p value ($p < 0.0001$)

response to intracellular iron. *Fur* is a key regulator of iron metabolism, acting as a classical iron-dependent repressor of protein-coding genes (Lewin *et al.* 2002). Transcriptional studies on *P. fluorescens* Pf-5, showed that among the up-regulated pyoverdine biosynthesis gene clusters, the highly expressed gene was *pvdS* which encoded for the extra-cytoplasmic function sigma factor, a transcriptional regulator of pyoverdine biosynthesis genes (Lim *et al.* 2012). Bacterial evolution demands mutations to permit adaptation through sequence variation by still preserving genome integrity. UV, the customary non-ionizing radiation, can induce adjacent pyrimidine bases in a DNA strand to become a covalent dimer causing oxidative damage to DNA (Kozmin *et al.* 2008). Both of the mutations could not stimulate higher siderophore production in bacterial isolates. It was observed that mutation in bacterial genome resulted in suppression of ferric homeostasis, reported by Schwyn and Neilands

(1987). Genomic mutation on FP, major rhizosphere group bacteria, which affects siderophore-iron binding efficiency, has to be analysed in detail.

Mycelial inhibition traits and phytoaugmentation traits

All the isolates were checked for their ability to inhibit the pathogenic *F. oxysporum* strain and the results are represented in Table 2. It is worth pointing out that all the high siderophore producers inhibited the *F. oxysporum*, where M140 demonstrated maximum inhibition (55.0 ± 2.6 mm). Hence, it can be speculated that production of siderophores, possibly in combination with antimicrobial agents, produced by FPs, leads to *F. oxysporum* inhibition. Plant protection through siderophore producing rhizobacteria has emerged as a possibility in sustainable crop management (Karimi *et al.* 2012; Solanki *et al.* 2014). The rhizobacterially synthesized siderophores are excellent competitors for Fe, thus reducing the Fe availability to the phytopathogens (Beneduzi *et al.* 2012). This is a very common and significantly useful action of rhizobacteria in sustainable agriculture.

Growth promotion study results showed that the FPs are capable of inducing root and shoot growth with higher germination percentages than in control plants. Bacterial seed treatment resulted in improved physical parameters, where M140 showed the longest shoot length 53.8 ± 02.4 mm and root length 209.6 ± 05.9 mm, against control.

Plant growth promotion effect exerted by siderophore producing rhizobacteria, especially FPs has been reported previously in different plant systems (Sharma and Johri 2003; Ahmed and Holmström 2014). A report by Anitha and Kumudini (2014) clearly demonstrated the impact of secondary metabolites, especially siderophores, in the growth promotion of tomato plants, using FPs.




Fig. 6. Effect of (A) UV mutagenesis and (B) EtBr on time course production of hydroxamate siderophores by fluorescent *Pseudomonas* isolates. Data presented are mean \pm standard error of three independent replicate experiments. Means were compared between treatments using two-way ANOVA. Same letters are not significantly different according to Tukey's HSD-test ($p \leq 0.0001$) between treatments. Capital letters indicate significant mean difference between concentration and small letters indicate significance between time

Table 2. *In vitro* antifungal activity against *Fusarium oxysporum* and phytoaugmentation traits of fluorescent *Pseudomonas* isolates

<i>Pseudomonas</i> isolates	Zone of inhibition [mm]	Shoot length [mm]	Root length [mm]
P88	0	53.9 ± 03.5	160.4 ± 10.6
T125	0	64.8 ± 03.6	170.4 ± 05.6
T106	0	52.2 ± 04.5	154.2 ± 09.2
M131	24.3 ± 2.8	38.1 ± 02.3	86.0 ± 08.7
M132	27.2 ± 2.3	51.2 ± 03.7	161.8 ± 06.5
M136	40.3 ± 0.9	34.4 ± 03.3	184.4 ± 07.6
M138	26.3 ± 2.7	48.3 ± 02.5	202.4 ± 06.9
M139	43.0 ± 2.6	44.9 ± 02.8	201.2 ± 05.5
M140	55.0 ± 2.6	53.8 ± 02.4	209.6 ± 05.9
R40	0	34.1 ± 06.9	124.0 ± 08.5
R44	0	38.7 ± 03.8	113.7 ± 08.2
R45	0	46.4 ± 16.2	113.3 ± 34.6
R61	0	37.5 ± 03.0	113.0 ± 06.8
C43	0	32.4 ± 02.1	155.7 ± 11.3
J49	0	41.6 ± 03.4	92.6 ± 08.3
A55	0	39.6 ± 03.5	102.4 ± 07.3
Control*	—	29.0 ± 02.3	97.4 ± 04.7

*distilled water treated control

Conclusions

There are various reports that pesticides, designed to target specific organisms, also affect non-target microbial flora also. Fungicides and insecticides are reported to be capable of exerting an adverse impact on natural soil microflora (Cycoń and Piotrowska-Seget 2007). However, there are very few studies on the impact of siderophore production, on various soil factors in the terrestrial habitat. The present study undertook a detailed analysis of siderophore characterization, and of the common factors which affect their production. It is known that maize plants (Gramineous family) produce phytosiderophores which offer a competence to the bacterial siderophores in rhizosphere terrain. This could be the reason for its high siderophore production nature. The difference in the quantities of siderophores produced by different organisms is logical. As the natural iron 'collectors' in the soil rhizosphere the chemical agents which are used need further study since they possibly spoil the natural biocontrol agents, and can have a long-term impact on nature.

Acknowledgements

The authors are thankful to the Jain University for financial and infrastructure support. The constructive comments of Prof. Leela Iyengar are gratefully acknowledged.

References

- Adler C., Corbala N.S., Seyedsayamdost M.R., Pomares M.F., de Cristobal R.E., Clardy J., Kolter R., Vincent P.A. 2012. Catecholate siderophores protect bacteria from pyochelin toxicity. *PloS One* 7 (10): e46754. DOI: <https://doi.org/10.1371/journal.pone.0046754>
- Ahmed E., Holmström S.J.M. 2014. Siderophores in environmental research: roles and applications. *Microbiology and Biotechnology* 7 (3): 196–208. DOI: <https://doi.org/10.1111/1751-7915.12117>
- Alexander B., Zuberer D.A. 1991. Use of chrome azurol S reagents to evaluate siderophore production by rhizosphere bacteria. *Biology and Fertility of Soils* 12 (1): 39–45. DOI: <https://doi.org/10.1007/bf00369386>
- Ali S.S., Vidhale N.N. 2011. Evaluation of siderophore produced by different clinical isolate *Pseudomonas aeruginosa*. *Indian Journal of Medical Research* 3 (3): 131–135. DOI: <https://doi.org/10.9735/0975-5276.3.3.131-135>
- Anitha G., Kumudini B.S. 2014. Isolation and characterization of fluorescent pseudomonads and their effect on plant growth promotion. *Journal of Environmental Biology* 35: 627–634.
- Arnou L.E. 1937. Colorimetric determination of the components of 3,4-dihydroxyphenylalanine tyrosine mixtures. *Journal of Biological Chemistry* 118: 531.
- Baakza A.A., Vala A.K., Dave B.P., Dube H.C. 2004. A comparative study of siderophore production by fungi from marine and terrestrial habitats. *Journal of Experimental Marine Biology and Ecology* 311 (1): 1–9. DOI: <https://doi.org/10.1016/j.jembe.2003.12.028>
- Beneduzi A., Ambrosini A., Passaglia L.M. 2012. Plant growth-promoting rhizobacteria (PGPR): their potential as antagonists and biocontrol agents. *Genetics and Molecular Biology* 35 (4): 1044–1051. DOI: <https://doi.org/10.1590/s1415-47572012000600020>
- Bhattacharya A. 2010. Siderophore mediated metal uptake by *Pseudomonas fluorescens* and its comparison to iron (iii)

- chelation. Ceylon Journal of Sciences 39 (2): 147–155. DOI: <https://doi.org/10.4038/cjsbs.v39i2.3001>
- Cappuccino J.C., Sherman N. 2013. Microbiology: A Laboratory Manual. 10th ed. Pearson Publishing, New York, USA, 560 pp.
- Cycoń M., Piotrowska-Seget Z. 2007. Effect of selected pesticides on soil microflora involved in organic matter and nitrogen transformations: Pot experiment. Polish Journal of Ecology 55 (2): 207–220.
- Duffy B.K., Défago G. 1999. Environmental factors modulating antibiotic and siderophore biosynthesis by *Pseudomonas fluorescens* biocontrol strains. Applied and Environmental Microbiology 65 (6): 2429–2438.
- Fallahzadeh-Mamaghani V., Ahmadzadeh M., Sharifi R. 2009. Screening systemic resistance-inducing fluorescent pseudomonads for control of bacterial blight of cotton caused by *Xanthomonas campestris* pv. *malvacearum*. Journal of Plant Pathology 91 (3): 663–670. DOI: <http://dx.doi.org/10.4454/jpp.v91i3.558>
- Gaonkar T., Nayak P.K., Garg S., Bhosle S. 2012. Siderophore-producing bacteria from a sand dune ecosystem and the effect of sodium benzoate on siderophore production by a potential isolate. Scientific World Journal 112: 1–8. DOI: <https://doi.org/10.1100/2012/857249>
- Gawel D., Maliszewska-Tkaczyk M., Schaaper J.P.R.M., Fijalkowska I.J. 2012. Lack of strand bias in UV-induced mutagenesis in *Escherichia coli*. Bacteriology 184 (6): 4449–4454. DOI: <https://doi.org/10.1128/jb.184.16.4449-4454.2002>
- Gupta V., Saharan K., Kumar R., Gupta R., Sahai V., Mittal A. 2008. Spectrophotometric ferric ion biosensor from *Pseudomonas fluorescens* culture. Biotechnology and Bioengineering 100 (2): 284–296. DOI: <https://doi.org/10.1002/bit.21754>
- Helmy M., Baddar D., El'Masry M.H. 2008. Affinity purification of a siderophore that exhibits an antagonistic effect against soft rot bacterium. Biochemistry 73 (7): 776–782. DOI: <https://doi.org/10.1134/s0006297908070055>
- Hemanth G., Kumar P.K.R., Niharika P.S., Kolli S.K. 2016. Fungicides effect on soil micro flora in Tekkali Mandal, Srikakulam (Dist). International Journal of Research and Development in Pharmacy and Life Sciences 5 (4): 2245–2250.
- Jenifer A.M.R., Reena A., Aysha O.S., Valli S., Nirmala P., Vinothkumar P. 2013. Isolation of siderophore producing bacteria from rhizosphere soil and their antagonistic activity against selected fungal plant pathogens. International Journal of Current Microbiology and Applied Science 2 (1): 59–65.
- Jin C.W., Li G.X., Yu X.H., Zheng S.J. 2010. Plant Fe status affects the composition of siderophore-secreting microbes in the rhizosphere. Annals of Botany 105 (5): 835–841. DOI: <https://doi.org/10.1093/aob/mcq071>
- Kamal F., Mehrgan H., Assadi M.M., Mortazavi S.A. 2003. Mutagenesis of *Xanthomonas campestris* and selection of strains with enhanced xanthan production. Iranian Biomedical Journal 7 (3): 91–98.
- Karimi K., Amini J., Harighi B., Bahramnejad B. 2012. Evaluation of biocontrol potential of *Pseudomonas* and *Bacillus* spp. against *Fusarium* wilt of chickpea. Australasian Journal of Crop Sciences 6 (4): 695–703.
- Khan H., Parmar N., Kahlon R.S. 2016. *Pseudomonas*-plant interactions I: Plant growth promotion and defense-mediated mechanisms. p. 419–469. In: “*Pseudomonas*: Molecular and Applied Biology” (R. Kahlon, ed.). Springer International Publishing, Switzerland.
- Kozmin S., Slezak G., Reynaud-Angelin A., Elie C., de Rycke Y., Boiteux S., Sage E. 2008. UVA radiation is highly mutagenic in cells that are unable to repair 7,8-dihydro-8-oxoguanine in *Saccharomyces cerevisiae*. Proceedings of National Academy of Sciences USA 102: 3538–3543.
- Krewulak K.D., Vogel H.J. 2008. Structural biology of bacterial iron uptake. Biochimica et Biophysica Acta 1778 (9): 1781–1804. DOI: <https://doi.org/10.1016/j.bbamem.2007.07.026>
- Kumudini B.S., Jayamohan N.S. & Patil S.V. 2017. Integrated mechanisms of plant disease containment by rhizospheric bacteria: Unraveling the signal cross-talk between plant and fluorescent *Pseudomonas*. p. 263–291. In: “Agriculturally Important Microbes for Sustainable Agriculture” (V. Meena, P. Mishra, J. Bisht, A. Pattanayak, eds.) Springer International Publishing. DOI: https://doi.org/10.1007/978-981-10-5343-6_9
- Lee W., van Baalen M., Jansen V.A.A. 2011. An evolutionary mechanism for diversity in siderophore-producing bacteria. Ecology Letters 15 (2): 119–125. DOI: <https://doi.org/10.1111/j.1461-0248.2011.01717.x>
- Lemanceau P., Bauer P., Kraemer S., Briat J.-F. 2009. Iron dynamics in the rhizosphere as a case study for analyzing interactions between soils, plants and microbes. Plant and Soil 321 (1–2): 513–535. DOI: <https://doi.org/10.1007/s11104-009-0039-5>
- Lewin A.C., Doughty P.A., Flegg L.A., Moore G.R., Spiro S. 2002. The ferric uptake regulator of *Pseudomonas aeruginosa* has no essential cysteine residues and does not contain a structural zinc ion. Microbiology 148 (8): 2449–2456. DOI: <https://doi.org/10.1099/00221287-148-8-2449>
- Lim C.K., Hassan K.A., Tetu S.G., Loper J.E., Paulsen I.T. 2012. The effect of iron limitation on the transcriptome and proteome of *Pseudomonas fluorescens* Pf-5. PLoS One 7 (6): e39139. DOI: <https://doi.org/10.1371/journal.pone.0039139>
- Masaki J., Fujita M.J., Nakano K., Sakai R., Bisucaberin B. 2013. A linear hydroxamate class siderophore from the marine bacterium *Tenacibaculum mesophilum*. Molecules 18 (4): 3917–3926. DOI: <https://doi.org/10.3390/molecules18043917>
- Meyer J.-M., Geoffroy V.A., Baida N., Gardan L., Izard D., Lemanceau P., Achouak W., Palleroni N.J. 2002. Siderophore typing, a powerful tool for the identification of fluorescent and nonfluorescent pseudomonads. Applied and Environmental Microbiology 68 (6): 2745–2753. DOI: <https://doi.org/10.1128/aem.68.6.2745-2753.2002>
- Miethke M., Marahiel M.A. 2007. Siderophore-based iron acquisition and pathogen control. Microbiology and Molecular Biology Reviews 71 (3): 413–451. DOI: <https://doi.org/10.1128/mmbr.00012-07>
- Mishra P.K., Mishra S., Selvakumar G., Bisht S.C., Bisht S.C., Bisht J.K., Kundu S., Gupta H.S. 2008. Characterisation of a psychrotolerant plant growth promoting *Pseudomonas* sp. strain PGERs17 (MTCC 9000) isolated from North Western Indian Himalayas. Annals of Microbiology 58 (4): 561–568. DOI: <https://doi.org/10.1007/bf03175558>
- Murugappan R.M., Aravindh A., Karthikeyan M. 2011. Chemical and structural characterization of hydroxamate siderophore produced by marine *Vibrio harveyi*. Journal of Indian Microbiology and Biotechnology 38 (2): 265–273. DOI: <https://doi.org/10.1007/s10295-010-0769-7>
- Nagata T., Oobo T., Aozasa O. 2013. Efficacy of a bacterial siderophore, pyoverdine, to supply iron to *Solanum lycopersicum* plants. Journal of Bioscience and Bioengineering 115 (6): 686–690. DOI: <https://doi.org/10.1016/j.jbiosc.2012.12.018>
- Neilands J.B. 1981. Microbial iron compounds. Annual Review of Biochemistry 50: 715–731.
- Rajkumar M., Ae N., Prasad N.M.V., Freitas. H. 2009. Potential of siderophore-producing bacteria for improving heavy metal phytoextraction. Trends in Biotechnology 28 (3): 142–149. DOI: <https://doi.org/10.1016/j.tibtech.2009.12.002>
- Mendes R., Garbeva P., Raaijmakers J.M. 2013. The rhizosphere microbiome: significance of plant beneficial, plant pathogenic, and human pathogenic microorganisms. FEMS Microbiology Reviews 37 (5): 634–663.
- Sah S., Singh R. 2015. Siderophore: Structural and functional characterisation – A comprehensive review. Agriculture 61 (3): 97–114. DOI: <https://doi.org/10.1515/agri-2015-0015>
- Saraf M., Pandya U., Thakkar A. 2014. Role of allelochemicals in plant growth promoting rhizobacteria for biocontrol of

- phytopathogens. *Microbiology Research* 169 (1): 18–29. DOI: <https://doi.org/10.1016/j.micres.2013.08.009>
- Sayyed R.Z., Badguzar M.D., Sonawane H.M., Mhaske M.M., Chincholkar S.B. 2005. Production of microbial iron chelators (siderophores) by fluorescent pseudomonads. *Indian Journal of Biotechnology* 4: 484–490.
- Schwyn B., Neilands J.B. 1987. Universal chemical assay for the determination of siderophores. *Analytical Biochemistry* 56: 47–56.
- Sharma A., Johri B.N. 2003. Growth promoting influence of siderophore-producing *Pseudomonas* strains GRP3A and PRS9 in maize (*Zea mays* L.) under iron limiting conditions. *Microbiological Research* 158 (3): 243–248. DOI: <https://doi.org/10.1078/0944-5013-00197>
- Snow G.A. 1954. Mycobactin, a growth factor for *Mycobacterium johnei*: II. Degradation and identification of fragments. *Journal of Chemical Society*: 2588–2596. DOI: <https://doi.org/10.1039/jr9540002588>
- Solanki M.K., Singh R.K., Srivastava S., Kumar S., Kashyap P.L., Srivastava A.K., Arora D.K. 2014. Isolation and characterization of siderophore producing antagonistic rhizobacteria against *Rhizoctonia solani*. *Journal of Basic Microbiology* 54 (6): 585–596. DOI: <https://doi.org/10.1002/jobm.201200564>
- Subramanian J., Satyan K. 2014. Isolation and selection of fluorescent pseudomonads based on multiple plant growth promotion traits and siderotyping. *Chilean Journal of Agricultural Research* 74 (3): 319–325. DOI: <https://doi.org/10.4067/s0718-58392014000300010>
- Taylor A.J., Joshi B.H. 2012. Characterization and optimization of siderophore production from *Pseudomonas fluorescens* strain isolated from sugarcane rhizosphere. *International Journal of Engineering Research and Development* 6 (3): 688–694.
- Tripathi M., Munot H.P., Shouche Y., Meyer J.M., Goel R. 2005. Isolation and functional characterization of siderophore-producing lead- and cadmium-resistant *Pseudomonas putida* KNP9. *Current Microbiology* 50 (5): 233–237. DOI: <https://doi.org/10.1007/s00284-004-4459-4>
- Varsha T., Kumudini B.S. 2016. Fluorescent *Pseudomonas* mediated alleviation of trivalent chromium toxicity in ragi through enhanced antioxidant activities. *Proceedings of National Academy of Sciences India, Sect B Biological Sciences*: 1–9. DOI: <https://doi.org/10.1007/s40011-016-0816-x>
- Widmer F., Seidler R.J., Watrud L.S., Gillevet P.M., Di Giovanni G.D. 1998. A highly selective PCR protocol for detecting 16S rRNA genes of the genus *Pseudomonas* (*sensu stricto*) in environmental samples. *Applied and Environmental Microbiology* 64 (7): 2545–2553.
- Ye L., Ballet S., Hildebrand F., Laus G., Guillemyn K., Raes J., Matthijs S., Martins J., Cornelis P. 2014. A combinatorial approach to the structure elucidation of a pyoverdine siderophore produced by a *Pseudomonas putida* isolate and the use of pyoverdine as a taxonomic marker for typing *P. putida* subspecies. *BioMetals* 26 (4): 561–575. DOI: <https://doi.org/10.1007/s10534-013-9653-z>
- Yimer A.M. 2015. Review on marvelous incidence of the iron observable in livelihood and its history. *Natural Products Chemistry and Research* 3 (5): 3–5. DOI: <https://doi.org/10.4172/2329-6836.1000185>
- Zongzheng Y., Xin L., Zhong L., Jinzhao P., Jin Q., Wenyan Y. 2009. Effect of *Bacillus subtilis* SY1 on antifungal activity and plant growth. *International Journal of Agricultural and Biological Engineering* 2 (4): 55–56. DOI: 10.3965/j.issn.1934-6344.2009.04.055-061