

ORIGINAL ARTICLE

Endophytic colonization by *Beauveria bassiana* and *Metarhizium anisopliae* induces growth promotion effect and increases the resistance of cucumber plants against *Aphis gossypii*

Roshan S. Shaalan^{1,3*} , Elvis Gerges², Wassim Habib², Ludmilla Ibrahim³¹ Department of Plant Protection, University of Forestry, Sofia, Bulgaria² Department of Plant Protection, Lebanese Agricultural Research Institute, Lebanon³ Department of Plant Protection, Lebanese University, Beirut, Lebanon

Vol. 61, No. 4: 358–370, 2021

DOI: 10.24425/jppr.2021.139244

Received: February 2, 2021

Accepted: May 27, 2021

Online publication: December 20, 2021

*Corresponding address:
roshan.shaalan23@hotmail.comResponsible Editor:
Natasza Borodynko-Filas

Abstract

The entomopathogenic fungi (EPF) are characterized as fungi with various functions and numerous mechanisms of action. The ability to establish themselves as beneficial endophytes provides a sound ground for their exploitation in crop production and protection. The purpose of this study was to evaluate the entomopathogenic strains of *Beauveria bassiana* and *Metarhizium anisopliae* for their potential to colonize cucumber plants under natural environmental conditions in non-sterile substrate. Seed submersion in conidial suspension resulted in systemic colonization of cucumber plants 28 days post-inoculation. Scanning electron microscope micrographs demonstrated that conidia of both fungal genera have adhered, germinated and directly penetrated seed epidermal cells 24 hr post-submersion. Treated with EPF cucumber seeds resulted seedlings tissues of which contained a significantly higher amount of total phenolic compounds and unchanged amounts of chlorophylls. There was a significant negative effect of endophytic colonization on the *Aphis gossypii* population size after 5 days of exposure as well as a positive effect on cucumber growth and development 7 weeks post-inoculation. We suggest that reduction of *A. gossypii* population on mature *Cucumis sativus* plants is caused via an endophyte-triggered improvement of plant's physiological parameters such as enhanced plant growth with subsequent increase in plant resistance through augmented production of phenolic compounds.

Keywords: aphid biocontrol, entomopathogenic fungi as endophytes, growth promoters, phenolic compounds

Introduction

Cucumber, *Cucumis sativus* L. (Cucurbitaceae), is an important cash crop cultivated around the world, contributing to the total income and living standards of many societies through increased foreign exchange (Vandre 2013). The popularity of its fruit rests not only on its availability but also on its essential nutrients (e.g., vitamins and minerals) and other beneficial substances (e.g., soluble fibers and antioxidants). Increasing demand and sustainable availability of this commodity depend on its enhanced productivity. Since cucumber

is one of the most important greenhouse crops, its intensive production is more susceptible to the rapid spread of problematic pests such as cotton aphid, *Aphis gossypii* Glover (Hemiptera: Aphididae). The damages caused by sucking the sap of its host plants, honeydew production, and by the transmission of plant viruses (Blackman and Eastop 2000) are problems that pose a serious concern about yield and quality of greenhouse crops (Hermoso de Mendoza *et al.* 2001). This pest is mainly controlled by chemical insecticides. However,

persistent use of these insecticides has resulted in the development of resistance by the pest (Shi *et al.* 2012), in the decrease of its natural enemy populations (Godfrey *et al.* 2000), in the resurgence of secondary pests, as well as in posing environmental and economic hazards. These problems have brought about the need to introduce other sustainable, effective, and safe alternative control strategies against *A. gossypii*.

Over the years, various biological control tactics have been evaluated in many parts of the world for the management of this pest. The most commonly used approach is augmentation of the seven-spot beetle, *Coccinella septempunctata* L. (Coleoptera: Coccinellidae) in cotton fields (Omkar 2004) and the aphid parasitoid, *Aphidius colemani* Viereck (Hymenoptera: Braconidae) in greenhouses (Gissella *et al.* 2006). However, entomopathogenic fungi (EPF), *Beauveria bassiana* (Balsamo) Vuillemin and *Metarhizium anisopliae* (Metchnikov) Sorokin (Ascomycota: Hypocreales) have also been shown to be promising alternatives. They are currently being developed, registered, marketed and used as biopesticides against many insect species (Lacey 2016) including aphids (Ibrahim *et al.* 2015). These and many other fungal entomopathogens have been inundated as foliar and/or soil applications to manage agricultural insect pests more effectively (Skinner *et al.* 2014). Their effectiveness, however, is limited by adverse environmental conditions such as UV, temperatures and low humidity (Wraight *et al.* 2007; Vega *et al.* 2012). Therefore, this method relies on the direct action of the released agent rather than on secondary effects on successive pest generations (Vincent *et al.* 2007).

As a result, recent research is aimed at introducing fungal entomopathogens as endophytes that allow colonization of internal plant tissues without causing apparent harm to the host (Wilson 1995). There is a general opinion that entomopathogenic endophytes, while internally protected, could cope better with negative environmental factors (Vega *et al.* 2009). Although the ecological function of endophytic EPFs remains largely unknown, some studies have implicated them in plant growth as probiotics (Dora 2013), herbivore and plant disease resistance (Vega *et al.* 2008), increased stress tolerance of plants to abiotic factors (Rodriguez *et al.* 2009) and bioremediation of heavy metals (Bajan *et al.* 1998). Some of the fungi have been reported as naturally occurring endophytes while others have been introduced into plants using different techniques (Vega *et al.* 2008; Bamisile *et al.* 2018). Among the most studied fungal endophytes are the grass endophytes in the genus *Neotyphodium* (Ascomycota) which has shown different levels of activity against grass herbivores including grubs (Saikkonen *et al.* 2006). However, recent findings show that EPFs could be successfully explored as endophytes for the management of many insect pests (Vidal and Jaber

2015) including aphids (Castillo-Lopez *et al.* 2014). The endophytic colonization of *B. bassiana* and *M. anisopliae* in cucumber plants was achieved through artificial inoculation of seeds (Shaalan and Ibrahim 2019). *B. bassiana* has been reported as an endophyte in various crop plants, but its endophytic activities against *A. gossypii* have been studied only on cotton *Gossypium hirsutum* L. (Malvales: Malvaceae), pumpkin, *Curcubita pepo* (Cucurbitaceae) (Gurulingappa *et al.* 2011; Castillo-Lopez *et al.* 2014), and melon, *Cucumis melo* (Cucurbitaceae) (González-Mas *et al.* 2019). A significant reduction in the reproduction rate of *A. gossypii* and populations of bean aphid, *Aphis fabae* Scopoli (Hemiptera: Aphididae) and pea aphid, *Acyrtosiphon pisum* Harris (Hemiptera: Aphididae) was reported on *B. bassiana* colonized cotton plants and fava bean, *Vicia faba* L. (Fabales: Fabaceae), respectively (Akello and Sikora 2012; Castillo-Lopez *et al.* 2014). However, *M. anisopliae* had no effect on bean aphids. On the other hand, population sizes of soybean aphid, *Aphis glycines* Matsumura increased on soybean plants colonized by *M. brunneum* but did not change in similar treatments with *B. bassiana* (Clifton *et al.* 2018). Such discrepancy suggests that endophytic EPF colonizing the plant material may induce or suppress plant systemic resistance to insect attack by altering a plant's kairomones (plant signaling molecules) or secondary plant metabolites (e.g., terpenoids) (Vega 2018). The abilities of endophytes to prime plant defense mechanisms with chemical or biological inducers against future insect attack depend on microbial, biochemical and physiological conditions of plant material (Jaber and Enkerli 2016), microbial and plant host genotypes, time of the analysis and their interactions (Pańka *et al.* 2013). To date, there is no information in the literature on the endophytic effects of *M. anisopliae* on *A. gossypii* in cucumber production.

Therefore, the aims of the present study were to: (1) evaluate the ability of *B. bassiana* and *M. anisopliae* to establish themselves in cucumber plants through seed inoculation; (2) determine if EPF colonization modifies the physiological and biochemical status of cucumber plants and (3) investigate the effects of seed inoculation on the population of *A. gossypii* on cucumber.

Materials and Methods

Source of fungal inoculum

Two strains, *B. bassiana* and *M. anisopliae*, isolated from fallow soil (Ibrahim *et al.* 2011) were used in this study. Long term storage of the isolates was achieved by freezing conidia in 35% w/w glycerol at -80°C . When prepared, frozen conidia were re-hydrated by suspending in a small volume of sterile 0.03% Tween

20 solution, placed on potato dextrose agar (PDA) and incubated at $23 \pm 2^\circ\text{C}$ in the dark for 14 days (Shaalán and Ibrahim 2019).

Molecular identification of *B. bassiana* and *M. anisopliae*

Mycelial plugs from 2-day-old single spore cultures of *B. bassiana* and *M. anisopliae* were placed on malt extract agar (MEA) overlaid with a disc of sterilized cellophane and incubated at $21 \pm 1^\circ\text{C}$ in the dark for 2 days. The resultant mycelia were reduced to powder using iron balls, liquid nitrogen and tissue lyser Qiagen (Hilden, Germany) for subsequent DNA extraction using the CTAB method (Lee *et al.* 1988; Wu *et al.* 2001). DNA from both isolates were purified with 1% RNAase A ($10 \text{ mg} \cdot \text{ml}^{-1}$), and precipitated with 5 M ammonium acetate and absolute ethanol. A NanoDrop spectrophotometer (Shimadzu, Japan) was used to quantify the amount of purified DNA required for amplification. The internally transcribed spacer (ITS) region was amplified using ITS4 (5'-TCC TCCGCTTATTGATATGC-3') and ITS5 (5'-GGAAG TAAAAGTCGTAACAAGG-3') primers (White *et al.* 1990) using Taq PCR Master Mix (Qiagen, USA). The specific mixtures of this reaction contained 1 X ready Master Mix, 3 mM MgCl_2 , 0.2 μM of each primer, and 25 ng of template DNA in 25 μl . The following PCR conditions were adopted: initial denaturation step of 5 min at 95°C followed by 25 cycles, each consisting of a 1 min denaturation step at 95°C , a 1 min annealing step at 58°C , and a 1 min extension step at 72°C , and a final extension step at 72°C for 7 min. The polymerase chain reaction (PCR) products were purified using QIAquick PCR amplification kit (Qiagen) and sequenced using Sanger sequencing at the Medical Genetics Unit of Saint Joseph University of Beirut (USJ). For species identification, sequences were compared with the available sequences in the NCBI BLAST database. The chromatographs were viewed and edited as needed using Chromas software (<https://chromas.software.informer.com/download/>).

Sterilization of cucumber seeds

Variety Beit Alpha hybrid cucumber seeds were used in this experiment. In order to remove possible epiphytic microorganisms by denaturing their DNA from the aerial surfaces, the cucumber seeds were surface sterilized prior to their artificial inoculation. Seeds were soaked in sterile distilled water (dH_2O) for 15 min, and then surface-sterilized by immersion in 3% sodium hypochlorite for 2 min with subsequent rinsing with sterile dH_2O and additional immersion in 70% ethanol for 2 min, followed by 3 rinses with sterile dH_2O . The axenically treated seeds were stored overnight at 5°C for synchronization of seed

germination and growth. To assess the effectiveness of the sterilization method, 20 randomly selected seeds from the seeds used in the screening test were placed onto PDA, incubated at 25°C in darkness and then examined after 3, 5 and 10 days. The disinfection was considered successful when no fungal growth was observed on the PDA plates (Shaalán and Ibrahim 2019).

Preparation of fungal inoculum and seed inoculation

Conidia from 14-day-old cultures grown on PDA were harvested by scraping the agar surface with a sterile spatula, suspended in 0.03% Tween 20 and shaken at 354 rpm for 30 min. Resultant suspensions were filtered through sterile cheesecloth under sterile conditions in order to remove any hyphal fragments. Conidial concentrations were determined using a light microscope and hemocytometer (Fuchs-Rosenthal), and later adjusted to 1×10^6 conidia $\cdot \text{ml}^{-1}$ with sterile 0.03% Tween 20 solution. Axenic seeds were immersed in fungal conidial suspensions of *B. bassiana* and *M. anisopliae* for 2 h. Preliminary tests indicated that a conidial concentration of 10^6 conidia $\cdot \text{ml}^{-1}$ and a 2 h immersion time were the optimal parameters for the most successful colonization of cucumber seeds and plantlets tested by *B. bassiana* and *M. anisopliae* isolates (Shaalán and Ibrahim 2019). Inoculated seeds were then sown on seed trays containing non-sterile compost supplemented with N $150 \text{ mg} \cdot \text{l}^{-1}$, P_2O_5 $185 \text{ mg} \cdot \text{l}^{-1}$, and K_2O $250 \text{ mg} \cdot \text{l}^{-1}$ (pH (H_2O) 5.2–6.0, neutral peat, MIKSKAAR) and grown under natural fluctuating night/day temperatures ($23\text{--}29^\circ\text{C}$) and relative humidity (RH, 30–80%). For the control group the seeds were immersed in sterile 0.03% Tween 20 only. Prior to planting, the compost was analyzed for the presence of soilborne fungal communities using the soil dilution method (Ratna Kumar *et al.* 2015; Umboh *et al.* 2016) with slight modification. For this purpose, 10 g of used peat were resuspended in 100 ml of sterile water. The resultant stock suspension was then shaken for 1 h at 200 rpm. Tenfold diluted (10^{-1}) samples were plated on 1/4-strength PDA media containing $20 \text{ mg} \cdot \text{l}^{-1}$ amoxicilline, $2 \text{ ml} \cdot \text{l}^{-1}$ Tween 20, and additional agar of 1.5% (w/v) final concentration. The test was replicated three times. The plates were incubated for 2 weeks at 25°C , and the fungal colonies which developed were then isolated further to be observed under a microscope and identified.

Assessment of fungal colonization

Assessment of inoculated seeds using scanning electron microscope (SEM)

Inoculated seeds were incubated at room temperature under sterile conditions for 24 h and then fixed in glutaraldehyde, dehydrated through an alcohol-

acetone series, dried in a critical-point drying apparatus, mounted on stubs in different positions and coated with gold in a gold metallization with a Cressington 108 auto sputter coater (Pathan *et al.* 2010). The specimens were observed and photographed with SEM (Seron ALS2100, Korea). There were 11 seeds examined for each treatment and the numbers of attached and germinated spores, and spores producing an appressorium or penetrating the seed coat were recorded on a randomly chosen surface area ($100 \mu\text{m}^2$) of observation on each seed.

Assessment of colonization using re-isolation

Five randomly selected 4-week-old seedlings (3–5 true leaf stage) were harvested from the trays, cleaned and surface-sterilized. First, the seedlings were immersed in 0.5% sodium hypochlorite for 2 min, and then in 70% ethanol for 30 s, followed by three rinses in sterile dH_2O . Disinfected seedlings were air-dried in a laminar flow hood for 15 min, the root, stem and leaf parts were cut using a sterile scalpel. The obtained plant materials were placed on PDA selective media amended with antibiotics ($20 \text{ mg} \cdot \text{l}^{-1}$ Amoxicilline), incubated at 25°C for 4 weeks and inspected regularly to observe fungal outgrowth. The success of the disinfection process was assessed by plating 0.5 ml of residual rinse water on PDA and by making imprints of surface sterilized plant tissue. Fungal outgrowth from the plated plant samples were identified as *B. bassiana* or *M. anisopliae* based on differential growth on selective media, colony morphology, and microscopic examination of conidia (Humber 1997). Percent colonization of different seedling parts by the respective inoculated fungus was calculated following the Petrini and Fisher (1987) formula: % colonization = number of sampled plant tissue showing fungal outgrowth divided by the total number of plated plant tissue samples $\times 100$. The test was repeated three times.

The effects of endophytic EPFs on the population size of *Aphis gossypii* on cucumber plants.

Propagation of cucumber plants

Cucumber seedlings at 1–2 true leaf stage were transplanted from seed trays into plastic pots (10 cm diameter and 15 cm depth) containing unsterilized compost and propagated for 5 weeks under the same conditions as described previously (see 2.4 section). The potted plants were regularly irrigated. Fifteen plants for each treatment were used.

Aphis gossypii culture

Prior to laboratory establishment of pure *A. gossypii* culture, aphid populations collected from different plant hosts and locations were identified using

morphological and molecular techniques. Primary morphological identifications were carried out according to Stoetzel *et al.* (1996) and Blackman (2010). For molecular identification, a single aphid body from each sample was used and total DNA was extracted according to the CTAB method (Gawel and Jarret 1991). Two sets of species-specific primers for *A. gossypii* (KBR(AG)-F-5'-TCTTCTCTAGAATTT TAATCCGATTA-3' and KBR(AG)-R-5'-AAGAATAGGGTCTCCCCCACCT-3' (Rebijith *et al.* 2012)) and for *Myzus persicae* (KBR(MP)-F-5'-TCATCACT TAGAATCTTAATTTCGTCTT-3' and KBR(MP)-R-5'-TGGTATTATATTTAAGATTGTACAAATA-3' (Rebijith *et al.* 2012)) were used in PCR reactions with genomic DNA as the template using the following conditions: initial denaturation step of 4 min at 94°C , followed by 35 cycles, each consisting of a 30 s denaturation step at 94°C , a 45 s annealing step at 64°C , and a 40 s extension step at 72°C , with a final extension step for 20 min at 72°C (Matallanas *et al.* 2013). PCR products were run on 1.2% agarose gel, stained with UVView loading dye (Bio-Rad, USA) and visualized under UV lights by the Molecular imager system Gel Doc 1000 (Bio-Rad) using ImageLab software (Bio-Rad). Colony individuals that were identified as *A. gossypii* were used for laboratory rearing and maintained on healthy cucumber plants grown in pots and in wooden cages ($70 \text{ cm} \times 50 \text{ cm} \times 50 \text{ cm}$) at $21 \pm 5^\circ\text{C}$. Adults of this colony were used in the bioassays.

Effect of endophytic EPFs on *Aphis gossypii* population size

Five-week-old plants were placed in individual clear plastic cages (20 cm dim. \times 30 cm high) and 10 *A. gossypii* adults were introduced to a middle leaf of each experimental plant. Cages were covered with lids and sealed with no-see-um mesh to prevent aphids from escaping. There were 36 caged plants (12 replicate plants for each treatment) randomly distributed and maintained under natural conditions ($25\text{--}30^\circ\text{C}$ and 30–80% RH).

After 5 days of exposure to the aphids, tested plants were observed and the total number of aphids (adults + nymphs) on each plant was recorded using a magnifier. This experiment was repeated twice.

The effects of endophytic EPFs on physiological and biochemical parameters of cucumber plants

The 6-week-old cucumber plants were examined, the number of fully developed leaves and flowers was measured and counted for each tested plant. One week later, the number of small cucumber fruits formed, and the height of each plant were also recorded.

Analysis of total phenolic compounds

Some studies have reported that grass endophytes indirectly influence the defense reaction of the host plant (Pańka *et al.* 2013) by involving e.g., synthesis of phenolic compounds. Phenolic compounds are the main class of secondary metabolites in plants and are divided into phenolic acids and polyphenols. They perform a variety of functions, including acting as antioxidants (Grace and Logan 2000). They may exist in free, soluble conjugated (acid and alkaline hydrolysable) and insoluble-bound forms (Perea-Domínguez *et al.* 2018). Cucumber plant is rich in polyphenolic compounds such as p-coumaric, caffeic, and ferulic acids and p-coumaric acid methyl ester (Daayf *et al.* 2000). The Folin-Ciocalteu (F-C) assay provides a convenient, rapid and simple estimation of the content of total phenolics and other oxidation substrates in plant extracts. This assay also provides a high-throughput tool for screening the response of plants to biotic factors and environmental stress. In this study, to see if fungal endophytes colonizing cucumber seedlings can alter the total content of phenolic compounds (TPCs), the F-C procedure for plant material (Ainsworth and Gillespie 2007) with caffeic acid (CA) as a standard was applied.

The TPC for each sample was determined using formula: $TPC = C \times V/M$, where: TPC is the total phenolic content ($\text{mg} \cdot \text{g}^{-1}$) of the extracts as CA equivalents, C is the concentration of CA established from the calibration curve ($\text{mg} \cdot \text{ml}^{-1}$), V is the volume of the extract solution (ml) and M is the weight of the sample (g). The assay consisted of one randomly sampled leaf from 4-week-old cucumber plants with three plants for each treatment. The assay was repeated three times.

Analysis of chlorophyll and carotenoids content

Chlorophylls and carotenoids are abundant pigments directly influencing cucumber growth through photosynthetic activities. In cucumber, chlorophylls are synthesized in green tissues, whereas carotenoids are synthesized in the leaves, flowers and fruits. In the cucumber leaf tissues carotenoids (mainly α -carotene and β -carotene) act as photoprotectors. The presence of the β -carotene and xanthophylls confer the characteristic yellow coloration to the flowers and to the fruit's meso- and endocarps (Hui Song *et al.* 2010). To determine the effects of fungal colonization on the accumulation of total chlorophyll (chlorophyll a and b) and carotenoids content in 7-week-old cucumber seedlings, leaves (0.5 g) were randomly sampled from three replicate plants of each treatment, chopped and immediately homogenized in 90% ethanol using mortar and pestle with gradual addition of 17.5 ml of ethanol. The homogenate was centrifuged at 1,500 rpm for 15 min. The absorbance of chlorophyll and carotenoid extracts was read after 90% aqueous ethanol was used as the blank, to zero the instrument initially and after

every wavelength resetting. Chlorophyll content was determined according to the method described by Barnes *et al.* (1992). Chlorophyll a and b absorbencies were measured at 664 and 649 nm, respectively, with a spectrophotometer (Thermospectronic) and calculated for chlorophyll concentration in fresh weight for three replicates. Formulas for chlorophyll content were as follows: $C_a = 13.36 \times A_{665} - 5.19 \times A_{649}$ and $C_b = 27.43 \times A_{649} - 8.12 \times A_{665}$, where: A_{665} = absorbance at a wavelength of 665 nm; A_{649} = absorbance at a wavelength of 649 nm. Carotenoids (mixture of α -carotenes and β -carotenes) absorbance was measured at 470 nm and carotenoid concentration was determined using the following formula: $C = (1,000A_{470} - 2.13 C_a - 97.63 C_b) / 209$, where A_{470} = absorbance at a wavelength of 470 nm (Braniša *et al.* 2014). The test was repeated three times.

Statistical analysis

All experiments were arranged in a complete randomized design. All data sets were analyzed with the statistical program IBM SPSS Statistics for Windows, Version 23.0 (2015) using one-way ANOVA after checking the assumptions for normality and the homogeneity of variance (Levene's test). Data sets from repeated tests for percentage values of fungal colonization with *M. anisopliae* and *B. bassiana* of different cucumber plant parts in addition to the aphid population size, TPCs, and chlorophyll contents were pooled together prior to one-way ANOVA analysis with the fungal strain as a main factor. A plant's physiological parameters were subjected to one-way ANOVA with the fungal strain as a main factor. When a significant F test was obtained at $p = 0.05$, separation of treatment means was performed using Duncan test.

Results

Molecular identification of microorganisms and aphids

Total DNA extracted from both fungal strains was PCR amplified by ITS4 and ITS5 primers. The amplification of the ITS region resulted in a single product (~ 600 bp) for both isolates. The ITS sequences of both isolates (BbL1, MaL1) were deposited in GenBank with the accession numbers. MT533246 and MT533250, respectively. The BLAST analysis of the ITS sequences of BbL1, MaL1 revealed 100% similarity with *B. bassiana* (accession No. KC753391) and 99.46% similarity with *M. anisopliae* (accession No. MG786739) in the NCBI database, respectively. This molecular identification confirmed the phenotypic identification of the fungal isolates used in this study.

Primer sets KBR (AG)-F and KBR (AG)-R successfully identified *A. gossypii* sp. by amplifying the expected fragment size of 600 bp in two samples only. None of the tested aphids was identified as *Myzus persicae*.

Assessment of colonization of *Beauveria bassiana* and *Metarhizium anisopliae* in cucumber plants

The efficiency of artificial inoculation of cucumber seeds was observed under scanning electron micrographs (SEM). Scanning electron micrographs revealed that

seeds of Beit Alpha hybrid were emarginate and elliptical with pointed ends. The seed sculpture of the studied taxa (*C. sativus*) showed that surfaces were smooth and covered with pits (hollows). The outer epidermal cells were of regular rectangular shapes separated with cell boundaries. Moreover, these micrographs showed that inoculated seed surfaces were covered with germinated and non-germinated conidia. All germinated spores were observed at different stages of development while their germ tubes spread over the seed's surfaces at different distances. Direct hyphal penetration through the epidermal cell wall was also observed (Fig. 1D)




Fig. 1. Scanning electron micrographs (SEM) of Tween (control), *M. anisopliae* and *B. bassiana* treated seeds 24 h post-inoculation. Micrograph A shows no fungal growth on control seed surface, whereas B shows *M. anisopliae* spores and hyphae adhered to the micropylar area showing ruptured covering layers (black arrows) and emergence of the radicle. Micrographs C and D of *M. anisopliae* treated seeds show attached conidia and conidial hyphae penetrating across cell boundary (black arrow), respectively. Micrographs E and F of *B. bassiana* treated seeds show hyphae growing on seed surface and directly penetrating the epidermal cell wall (black arrow), respectively

and F (black arrows) demonstrating the evidence of applied EPF becoming endophytic. The results of the number of attached conidia, germinated conidia, appressorium producing conidia and hyphae penetrating the seed cuticle are presented in Figure 2. No conidia were found on the surfaces of non-treated control seeds (Fig. 1A). The number of spores attached to the surface of *M. anisopliae* treated seeds was significantly higher ($13.2 \text{ conidia } 100 \mu\text{m}^{-2}$; $p < 0.0001$) than that on *B. bassiana* ($2.5 \text{ conidia } 100 \mu\text{m}^{-2}$) treated seeds (Fig. 1C and E). Moreover, a greater number of *M. anisopliae* spores adhered to the micropylar area of the examined seeds showing ruptured covering layers and emergence of radicles (Fig. 1B) than to the other parts of the same seed. In contrast, a more uniformed distribution of spores on cucumber seed surfaces was recorded on *B. bassiana* inoculated (Fig. 1E). However, not all attached spores germinated after 24 h and there was no difference in the number of germinated spores ($p = 0.014$), either penetrating ($p = 0.342$) or appressorium producing spores ($p = 0.38$) on the surfaces of treated seeds (Fig. 2). Moreover, approximately 11% and 13% of hyphae developed by *B. bassiana* and *M. anisopliae*, respectively, entered the seed through direct penetration. Appressoria were produced by 6.6% of the *Metarhizium* spores. There was no appressorium formation observed for *B. bassiana*.

All the fungal populations found in the peat compost were dominated by Ascomycetes, particularly




Fig. 2. Mean number ($100 \mu\text{m}^{-2}$) \pm SE of attached, germinated spores, appressorium and hyphae penetrating the seed epidermal cells of the treated seeds. Bars with different letters are significantly different at $p = 0.05$ (Duncan test, after one-way ANOVA)

by *Aspergillus*, *Penicillium*, and *Sclerotium* spp. The second most abundant group was the Mesomycetozoa (Protists), Basidiomycetes, and by a few unidentified yeasts. There were no *Metarhizium* or *Beauveria* spp. detected in the compost used for planting. Artificial inoculation of cucumber seeds carried out by their direct submersion in conidial suspensions provided successful colonization of plant parts with tested *B. bassiana* and *M. anisopliae* isolates. The presence of fungi in the internal tissues was confirmed by their recovery from roots, stems and leaf pieces on PDA selective medium. Results in Table 1 show that both, *B. bassiana* and *M. anisopliae*, isolates colonized cucumber seedlings. The most successful endophyte re-isolation frequency was observed from root tissues for both isolates (100%) ($p < 0.0001$). In addition, slightly higher percentage recoveries (50%, $p = 0.1$ and 25%, $p = 0.405$) were observed in stem and leaf tissues, respectively, colonized by *M. anisopliae*. However, no fungal recovery was achieved from stem and leaf tissues of *B. bassiana* inoculated plants. None of the leaf discs obtained from control plants showed signs of *M. anisopliae* or *B. bassiana* outgrowth, however, the control plants were observed to be colonized by naturally occurring endophytes such as *Penicillium*, *Aspergillus* as well as by other unidentified fungi.

The effects of endophytic EPFs on the population size of *Aphis gossypii* on cucumber plants

In general, there was a significant negative effect of cucumber plant colonization with both fungal entomopathogens on the population size of *A. gossypii* 5 days after exposure. The data (Fig. 3) show that the number of aphids in both treatments was significantly lower than that in the control plants ($p < 0.05$).

The aphid population size on *Metarhizium* colonized plants was reduced by almost 80 aphids per plant (35%) and on *Beauveria* colonized plants by 72 aphids (32%) (Fig. 3) resulting in healthier/fitter cucumber plants.

Table 1. Effects of conidial seed treatments on percentage recovery of *B. bassiana* and *M. anisopliae* from different parts of cucumber plants cultured outdoors in non-sterilized peat moss

Treatment	Leaves (% \pm SE)	Stem (% \pm SE)	Roots (% \pm SE)
Control	0.00 \pm 0.00 a*	0.00 \pm 0.00 a	0.00 \pm 0.00 a
<i>B. bassiana</i>	0.00 \pm 0.00 a	0.00 \pm 0.00 a	100 \pm 0.00 b
<i>M. anisopliae</i>	25 \pm 0.25 a	50 \pm 0.28 a	100 \pm 0.00 b

Medium 4 weeks post-inoculation

*means (% \pm SE) within a column followed by the same letter are not significantly different at $p = 0.05$ (Duncan test, after one-way ANOVA)




Fig. 3. Mean (\pm SE) number of aphids per cucumber plant inoculated with *B. bassiana*, *M. anisopliae* and Tween (control). Bars with different letters are significantly different at $p = 0.05$ (Duncan test, after one-way ANOVA)

The effects of endophytic EPFs on physiological and biochemical parameters of cucumber plants

Physiological parameters

The number of fully developed leaves and flowers was counted for each replicate and treatment 6 weeks post-inoculation. The number of small cucumber fruits and the plant's height were recorded for each potted plant 7 weeks post inoculation. The results show that there was a significant difference in the number of flowers ($p = 0.05$), leaves ($p < 0.005$) and the height of the plants ($p < 0.01$) colonized with tested fungal endophytes compared to uncolonized control plants (Fig. 4). On one hand, plants colonized with *Beauveria* isolate produced twice as many flowers and two times more leaves than control plants (Fig. 3). On the other hand, *Metarhizium* colonized plants were almost 12 cm taller and had two more developed cucumber fruits than control plants (Fig. 4).

Biochemical parameters

To see if the suggestion of Pańka et al. (2013) that endophytes could indirectly influence the defense reaction in the host plant by synthesizing more phenolic compounds than plants without endophytes, the total phenolic content was determined for 2-week-old seedlings. The results presented in Table 2 show that cucumber plants reacted to endophytic colonization by increased induction of soluble phenols. Thus, plants grown from treated seeds with *B. bassiana* accumulated a significantly ($p < 0.0001$) higher amount of TPC ($182 \text{ mg} \cdot \text{g}^{-1}$) than those plants treated with *M. anisopliae* ($101 \text{ mg} \cdot \text{g}^{-1}$) or Tween ($95 \text{ mg} \cdot \text{g}^{-1}$).

However, analyses of Ca, Cb and carotenoid content in the leaf blades of 7-week-old cucumber plants showed no effect of endophytic EPFs on measured parameters (Fig. 5). The leaf extracts from colonized

Table 2. Effects of endophytic entomopathogenic fungi (EPFs) on total phenolic compounds in 4-week-old plants grown from seeds treated with *B. bassiana*, *M. anisopliae* and Tween (control)

Treatment	Total phenol content [$\mu\text{g} \cdot \text{g}^{-1}$]
Control	$94.5 \pm 0.003 \text{ a}^*$
<i>M. anisopliae</i>	$101.3 \pm 0.003 \text{ a}$
<i>B. bassiana</i>	$181.8 \pm 0.001 \text{ b}$

*means ($\mu\text{g} \cdot \text{g}^{-1} \pm \text{SE}$) within a column followed by the same letter are not significantly different at $p = 0.05$ (Duncan test, after one-way ANOVA)




Fig. 4. Mean height (cm \pm SE) and the mean (\pm SE) number (N) of flowers, leaves and fruits of cucumber plants inoculated with *B. bassiana*, *M. anisopliae* and Tween (control). Bars with different letters are significantly different at $p = 0.05$ (Duncan test, after one-way ANOVA)




Fig. 5. Effect of fungal inoculation on chlorophylls Ca and Cb and carotenoids content (mean concentration, $\mu\text{g} \cdot \text{ml} \pm \text{SE}$) in 7-week-old cucumber plants. Bars with similar letters do not differ significantly at $p = 0.05$ (Duncan test, after one-way ANOVA)

and uncolonized cucumber tissues contained similar amounts of both chlorophylls ($p = 0.785$ and $p = 0.325$, respectively) and carotenoids ($p = 0.593$).

Discussion

The main aims of this study were to determine if *M. anisopliae* and *B. bassiana* can establish an endophytic relationship with cucumber plants under naturally fluctuating conditions in non-sterile peat compost, and if these fungi can act as plant growth promoters, or if inoculation of seeds with EPF affects the aphid, *A. gossypii*. Seed inoculation was shown to be very successful for EPF colonizing many plant tissues (Kabaluk and Ericsson 2007; Ownley *et al.* 2008; Powell *et al.* 2009; Akello and Sikora 2012; Akutse *et al.* 2013; Lopez *et al.* 2014; Russo *et al.* 2015) including cucumber seedlings (Shaalán and Ibrahim 2019; Rajab *et al.* 2020). The success of internal colonization is usually determined by re-isolation of tested fungi using the culture dependent method (CDM) (McKinnon *et al.* 2017). However, some other techniques such as molecular identification and visualization of endophytes within plant tissues using dyes, histological techniques, immunofluorescence, light and electron microscopy have also been shown to be reliable for endophyte detection (McKinnon *et al.* 2017; Ullrich *et al.* 2017; Vega 2018).

Seed submersion used in the current study resulted in the attachment of conidia to seed surfaces, their germination, hyphal growth across seed surfaces and direct penetration through epidermal cell walls 24 h post inoculation. Similarly, germinated conidia of *B. bassiana* produced hyphae that penetrated the leaf cuticle directly or grew through stomates into the corn leaf (Wagner and Lewis 2000). When a spore lands on the plant surface, it perceives the immediate environment, the topography of the host cuticle and nutrient availability, which are the essential factors that trigger spore attachment and stimulate germination (Tucker and Talbot 2001). Indeed, submerging cucumber seeds in spore suspension prompted an imbibition of water, which, in turn, caused a rupture of seed covering layers exposing micropylar endosperms and emerging radicles. Therefore, a germinating cucumber seed was likely to provide a perfect humid environment and accessible nutrients for conidia to adhere to and germinate. It is interesting that while *M. anisopliae* conidia in this study preferred to aggregate and attach near the micropylar area of radicle emergence, conidia of *B. bassiana* spread homogeneously across the entire seed surface. A previous study indicated that in the environment *Metarhizium* spp. appear to be restricted to the rhizosphere (Behie *et al.* 2015). Also, initial stages of plant-*Metarhizium* interaction through seed treatment would involve extensive rhizoplane colonization followed by endophytic growth (Barelli *et al.* 2018). The results of SEM observations also revealed that

around 13% of germinated spores of both spp. penetrated seeds through epidermal cells directly without using epidermal pits. This rate of initial seed colonization, perhaps, was enough to set off a mechanism for systemic colonization of the whole plant (Wagner and Lewis 2000).

Indeed, the fungal recovery from 4-week-old cucumber seedlings grown on non-sterile compost was achieved for both entomopathogens. *Beauveria bassiana* was isolated only from roots whereas *M. anisopliae* was isolated from roots, stems and leaves. In the previous study, however, the same fungal isolates were detected in all plant parts obtained from 10-day-old cucumber seedlings cultivated on sterile substrate (Shaalán and Ibrahim 2019). Likewise, in the study of Tefera and Vidal (2009), colonization was not recorded in stems and leaves of *Beauveria* seed-treated seedlings grown on non-sterile soil but occurred in vermiculite and sterile soil. The fungistatic effects of soil (Lingg and Donaldson 1981), biotic antagonism (Pereira *et al.* 1993) or microbial competition in the compost may have prevented or delayed *B. bassiana* from reaching the stem and leaves. On the other hand, the initial low number of attached spores (Fig. 2) as well as the transient nature of *B. bassiana* (Russo *et al.* 2015) could explain negative endophytism with cucumber stems and leaves. The application rate of conidia is also known to influence the rate of endophytic colonization (Ownley *et al.* 2008).

In addition, this study not only reports the development of entomopathogenic fungi introduced endophytically but also shows that endophytic presence in cucumber plant tissues can increase the resistance against melon aphids, *A. gossypii*. The results of the present work support previous studies where *B. bassiana* endophyte also negatively affected cotton aphid reproduction (Castillo-Lopez *et al.* 2014; González-Mas *et al.* 2019), but are in opposition to endophytic *B. bassiana* effects on other aphid species (Clifton *et al.* 2018; Jensen *et al.* 2019). The mechanisms by which herbivores can be negatively affected by clavicipitaceous obligate endophytes have been studied in a few different grass species and can vary from antixenosis and/or antibiosis mediated by constitutive production and/or induction of secondary compounds produced by the plant (Clay *et al.* 1993; Clay 1996; Carriere *et al.* 1998) or secondary metabolites produced by the endophytes themselves (Jaber and Vidal 2010; Gurunlingappa *et al.* 2010; Saari *et al.* 2010; Vega 2018). Another hypothesis for the mechanism by which endophytes can negatively affect herbivores is based on the idea that endophytes can alter the phytosterol profiles of plants and compete with insects for these compounds which are essential for their development (Dugassa-Gobena *et al.* 1996; Raps and Vidal 1998). It was also

suggested that plant responses to endophyte invasion may involve e.g., synthesis of phenolic compounds (Pańka *et al.* 2013). The presence of *Neotyphodium lolii* endophyte increased significantly the production of total phenolics in all ryegrass genotypes. In general, phenolic compounds and some reactive oxygen species are synthesized in plants partly as a response to ecological and physiological pressures (Ibrahim *et al.* 2001; Chung *et al.* 2003; Schulz and Boyle 2005; Diaz *et al.* 2010) and possess anti-herbivore properties (Fürstenberg-Hägg *et al.* 2013; Vega 2018). Total phenolic content analysis showed that *B. bassiana* and *M. anisopliae* endophytes colonizing 4-week-old cucumber seedlings influenced the defense reaction in the cucumber plant by synthesizing more phenolic compounds than seedlings without an endophyte. Such a response could have rendered plants more capable of self-defense against initial attack by *A. gossypii*. The mechanisms of plant defense responses to fungal colonization could be species-dependent, meaning that the same plants could react differently to different fungal endophytes. When the roots-inoculated tomato plants with *B. bassiana* and *Trichoderma koningiopsis* were analyzed by GC-MS and compared to non-inoculated plants, it was found that all successfully colonized roots, stems and leaf tissues contained modified tomato volatile organic compounds (VOCs) with significant differences between isolates (Pineda *et al.* 2010). However, the profiles of cucumber VOCs were not determined in colonized or non-colonized tissues for this study, and the hypothesis of modifications to the chemical signals produced by the plant (Vega 2018; Gange *et al.* 2019) remains to be tested as an explanation of the data in the current study.

Recent reviews by McKinnon *et al.* (2017), Jaber and Enkerli (2017) and Vega (2018) presented examples of many studies reporting the effects of endophytism on plant growth and development. It was evident that some plants were unresponsive (neutral) to endophytic colonization. Some responded with a positive outcome (e.g., improved growth (Vega *et al.* 2009; Liao *et al.* 2014; Jaber and Enkerli 2016) and increased yields (Maniania *et al.* 2003; Kabaluk and Ericsson 2007)), while a very few exhibited a negative response (e.g., alleviated iron chlorosis symptoms (Sánchez-Rodríguez *et al.* 2015)). Colonization of cucumber plants by *B. bassiana* and *M. anisopliae* in our study resulted in improved plant growth, greater numbers of leaves and flowers without significant effects on chlorophyll synthesis. It is possible that since seed treatment with these two isolates can significantly improve seed germination rates and strengthen root systems of resultant seedlings (Shaalan and Ibrahim 2019), this may explain the subsequent improved plant growth. In addition, the ability of entomopathogenic endophytes

to produce plant growth regulators (e.g., indole-3-acetic acid (IAA)) (Liao *et al.* 2017) as well as to improve nutrient absorption (Sasan and Bidochka 2012) may indicate a high potential of the isolates used in this study to promote plant growth (Vega *et al.* 2009).

Conclusions

Colonization of cucumber plant tissues by entomopathogenic endophytes, *B. bassiana* and *M. anisopliae*, began with direct hyphal penetration of the seed surface. Conidia of *M. anisopliae* preferred the micropylar area of cucumber seeds for their initial attachment and penetration, while conidia of *B. bassiana* spread continually across the entire seed surface. Colonization of cucumber plants occurred under natural environmental conditions and on a non-sterile substrate, greatly promoting plant growth and development. Here, we showed that cucumber plant colonization by fungal entomopathogens is generally negative to aphids, *A. gossypii*, reducing their overall population size. Activation of multiple mechanisms related to plant-mediated systemic resistance could suppress the invasion of piercing-sucking insects and, thus, improve quality and quantity of agricultural crops. Since cucumber growers regularly purchase plantlets from plant nurseries, it would be viable to commercialize the production of *B. bassiana* and *M. anisopliae* pre-colonized plants.

Acknowledgements

The authors wish to thank the National Council for Scientific Research (CNRS) and the Lebanese University for funding this study. We also appreciate the assistance of the plant protection team at the Lebanese Agricultural Institute (LARI), including Eng. C. Saab and Z. Moussa. We also wish to thank Dr. R. Habchi (Research Platform for Nanoscience and Nanotechnology, Fanar Campus 90656, Lebanese University) for helping with SEM work.

References

- Ainsworth E.A., Gillespie K.M. 2007. Estimation of total phenolic content and other oxidation substrates in plant tissues using Folin-Ciocalteu reagent. *Nature Protocols* 2: 875–877. DOI: 10.1038/nprot.2007.102
- Akello J., Sikora R. 2012. Systemic acropetal influence of endophyte seed treatment on *Acyrtosiphon pisum* and *Aphis fabae* offspring development and reproductive fitness. *Biological Control* 61: 215–221. DOI: 10.1016/J.BIOCONTROL.2012.02.007
- Akutse K., Maniania N., Fiaboe K., Van Den Berg J., Ekesi S. 2013. Endophytic colonization of *Vicia faba* and *Phaseolus vulgaris* (Fabaceae) by fungal pathogens and their effects on the life-history parameters of *Liriomyza huidobrensis*

- (Diptera: Agromyzidae). *Fungal Ecology* 6: 293–301. DOI: <https://doi.org/10.1016/j.funeco.2013.01.003>
- Bajan C., Tyrawska D., Popowska-Nowak E., Bienkowski P. 1998. Biological response of *Beauveria bassiana* strains to heavy metal pollution and their accumulative ability. *Ecological Chemistry and Engineering* 5 (8–9): 676–685.
- Bamisile B.S., Dash C.K., Akutse K.S., Keppan R., Afolabi O.G., Hussain M., Qasim M., Wang L. 2018. Prospects of endophytic fungal entomopathogens as biocontrol and plant growth promoting agents: An insight on how artificial inoculation methods affect endophytic colonization of host plants. *Microbiological Research* 217: 34–50. DOI: <https://doi.org/10.1016/j.micres.2018.08.016>
- Barelli L., Moreira C.C., Bidochka M.J. 2018. Initial stages of endophytic colonization by *Metarhizium* involves rhizoplane colonization. *Microbiology* 164: 1531–1540. DOI: [10.1099/mic.0.000729](https://doi.org/10.1099/mic.0.000729).
- Barnes J.D., Balaguer L., Manrique E., Elvira S., Davison A.W. 1992. A reappraisal of the use of DMSO for the extraction and determination of chlorophylls a and b in lichens and higher plants. *Environment* 32: 83–100. DOI: [http://dx.doi.org/10.1016/0098-8472\(92\)90034-Y](http://dx.doi.org/10.1016/0098-8472(92)90034-Y)
- Behie S.W., Jones S.J., Bidochka M.J. 2015. Plant tissue localization of the endophytic insect pathogenic fungi *Metarhizium* and *Beauveria*. *Fungal Ecology* 13: 112–119. DOI: [10.1016/j.funeco.2014.08.001](https://doi.org/10.1016/j.funeco.2014.08.001)
- Blackman R.L. 2010. Aphids – Aphidinae (Macrosiphini). *Handbook for the Identification of British Insects* 2 (7): 1–413.
- Blackman R.L., Eastop V.F. 2000. Aphids on the world's crops: An identification and information guide. John Wiley and Sons, Chichester, UK. XF2006251066.
- Braniša J., Jenišová Z., Porubská M., Jomová K., Valko M. 2014. Spectrophotometric determination of chlorophylls and carotenoids. An effect of sonication and sample processing. *Journal of Microbiology, Biotechnology and Food Sciences* 3 (2): 61–64.
- Carriere Y., Bouchard A., Bourassa S., Brodeur J. 1998. Effect of endophyte incidence in perennial ryegrass on distribution, host choice, and performance of the hairy chinch bug (Hemiptera: Lygaeidae). *Economic Entomology* 91: 324–328. DOI: <https://doi.org/10.1093/jee/91.1.324>
- Castillo-Lopez D., Zhu-Salzman K., Ek-Ramos M.J., Sword G.A. 2014. The entomopathogenic fungal endophytes *Purpureocillium lilacinum* (formerly *Paecilomyces lilacinus*) and *Beauveria bassiana* negatively affect cotton aphid reproduction under both greenhouse and field conditions. *PLoS ONE* 9 (8): e103891. DOI: [10.1371/journal.pone.0103891](https://doi.org/10.1371/journal.pone.0103891)
- Chung I.M., Park Chun J.C., Yun S.J. 2003. Resveratrol accumulation and resveratrol synthase gene expression in response to abiotic stresses and hormones in peanut plants. *Plant Science* 164: 103–109. DOI: [10.1016/S0168-9452\(02\)00341-2](https://doi.org/10.1016/S0168-9452(02)00341-2)
- Clay K. 1996. Interactions among fungal endophytes, grasses and herbivores. *Researches on Population Ecology* 38 (2): 191–201. DOI: <https://doi.org/10.1007/BF02515727>
- Clay K., Marks S., Cheplick G.P. 1993. Effects of insect herbivory and fungal endophyte infection on competitive interactions among grasses. *Ecology* 74 (6): 1767–1777. DOI: <https://doi.org/10.2307/1939935>
- Clifton E.H., Jaronski S.T., Coates B.S., Hodgson E.W., Gasmann A.J. 2018. Effects of endophytic entomopathogenic fungi on soybean aphid and identification of *Metarhizium* isolates from agricultural fields. *PLoS ONE* 13 (3): e0194815. DOI: <https://doi.org/10.1371/journal.pone.0194815>
- Daayf F., Ongena M., Boulanger R., El Hadrami I., Bélanger R.R. 2000. Induction of phenolic compounds in two cultivars of cucumber by treatment of healthy and powdery mildew infected plants with extracts of *Reynoutria sachalinensis*. *Journal of Chemical Ecology* 26 (7): 1579–1593. DOI: <https://doi.org/10.1023/A:1005578510954>
- Diaz Napal G.N., Defago M., Valladares G., Palacios S. 2010. Response of *Epilachna paenulata* to two flavonoids, pinocembrin and quercetin, in a comparative study. *Journal of Chemical Ecology* 36 (8): 898–904. DOI: [10.1007/s10886-010-9823-1](https://doi.org/10.1007/s10886-010-9823-1)
- Dugassa-Gobena D., Raps A., Vidal S. 1996. Einu von *Acremonium strictum* auf den Sterolhaushalt von Panzen: ein möglicher Faktor zum veränderten Verhalten von Herbivoren. *Mitteilungen aus der Biologischen Bundesanstalt für Land- und Forstwirtschaft* 321: 299.
- Fürstenberg-Hägg J., Zagrobelny M., Bak S. 2013. Plant defense against insect herbivores. *International Journal of Molecular Sciences* 14 (5): 10242–10297. DOI: [10.3390/ijms140510242](https://doi.org/10.3390/ijms140510242)
- Gange A.C., Koricheva J., Currie A.F., Jaber L.R., Vidal S. 2019. Meta-analysis of the role of entomopathogenic and unspecialized fungal endophytes as plant bodyguards. *New Phytologist* 223 (4): 2002–2010. DOI: <https://doi.org/10.1111/nph.15859>
- Gawel N.J., Jarret R.L. 1991. A modified CTAB DNA extraction procedure for musa and ipomoea. *Plant Molecular Biology Reporter* 9 (3): 262–266. DOI: [10.1007/BF02672076](https://doi.org/10.1007/BF02672076)
- Gissella M.V., David B.O., James R.B. 2006. Efficacy assessment of *Aphidius colemani* (Hymenoptera: Braconidae) for suppression of *Aphis gossypii* (Homoptera: Aphididae) in greenhouse-grown chrysanthemum. *Economic Entomology* 99 (4): 1104–1111. DOI: [10.1603/0022-0493-99.4.1104](https://doi.org/10.1603/0022-0493-99.4.1104)
- Godfrey L.D., Rosenheim J.A., Goodell P.B. 2000. Cotton aphid emerges as major pest in SVJ cotton. *California Agriculture* 54 (6): 32–34. DOI: [10.3733/ca.v054n06p26](https://doi.org/10.3733/ca.v054n06p26)
- González-Mas N., Sánchez-Ortiz A., Valverde-García P., Quesada-Moraga E. 2019. Effects of endophytic entomopathogenic ascomycetes on the life-history traits of *Aphis gossypii* Glover and its interactions with melon plants. *Insects* 10 (6): 165. DOI: [10.3390/insects10060165](https://doi.org/10.3390/insects10060165)
- Grace S.C., Logan B.A. 2000. Energy dissipation and radical scavenging by the plant phenylpropanoid pathway. *Philosophical Transactions: Biological Sciences* 355 (1402): 1499–1510. DOI: [10.1098/rstb.2000.0710](https://doi.org/10.1098/rstb.2000.0710)
- Gurulingappa P., McGee P.A., Sword G. 2011. Endophytic *Leucanicillium lecnii* and *Beauveria bassiana* reduce the survival and fecundity of *Aphis gossypii* following contact with conidia and secondary metabolites. *Crop Protection* 30 (3): 349–353. DOI: [10.1016/j.cropro.2010.11.017](https://doi.org/10.1016/j.cropro.2010.11.017)
- Gurulingappa P., Sword G.A., Murdoch G., McGee P.A. 2010. Colonization of crop plants by fungal entomopathogens and their effects on two insect pests when in planta. *BioControl* 55 (1): 34–41. DOI: <https://doi.org/10.1016/j.biocontrol.2010.06.011>
- Hermoso de Mendoza A., Belliure B., Carbonell E.A., Real V. 2001. Economic thresholds for *Aphis gossypii* (Hemiptera: Aphididae) on *Citrus clementina*. *Journal of Economic Entomology* 94 (2): 439–444. DOI: <https://doi.org/10.1603/0022-0493-94.2.439>
- Humber R.A. 1997. Fungi: identification. p. 153–185. In: “Manual of Techniques in Insect Pathology” (L.A. Lacey, ed.). Academic Press, London. DOI: <https://doi.org/10.1016/B978-012432555-5/50011-7>
- Ibrahim L., Hamieh A., Ghanem H., Ibrahim S. 2011. Pathogenicity of entomopathogenic fungi from Lebanese soils against aphids, whitefly and non-target beneficial insects. *International Journal of Agriculture Sciences* 3 (3): 156–164. DOI: [10.9735/0975-3710.3.3.156-164](https://doi.org/10.9735/0975-3710.3.3.156-164)
- Ibrahim L., Laham L., Tomma A., Ibrahim S. 2015. Mass production, yield, quality, formulation and efficacy of entomopathogenic *Metarhizium anisopliae* conidia. *British Journal of Applied Sciences and Technology* 9 (5): 427–440. DOI: [10.9734/bjast/2015/17882](https://doi.org/10.9734/bjast/2015/17882)
- Ibrahim L., Spackman V.M.T., Cobb A.H. 2001. An investigation of wound healing in sugar beet roots using light and fluorescence microscopy. *Annals of Botany* 88 (2): 313–320. DOI: <https://doi.org/10.1006/anbo.2001.1461>
- Jaber L.R., Enkerli J. 2016. Effect of seed treatment duration on growth and colonization of *Vicia faba* by endophytic *Beauveria bassiana* and *Metarhizium brunneum*. *Biologi-*

- cal Control 103: 187–195. DOI: <https://doi.org/10.1016/j.biocontrol.2016.09.008>
- Jaber L.R., Enkerli J. 2017. Fungal entomopathogens as endophytes: can they promote plant growth? *Biocontrol Science and Technology* 27 (1): 28–41. DOI: <https://doi.org/10.1080/09583157.2016.1243227>
- Jaber L.R., Vidal S. 2010. Fungal endophyte negative effects on herbivory are enhanced on intact plants and maintained in a subsequent generation. *Ecological Entomology* 35 (1): 25–36. DOI: <https://doi.org/10.1111/j.1365-2311.2009.01152.x>
- Jensen R.E., Enkegaard A., Steenberg T. 2019. Increased fecundity of *Aphis fabae* on *Vicia faba* plants following seed or leaf inoculation with the entomopathogenic fungus *Beauveria bassiana*. *PLoS ONE* 14 (10): 1–13, e0223616. DOI: <https://doi.org/10.1371/journal.pone.0223616>
- Kabaluk J.T., Ericsson J.D. 2007. *Metarhizium anisopliae* seed treatment increases yield of field corn when applied for wireworm control. *Agronomy Journal* 99 (5): 1377–1381. DOI: 10.2134/agronj2007.0017N
- Kumar P.K.R., Hemanth G., Niharika P.S., Kolli S.K. 2015. Isolation and identification of soil mycoflora in agricultural fields at Tekkali Mandal in Srikakulam District. *International Journal of Advances in Pharmacy, Biology and Chemistry* 4 (2): 484–490.
- Lacey L.A. 2016. Entomopathogens used as microbial control agents. p. 3–12. In: “Microbial Control of Insect and Mite Pests” (L.A. Lacey, ed.). Amsterdam: Elsevier/Academic Press. DOI: <https://doi.org/10.1016/B978-0-12-803527-6.00001-9>
- Lee S.B., Milgroom M.G., Taylor J.W. 1988. A rapid, high yield mini-prep method for isolation of total genomic DNA from fungi. *Fungal Genetics Newsletter* 35: 23–24. DOI: <https://doi.org/10.4148/1941-4765.1531>
- Liao J., Li X., Wong T.Y., Wang J.J., Khor C.C., Tai E.S., Aung T., Teo Y.Y., Cheng C.Y. 2014. Impact of measurement error on testing genetic association with quantitative traits. *PLoS ONE* 9 (1): e87044. DOI: <https://doi.org/10.1371/journal.pone.0087044>
- Liao X., Lovett B., Fang W., St. Leger R.J. 2017. *Metarhizium robertsii* produces indole-3-acetic acid, which promotes root growth in *Arabidopsis* and enhances virulence to insects. *Microbiology* 163 (7): 980–991. DOI: 10.1099/mic.0.000494
- Lingg A.J., Donaldson M.D. 1981. Biotic and abiotic factors affecting stability of *Beauveria bassiana* conidia in soil. *Journal of Invertebrate Pathology* 38 (2): 191–200. DOI: [https://doi.org/10.1016/0022-2011\(81\)90122-1](https://doi.org/10.1016/0022-2011(81)90122-1)
- Lopez D.C., Zhu-Salzman K., Ek-Ramos M.J., Sword G.A. 2014. The entomopathogenic fungal endophytes *Purpureocillium lilacinum* (formerly *Paecilomyces lilacinus*) and *Beauveria bassiana* negatively affect cotton aphid reproduction under both greenhouse and field conditions. *PLoS One* 9: e103891. DOI: 10.1371/journal.pone.0104342
- Maniania N.K., Sithanatham S., Ekesi S., Ampong-Nyarko K., Baumgärtner J., Löhr B., Matoka C.M. 2003. A field trial of the entomogenous fungus *Metarhizium anisopliae* for control of onion thrips, *Thrips tabaci*. *Crop Protection* 22 (3): 553–559. DOI: [https://doi.org/10.1016/S0261-2194\(02\)00221-1](https://doi.org/10.1016/S0261-2194(02)00221-1)
- Matallanas B., Lantero E., M'Saad M., Callejas C., Ochando M.D. 2013. Genetic polymorphism at the cytochrome oxidase I gene in mediterranean populations of *Batrocera oleae* (Diptera: Tephritidae). *Applied Entomology* 137 (8): 624–630. DOI: <https://doi.org/10.1111/jen.12037>
- McKinnon A.C., Saari S., Moran-Diez M.E., Meyling N.V., Raad M., Glare T.R. 2017. *Beauveria bassiana* as an endophyte: a critical review on associated methodology and biocontrol potential. *BioControl* 62: 1–17. DOI: 10.1007/s10526-016-9769-5.
- Omkar P.A. 2004. Predaceous coccinellids in India: predator-prey catalogue. *Oriental Insects* 38 (1): 27–61. DOI: 10.1080/00305316.2004.10417373
- Ownley B.H., Dee M.M., Gwinn K. 2008. Effect of conidial seed treatment rate of entomopathogenic *Beauveria bassiana* 11-98 on endophytic colonization of tomato seedlings and control of *Rhizoctonia* disease. *Phytopathology* 98 (6): S118–S118.
- Pańka D., Piesik D., Jeske M., Musiał N., Koczwara K. 2013. Production of phenolic compounds by perennial ryegrass (*Lolium perenne* L.)/*Neotyphodium lolii* association as a defense reaction towards infection by *Fusarium poae* and *Rhizoctonia solani*. p. 124–125. In: “Endophytes for Plant Protection: the State of the Art” (C. Schneider, C. Leifert, F. Feldmann, eds.). Deutsche Phytomedizinische Gesellschaft, Braunschweig.
- Pathan A.K., Bond J., Gaskin R.E. 2010. Sample preparation for SEM of plant surfaces. *Materials Today* 12 (1): 32–43. DOI: 10.1016/S1369-7021(10)70143-7
- Perea-Domínguez X.P., Hernández-Gastelum L.Z., Olivás-Olguin H.R., Espinosa-Alonso L.G., Valdez-Morale M., Medina-Godoy S. 2018. Phenolic composition of tomato varieties and an industrial tomato by-product: free, conjugated and bound phenolics and antioxidant activity. *Journal of Food Science and Technology* 55 (9): 3453–3461. DOI: <https://doi.org/10.1007/s13197-018-3269-9>
- Pereira R.M., Stimac J.L., Alves S.B. 1993. Soil antagonism affecting the dose – response of workers of the red imported fire ant, *Solenopsis invicta*, to *Beauveria bassiana* conidia. *Journal of Invertebrate Pathology* 61 (2): 156–161. DOI: <https://doi.org/10.1006/jipa.1993.1028>
- Petrini O., Fisher P.J. 1987. Fungal endophytes in *Salicornia perennis*. *Transactions of the British Mycological Society* 87 (4): 647–651. DOI: [https://doi.org/10.1016/S0007-1536\(86\)80109-7](https://doi.org/10.1016/S0007-1536(86)80109-7)
- Pineda A., Zheng S.-J., van Loon J.J.A., Pieterse C.M.J., Dicke M. 2010. Helping plants to deal with insects: the role of beneficial soil-borne microbes. *Trends in Plant Science* 15 (9): 507–514. DOI: <https://doi.org/10.1016/j.tplants.2010.05.007>
- Powell W.A., Klingeman W.E., Ownley B.H., Gwinn K.D. 2009. Evidence of endophytic *Beauveria bassiana* in seed-treated tomato plants acting as a systemic entomopathogen to larval *Helicoverpa zea* (Lepidoptera: Noctuidae). *Journal of Entomological Science* 44 (4): 391–396. DOI: <https://doi.org/10.18474/0749-8004-44.4.391>
- Rajab L., Ahmad M., Gazal I. 2020. Endophytic establishment of the fungal entomopathogen, *Beauveria bassiana* (Bals.) Vuil., in cucumber plants. *Egyptian Journal of Biological Pest Control* 30: 143. DOI: <https://doi.org/10.1186/s41938-020-00344-8>
- Raps A., Vidal S. 1998. Indirect effects of an unspecialized endophytic fungus on specialized plant-herbivorous insect interactions. *Oecologia* 114 (4): 541–547. DOI: 10.1007/s004420050478
- Rebijith K.B., Asoka R., Krishna V., Krishna Kumar N.K., Ramamurth V.V. 2012. Development of species-specific markers and molecular differences in mitochondrial and nuclear DNA sequences of *Aphis gossypii* and *Myzus persicae* (Hemiptera: Aphididae). *Florida Entomologist* 95 (3): 674–682. DOI: 10.1653/024.095.0318
- Rodriguez R.J., White Jr. J.F., Arnold A.E., Redman R.S. 2009. Fungal endophytes: Diversity and functional roles. *New Phytologist* 182 (2): 314–330. DOI: <http://dx.doi.org/10.1111/j.1469-8137.2009.02773.x>
- Russo M.L., Pelizza S.A., Cabello M.N., Stenglein S.A., Scorsetti A.C. 2015. Endophytic colonisation of tobacco, corn, wheat and soybeans by the fungal entomopathogen *Beauveria bassiana* (Ascomycota, Hypocreales). *Biocontrol Science and Technology* 25 (4): 475–480. DOI: 10.1080/09583157.2014.982511
- Saari S., Helander M., Faeth S.H. 2010. The effects of endophytes on seed production and seed predation of tall fescue and meadow fescue. *Microbial Ecology* 60: 928–934. DOI: <https://doi.org/10.1007/s00248-010-9749-8>
- Saikkonen K., Lehtonen P., Helander M., Koricgeva J., Faeth S.H. 2006. Model systems in ecology: dissecting the endophyte-

- grass literature. *Trends in Plant Science* 11 (9): 428–433. DOI: 10.1016/j.tplants.2006.07.001
- Sánchez-Rodríguez A.R., Del Campillo M.C., Quesada-Moraga E. 2015. *Beauveria bassiana*: An entomopathogenic fungus alleviates Fe chlorosis symptoms in plants grown on calcareous substrates. *Scientia Horticulturae* 197: 193–202. DOI: <https://doi.org/10.1016/j.scienta.2015.09.029>
- Sasan R.K., Bidochka M.J. 2012. The insect-pathogenic fungus *Metarhizium robertsii* (Clavicipitaceae) is also an endophyte that stimulates plant root development. *American Journal of Botany* 99 (1): 101–107. DOI: 10.3732/ajb.1100136
- Schulz B., Boyle C. 2005. Fungal endophyte continuum. *Mycological Research* 109 (6): 661–686. DOI: <https://doi.org/10.1017/S095375620500273X>
- Shalan R., Ibrahim L. 2019. Entomopathogenic fungal endophytes: can they colonize cucumber plants? p. 853–860. In: “Book of Proceedings of the IX International Scientific Agriculture Symposium AGROSYM 2018”. 04–07 October 2018, Bosnia and Herzegovina.
- Shi X.G., Zhu Y.K., Xia X.M., Qiao K., Wang H.Y., Wang K.Y. 2012. The mutation in nicotinic acetylcholine receptor $\beta 1$ subunit may confer resistance to imidacloprid in *Aphis gossypii* (Glover). *Journal of Food, Agriculture and Environment* 10 (2): 1227–1230.
- Skinner M., Parker B.L., Kim J.S. 2014. Role of entomopathogenic fungi in integrated pest management. p. 169–191. In: “Integrated Pest Management: Current Concepts and Ecological Perspectives” (D.P. Abrol, ed.). Academic Press, San Diego. DOI: <https://doi.org/10.1016/B978-0-12-398529-3.00011-7>
- Song H., Chen J., Staub J., Simon P. 2010. QTL analyses of orange color and carotenoid content and mapping of carotenoid biosynthesis genes in cucumber (*Cucumis sativus* L.). *Acta Horticulturae* 871: 607–614. DOI: 10.17660/ACTAHORTIC.2010.871.83
- Stoetzel M.B., Miller G.L., O'Brien P.J., Graves J.B. 1996. Aphids (Homoptera: Aphididae) colonizing cotton in the United States. *Florida Entomologist* 79 (2): 193–205. DOI: 10.2307/3495817
- Tefera T., Vidal S. 2009. Effect of inoculation method and plant growth medium on endophytic colonization of sorghum by the entomopathogenic fungus *Beauveria bassiana*. *BioControl* 54: 663–669. DOI: 10.1007/s10526-009-9216-y
- Tucker S.L., Talbot N.J. 2001. Surface attachment and pre-penetration stage development by plant pathogenic fungi. *Annual Review of Phytopathology* 39: 385–417. DOI: 10.1146/annurev.phyto.39.1.385
- Ullrich C.I., Koch E., Matecki C., Schäfer J., Burkl T., Rabenstein F., Kleespies R.G. 2017. Detection and growth of endophytic entomopathogenic fungi in dicot crop plants. *Journal für Kulturpflanzen* 69 (9): 291–302. DOI: 10.1399/JfK.2017.09.02
- Umboh S.D., Salaki C.L., Tulung M., Mandey L.C., Maramis R.T.D. 2016. The isolation and identification of fungi from the soil in gardens of cabbage were contaminated with pesticide residues in subdistrict Modinding. *International Journal of Research in Engineering and Science* 4 (7): 25–32.
- Vandre W. 2013. Cucumber production in greenhouses. University of Alaska Fairbanks Cooperative Extension Service. Available on: www.uaf.edu/ces
- Vega F.E. 2018. The use of fungal entomopathogens as endophytes in biological control: a review. *Mycologia* 110 (1): 4–30. DOI: <https://doi.org/10.1080/00275514.2017.1418578>
- Vega F.E., Goettel M.S., Blackwell M., Chandler D., Jackson M.A., Keller S., Koike M., Maniania N.K., Monzón A., Ownley B.H. et al. 2009. Fungal entomopathogens: New insights on their ecology. *Fungal Ecology* 2 (4): 149–159. DOI: <https://doi.org/10.1016/j.funeco.2009.05.001>
- Vega F.E., Meyling N.V., Luangsa-Ard J.J., Blackwell M. 2012. Fungal entomopathogens. p. 171–220. In: “Insect Pathology” 2nd ed. (F.E. Vega, H.K. Kaya, eds.). Academic Press, San Diego. DOI: 10.1016/B978-0-12-384984-7.00006-3
- Vega F.E., Posada F., Aime M.C., Pava-Ripolli M., Infante F., Rehner S.A. 2008. Entomopathogenic fungal endophytes. *Biological Control* 46 (1): 72–82. DOI: <https://doi.org/10.1016/j.biocontrol.2008.01.008>
- Vidal S., Jaber L. 2015. Entomopathogenic fungi as endophytes: plant–endophyte–herbivore interactions and prospects for use in biological control. *Current Science* 109 (1): 46–54.
- Vincent C., Goettel M.S., Lazarovits G. (eds.) 2007. *Biological Control: A Global Perspective*. CAB International/A AFC, Wallingford, United Kingdom. DOI: 10.1079/9781845932657.0000
- Wagner B.L., Lewis L.C. 2000. Colonization of corn, *Zea mays*, by the entomopathogenic fungus *Beauveria bassiana*. *Applied and Environmental Microbiology* 66 (8): 3468–3473. DOI: 10.1128/aem.66.8.3468-3473.2000
- White T.J., Bruns T.D., Lee S.B., Taylor J.W. 1990. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. p. 315–322. In: “PCR Protocols: A Guide to Methods and Applications” (M.A. Innis, D.H. Gelfand, J.J. Sninsky, T.J. White, eds.). Academic Press, San Diego. DOI: <http://dx.doi.org/10.1016/B978-0-12-372180-8.50042-1>
- Wilson D. 1995. Endophyte – the evolution of a term, and clarification of its use and definition. *Oikos* 73: 274–276. DOI: <https://doi.org/10.2307/3545919>
- Wright S.P., Inglis G.D., Goettel M.S. 2007. Fungi. p. 223–248. In: “Field Manual of Techniques in Invertebrate Pathology: Application and Evaluation of Pathogens for Control of Insects and other Invertebrate Pest”. 2nd ed. (L.A. Lacey, H.K. Kaya, eds.). Springer, Dordrecht, the Netherlands.
- Wu Z.H., Wang T.H., Huang W., Qu Y.B. 2001. A simplified method for chromosome DNA preparation from filamentous fungi. *Mycosystema* 20 (4): 575–577.