

Fungal communities in barren forest soil after amendment with different wood substrates and their possible effects on trees', pathogens, insects and nematodes

Monika Matecka¹, Hanna Kwaśna^{2*}, Wojciech Szewczyk²

¹ Department of Forest Protection, Forest Research Institute, Sękocin Stary, Braci Leśnej 3, 05-090 Raszyn, Poland

² Department of Forest Pathology, Poznań University of Life Sciences, Wojska Polskiego 71c, 60-625 Poznań, Poland

Received: April 14, 2015

Accepted: June 26, 2015

Abstract: Scots pine sawdust, composted bark or coarse, post-harvest woody debris from conifers had been spread over the surface of barren forest soil before planting with Scots pine. The effects of the Scots pine sawdust, composted bark or coarse, post-harvest woody debris from conifers on the abundance and diversity of culturable fungi were investigated. The amendments were aimed at increasing the soil suppressiveness to *Armillaria* and *Heterobasidion*. The classical soil-dilution method was chosen for qualitative and quantitative analyses of fungal communities in soils because of its proven reliability and consistency. The soil was inhabited by saprotrophic fungi from Ascomycota and Zygomycota, including species known to be potential antagonists of *Armillaria* or *H. annosum* (i.e. *Clonostachys* + *Trichoderma* spp., *Penicillium commune*, *P. daleae*, *P. janczewskii*) or stimulants of *Armillaria* (i.e. *Pseudogymnoascus roseus*, *Trichocladium opacum*). Eleven years after treatment, the abundance and diversity of fungi, the abundance of *P. commune*, and locally the abundance of *P. janczewskii* increased, while *Clonostachys* + *Trichoderma* spp., and locally, *P. daleae* and *T. opacum* decreased. Amending the barren soil with organic matter does not guarantee effective, long-term suppressiveness of the sandy loam soil to *Armillaria* and *Heterobasidion*. Increased abundance of entomopathogenic and nematophagous species, 11 years after treatment, does suggest the long-term possibility of insect or nematode control in soil.

Key words: *Armillaria*, *Heterobasidion*, insects, nematodes, organic matter, soil fungi

Introduction

Poland has a 28.8% forest cover, which is less than the European average of 32%. At the same time, 25% of land (4.6 million ha) that is potentially available for arable use is unsuitable for agricultural production. Considering Poland's present stage of development, land-use structure, and environmental, geographic and climatic conditions, forest cover should be 33–34%. In 2002, the National Program for the Augmentation of Forest Cover (KPZL) was revised. The Program anticipates afforestation of 1.5 million ha of post-agricultural land and an increase in forest cover to 30% by 2020 and 33% by 2050 (Zajac and Kwiecień 2001). Land designated for afforestation includes 2.3 million ha of marginal land. Such lands include soils which are non-fertile, contaminated with chemicals, destroyed or mechanically transformed, and fallow land or land with unfavourable natural and topographic conditions.

First and second generation Scots pine (*Pinus sylvestris* L.) grown on soils formerly under agricultural use become seriously damaged by *Heterobasidion annosum* (Fr.) Bref. Young Scots pine plantations also become infested with *Armillaria* spp. (Fr. Fr.) Staude (Sierota 1995, 2013).

Armillaria ostoyae (Romagn.) Herink and *H. annosum* are among the most dangerous pathogens causing butt and root rot in conifers of temperate and boreal regions of the world, in Europe, North America, and Asia (Gregory *et al.* 1991; Korhonen and Stenlid 1998; Korhonen *et al.* 1998; Dettman and van der Kamp 2001; Kile *et al.* 2001; Dai *et al.* 2003, 2006; Żółciak 2007; Otrosina and Garbelotto 2010; Worrall *et al.* 2010). In Europe alone, *H. annosum* is responsible for the loss of 800 million euros annually. In Poland, in 2013, serious occurrence of butt and root rot caused by *Armillaria* and *H. annosum* were recorded on 99,215 and 146,379 ha, respectively (Tarwacki 2013).

Favourable conditions for pathogens include alkaline soils, soils with a disease history, and soils with a deficiency of competitive and antagonistic microorganisms. In these conditions, saplings of conifers can become infected very soon after being planted.

Damage by *H. annosum* is greater on former agricultural and fallow lands partly because of their higher nitrogen content, lower C : N ratio, deficiency in potassium and phosphorus, higher pH, lower capillary ascent of water and lower aggregate stability of soil under the "plough layer". Shortage of lignified tissues, which are the specific carbon and energy source for the forest soil microbiota,

*Corresponding address:

kwasna@up.poznan.pl

creates an unfavourable habitat for microorganisms that could contribute to forest site-forming processes (Trojanowski and Heider 1975).

The aim of the research was to determine the abundance, diversity, and temporal distribution of culturable fungi in barren forest soils amended with different forms of organic matter, i.e. Scots pine sawdust, composted bark and coarse, post-harvest coniferous woody debris. The hypothesis was that organic matter amendment would change the physical and chemical properties of soil, which in turn would initiate and stimulate microbiological shifts that would increase the long-term suppressiveness of soil to *Armillaria* and *Heterobasidion*. The classical soil-dilution method was chosen for qualitative and quantitative analyses of fungal communities in soils because of the method's proven reliability and consistency. The affiliation, activity and microbiological relationships of many soil fungi detected with such culture-based methods are already known. The microbiological status of soils so determined, can be used for comparing habitat qualities/properties. Alternative methods of profiling based on DNA sequences were not used because they tend to detect different communities. The communities most often detected are of non-culturable or slow-growing, non-sporulating organisms with unknown affiliation or activity and less obvious relevance in microbiological interactions (Kwaśna *et al.* 2008; van Elsas and Boersma 2011).

Materials and Methods

Site description and treatments

The experiment was established on barren, uncultivated soil in the Bielsk Forest District (north-east Poland, 23°11'11"E, 52°45'54"N). In October 2001, an area of 0.4 ha

was divided into 20 plots of 200 m² (10 × 20 m). Scots pine sawdust, composted bark or coarse, post-harvest coniferous woody debris was spread on the surface of the plots, along the proposed planting rows. The amount of amendment used was 7.5 dm³ per 1 m × 0.4 m area. One-year-old Scots pine seedlings were planted 0.6 m apart, in rows 1.2 m apart. The seedlings were planted in spring 2002 which was eight months after amendments were applied. In an additional treatment, 0.4 dm³ of composted bark was placed under the roots of each of the one-year-old Scots pine seedlings while they were being planted in spring 2002. In control plots, Scots pine seedlings were planted into non-amended soil. Treatment and control plots were replicated four times in a randomised block design.

Composted bark was prepared in windrows (4–5 m wide, 2.5–3 m high) and consisted exclusively of Scots pine bark. Composting occurred in three phases: 1) an initial phase of 1–2 days during which easily degradable compounds were decomposed; 2) a thermophilic phase, lasting several months, with temperatures 40–80°C, in which cellulose was primarily degraded; 3) a stabilisation phase, with a decreased rate of decomposition, lower temperature and colonisation by mesophilic microorganisms. Composted bark contained 5% readily degradable carbon (cellulose).

The soil in the plots was sandy-loam (10.8% clay, 13.6% silt, and 75.6% sand), with 0.353% organic carbon content, and a pH of 4.41 in water (Table 1). Increased amounts of carbon, nitrogen, phosphorus (3–4×), and potassium were present 11 years after treatment. The first year of the experiment (2002) was slightly warmer (mean annual temperature 8.25°C) and dryer (total annual rainfall 489.4 mm) than the last year (2012) when the mean temperature was 7.0°C and total annual rainfall was 611.0 mm.

Table 1. Physicochemical properties of the barren forest topsoil (3–20 cm layer) before organic substrate amendments (2001) and 11 years after amendments (2012)

Soil characteristics	Barren forest soil (2001)	11-year-old <i>Penicillium sylvestris</i> plantation (2012)				
		Scots pine sawdust-spread	composted bark		coarse coniferous woody debris-spread	the control
			under roots	spread		
pH in KCl ¹	3.970	3.945	4.030	3.940	3.955	4.000
pH in H ₂ O	4.410	4.710	4.720	4.695	4.510	4.585
Organic carbon (%) ²	0.353	0.470 ^{abcd}	0.357 ^{aefg}	0.553 ^{behi}	0.419 ^{efhj}	0.453 ^{dgi}
Total nitrogen (%) ³	0.029	0.049	0.040	0.056	0.046	0.048
C : N	12.10	9.40	9.15	9.80	9.15	9.15
Extractable phosphorus P ₂ O ₅ [mg · kg ⁻¹] ⁴	48.0	193.0	172.0	207.0	195.0	200.0
Extractable potassium [mg · kg ⁻¹] ⁵	7.00	11.70	10.95	12.10	10.00	11.25
Extractable magnesium [mg · kg ⁻¹] ⁵	1.00	1.10	1.05	1.35	1.00	1.20
Extractable calcium [mg · kg ⁻¹] ⁵	7.00	8.00 ^{abc}	4.30 ^{adef}	13.10 ^{bdgh}	5.10 ^{cegi}	8.25 ^{thi}
Extractable sodium [mg · kg ⁻¹] ⁵	4.00	1.65 ^{abc}	0.75 ^a	1.00 ^b	0.90 ^c	0.90 ^d

¹analysed with potentiometer according to norm PN-ISO 103390:1997

²analysed chemically according to norm PN-ISO 10694:2002

³analysed chemically according to norm PN-13878:2002

⁴analysed chemically with method of Egner-Riehm

⁵analysed chemically according to procedure PB-05 ed. 2

The same letter indicates a statistically significant difference according to one-way ANOVA at $p \leq 0.001$ or at $p \leq 0.05$

Collection of samples

In September 2002 (10 months after treatment) and 2012 (11 years after treatment), six 7.5-cm-diam. cores of non-rhizosphere soil were collected from the A horizon (0–20 cm deep) along a row of the Scots pine trees in each of the four randomised plots in each treatment. The sample from a single plot were bulked together and mixed by rotating for 10 h.

Isolation and identification of fungi

Soil fungi were isolated using the soil-dilution method (Warcup 1950, 1955). From the four bulked sub-samples of each plot, 1 g of soil was diluted in 149 g of sterile quartz sand and 0.02 g of the mixture was put into a Petri dish, then covered with liquid (50°C) Johnson-Martin's agar (JMA; glucose 10 g · l⁻¹, peptone 5 g · l⁻¹, KH₂PO₄ 1 g · l⁻¹, MgSO₄ · 7H₂O 0.5 g · l⁻¹, rose Bengal 0.03 g · l⁻¹, aureomycin 0.0025 g · l⁻¹, agar 20 g · l⁻¹). Eight replicates (Petri dishes) were made for each plot (= 32 Petri dishes for a treatment). All plates were incubated for 20 days at 25°C.

All colonies on each plate were examined macro- and microscopically and distinguished on the basis of colour, growth rate, hyphal characteristics, and sporulation. Colonies of each species were counted and representatives of fungi were identified by morphotyping on Potato Dextrose Agar (PDA; Difco PDA 39 g · l⁻¹, pH 5.5) and Synthetic Nutrient Agar (SNA; KH₂PO₄ 1 g · l⁻¹, KNO₃ 1 g · l⁻¹, MgSO₄ · 7H₂O 0.5 g · l⁻¹, KCl 0.5 g · l⁻¹, glucose 0.2 g · l⁻¹, sucrose 0.2 g · l⁻¹, agar 20 g · l⁻¹). The *Aspergillus* and *Penicillium* species were identified on Czapek Yeast Autolysate agar (CYA; sucrose 30 g · l⁻¹, powdered yeast extract 5 g · l⁻¹, KH₂PO₄ 1 g · l⁻¹, Czapek concentrate 10 ml · l⁻¹, agar 15 g · l⁻¹) and 2% Malt Extract Agar (MEA; powdered malt extract 20 g · l⁻¹, glucose 20 g · l⁻¹, peptone 1 g · l⁻¹, agar 20 g · l⁻¹). Identification was made using mycological keys (Raper and Thom 1949; Barron 1968; Domsch *et al.* 1980; Pitt 1979; Klich and Pitt 1992). Fungi abundance was defined as the number of colony forming units (*cfu*) in a treatment. Frequency of an individual species or group of species was defined as percentage of isolates in the total number of isolates. Diversity was defined as the number of species in a treatment. A species, or group of related species of fungi, was considered as: (i) eudominant, with a frequency of 10–100%; (ii) dominant, with a frequency of 5–10%; (iii) subdominant, with a frequency of 2–5%; (iv) recedent, with a frequency of 1–2%; (v) subrecedent, with a frequency of 0–1%, in at least one treatment (Tischler 1949).

Statistics

Differences in soil chemical properties and numbers of *cfu* from two different samples, were analysed with one-way analysis of variance (ANOVA) using Statgraphics™ Centurion (Statpoint Technologies, Inc. Warrenton, VA, USA). Diversity within and between microbial communities was compared using diversity indices calculated for each community (Magurran 1988). Species richness was indicated by the total number of species in the commu-

nity, Margalef's index (DMg), Shannon's diversity index (H'), and Simpson's diversity index (D). Evenness and dominance were indicated by Shannon's evenness index (E), and Berger-Parker's index (d). Similarity between fungal communities in two systems was determined by calculating the qualitative Sorensen's similarity index (CS). Relationships between the soil chemical properties and the abundances of fungi were estimated using Pearson's correlation coefficient.

Results

A total of 93 species of saprotrophic soil fungi from Ascomycota and Zygomycota were recorded in the barren, non-rhizosphere forest soil 10 months and 11 years after amending the soil with Scots pine sawdust, composted bark or coarse coniferous woody debris and after having been planted with Scots pines (Table 2). There were increases between 2002 and 2012 in total abundance of fungi, diversity of fungi, abundance of *Acremonium* + *Sagenomella* spp., *Aspergillus* spp., *Oidiodendron* spp. and *Penicillium* + *Talaromyces* spp., and often of the entomopathogenic and nematophagous species. There were decreases in abundance of *Mortierella* spp. and *Clonostachys* + *Trichoderma* spp. (Fig. 1, Table 3). There were 16 eudominant, 28 dominant, eight subdominant, five recedent, and 36 subrecedent species. Twenty-one species, including *A. kiliense* Grütz, *A. pteridii* W. Gams, *A. strictum* W. Gams, *A. ochraceus* Wilhelm, *C. albicans* Berkhout, *C. merdarium* (Link ex Grev), *C. candelabrum* (Bonord.) Schroers, *Mortierella* spp., *P. lilacinus* (Thom) Samson, *P. adametzii* Zaleski, *P. brevicompactum* Dierckx, *P. citreo-viride* Biourge, *P. cyclopium* Westling, *P. lilacinum* Thom, *T. opacum* (Corda) Hughes, *T. polysporum* (Link ex Pers.) Rifai, *T. mentagrophytes* (Robin) Blanchard, and *U. nana* (Linnem.) Arx already occurred 10 months after the treatment. Fifty-six species, including *A. fusidioides* (Nicot) W. Gams, *A. niger* Tiegh., *C. cladosporioides* (Fresen.) G.A. de Vries, *O. griseum* Robak, *P. commune* Thom, *P. steckii* Zaleski and *P. vinaceum* J.C. Gilman & E.V. Abbott (eudominants and dominants), and all subdominants, recedents and subrecedents, only began to occur 11 years after treatment. The most frequent *Penicillium* species were *P. commune*, *P. daleae* Zaleski, *P. janczewskii* Zaleski, *P. lilacinum* Thom, *P. steckii*, and *P. vinaceum*. From the *Trichoderma*, only *T. polysporum* was among the eudominants. Another three *Trichoderma* species, i.e. *T. hamatum* (Bonord.) Bain, *T. koningii* Oudem., and *T. viride* Pers. were among the dominants.

The relatively small number of fungal taxa and the infrequent occurrence of many taxa soon after treatment, resulted in relatively small diversity indices based on species richness (DMg) and proportional abundance of species (H') (Table 4). There was significantly more diversity after 11 years than 10 months after treatment. Communities were more evenly distributed after 11 years. Dominance, to a greater or lesser extent, of single taxa after 11 years resulted in smaller values for Shannon's evenness index (E) and often higher values for the dominance index (d). Sorensen's qualitative similarity index (CS), when used for comparing fungal communities between the two samples, showed there was the most similarity after the

Table 2. Frequency (%) of fungi in barren forest soil recently amended with organic substrates and planted with Scots pines (2002), and the frequency 11 years after amending (2012)


No.	Taxon	Organic matter amendment									
		Scots pine sawdust-spread		composted bark				coarse coniferous woody debris-spread		the control	
		2002	2012	2002	2012	2002	2012	2002	2012	2002	2012
Eudominants (frequency 10–100% in at least one treatment)											
1	<i>Acremonium fusidioides</i> (Nicot) W. Gams	–	15.1 ^{ab}	–	9.1 ^c	–	3.7 ^a	–	18.6 ^c	–	5.6 ^b
2	<i>Cladosporium</i> spp.	–	–	–	–	–	0.5 ^a	3.5	0.7 ^b	–	13.8 ^{ab}
3	<i>Clonostachys candelabrum</i> (Bonord.) Schroers	–	–	–	–	–	–	25.9	–	–	–
4	<i>Oidiendron tenuissimum</i> (Peck) S. Hughes	7.1 ^a	32.8 ^{abe}	–	11.4 ^{bcdf}	–	21.6 ^{eg}	–	26.6 ^{dh}	–	5.6 ^{efgh}
5	<i>Penicillium citrinum</i> Thom	–	–	–	–	–	–	–	–	10.4	–
6	<i>P. commune</i> Thom	–	10.6 ^{ab}	–	22.8 ^{ad}	–	31.6 ^{bce}	–	18.2 ^{ce}	–	8.2 ^{de}
7	<i>P. cyclopium</i> Westling	–	–	–	–	5.7 ^a	–	–	–	23.6 ^a	–
8	<i>P. daleae</i> Zaleski	–	–	7.7	–	5.7	–	29.2	1.0	–	–
9	<i>P. janczewskii</i> Zaleski	7.1 ^a	6.6	10.1 ^b	0.5 ^{bef}	11.5 ^c	2.9 ^c	17.2 ^{ad}	8.8 ^{df}	–	5.6 ^e
10	<i>P. lilacinum</i> Thom	–	–	28.3	–	17.1	–	–	–	–	–
11	<i>P. purpurescens</i> Sopp	25.1 ^a	–	–	–	–	–	–	–	10.4 ^a	–
12	<i>P. steckii</i> Zaleski	–	1.4 ^a	–	10.3 ^{abc}	–	1.1 ^b	–	2.6 ^c	–	5.6
13	<i>P. vinaceum</i> J.C. Gilman & E.V. Abbott	–	12.6 ^a	–	9.7 ^b	–	16.9 ^{bcd}	–	4.3 ^{ace}	–	8.2 ^{de}
14	<i>Pseudogymnoascus roseus</i> Raïllo	7.1	–	–	–	–	0.8 ^a	–	–	13.2 ^a	19.4
15	<i>Trichocladium opacum</i> (Corda) Hughes	14.4 ^a	–	–	–	–	–	–	–	5.3 ^a	–
16	<i>Trichoderma polysporum</i> (Link ex Pers.) Rifai	17.9 ^{ab}	–	–	–	14.3 ^{ac}	–	6.9 ^{bc}	–	–	–
Dominants (frequency 5–10% in at least one treatment)											
17	<i>A. kiliense</i> Grütz	–	–	10.0	–	–	–	–	–	5.3	–
18	<i>A. pteridii</i> W. Gams	7.1	–	–	–	–	–	–	–	–	–
19	<i>A. roseogriseum</i> (S.B. Saksena) W. Gams	–	0.4	–	–	–	0.3	–	–	5.3	–
20	<i>A. strictum</i> W. Gams	–	–	–	–	8.6	–	–	–	–	–
21	<i>Alternaria alternata</i> (Fr.) Keissl.	7.1	–	–	–	–	–	–	–	–	2.8
22	<i>Aspergillus niger</i> Tiegh.	–	1.0	–	5.2	–	2.9	–	0.7	–	–
23	<i>A. ochraceus</i> Wilhelm	–	–	5.2	–	–	–	–	–	–	–
24	<i>Candida albicans</i> Berkhout	–	–	–	–	5.7	–	–	–	–	–
25	<i>Chrysosporium merdarium</i> (Link ex Grev)	–	–	–	–	5.7	–	–	–	–	–
26	<i>C. cladosporioides</i> (Fresen.) G.A. de Vries	–	0.7	–	–	–	–	–	0.7	–	5.6
27	<i>Lecanicillium lecanii</i> (Zimm.) Zare & W. Gams	–	0.4	7.7	2.9	5.7	6.4	–	3.2	–	–
28	<i>Leptographium lundbergii</i>	–	–	–	–	–	–	–	–	5.3	–
29	<i>Mortierella macrocystis</i> W. Gams	–	–	5.2	–	–	–	–	–	–	–
30	<i>M. spinosa</i> Linnem	–	–	7.7	–	–	–	–	–	–	–
31	<i>O. griseum</i> Robak	–	1.8	–	9.1	–	1.9	–	5.8	–	–
32	<i>Paecilomyces carneus</i> (Duché & R. Heim) A.H.S. Br. & G. Sm.	–	1.4	–	0.5	–	–	–	0.3	5.3	–
33	<i>P. lilacinus</i> (Thom) Samson	–	–	–	–	5.7	–	–	–	–	–
34	<i>P. adametzii</i> Zalewski	–	–	5.2	–	–	–	–	–	–	–
35	<i>P. brevicompactum</i> Dierckx	–	–	–	–	8.6	–	–	–	–	–
36	<i>P. canescens</i> Sopp	–	–	–	–	–	–	–	–	5.3	–
37	<i>P. citreo-viride</i> Biourge	–	–	–	–	–	–	5.2	–	–	–
38	<i>Simplicillium lamellicola</i> (F.E.V. Sm.) Zare & W. Gams	–	–	–	–	–	–	–	–	5.3	–
39	<i>Trichoderma hamatum</i> (Bonord.) Bain	–	–	–	–	–	–	–	–	5.3	–
40	<i>T. koningii</i> Oudem.	–	–	–	–	5.7	–	3.5	0.7	–	2.8
41	<i>T. viride</i> Pers.	–	–	7.7	–	–	–	–	0.3	–	–
42	<i>Trichophyton mentagrophytes</i> (Robin) Blanchard	7.1	–	–	–	–	–	–	–	–	–
43	<i>Ulocladium consortiale</i> (Thüm.) Simmons	–	–	5.2	–	–	–	–	–	–	–
44	<i>Umbelopsis nana</i> (Linnem.) Arx	–	–	–	–	–	–	8.6	–	–	–
Subdominants (frequency 2–5% in at least one treatment)											
45	<i>Acrodontium salmonium</i> de Hoog	–	–	–	–	–	–	–	–	–	2.8
46	<i>Cladosporium herbarum</i> (Pers.) Link	–	0.4	–	–	–	0.5	–	0.7	–	2.8
47	cf. <i>Cylindrocladiella hahajimaensis</i> (Ts. Watanabe) Inderb., R.M. Bostock & K.V. Subbarao	–	–	–	–	–	0.5	–	–	–	–
48	<i>Metacordyceps chlamydosporia</i> (H.C. Evans) G.H. Sung, J.M. Sung, Hywel-Jones & Spatafora	–	–	–	–	–	–	–	–	–	2.8
49	<i>Metarhizium anisopliae</i> (Metschn.) Sorokin	–	2.1	–	0.5	–	0.5	–	–	–	–
50	<i>Paraconiothyrium fuckelii</i> (Sacc.) Verkley & Gruyter	–	2.1	–	1.2	–	–	–	0.3	–	–
51	<i>Sagenomella diversispora</i> (J.F.H. Beyma) W. Gams	–	–	–	3.4	–	0.5	–	0.7	–	–
52	<i>Talaromyces variabilis</i> (Sopp) Samson, Yilmaz, Frisvad & Seifert	–	–	–	2.8	–	–	–	0.3	–	–

Table 2. Frequency (%) of fungi in barren forest soil recently amended with organic substrates and planted with Scots pines (2002), and the frequency 11 years after amending (2012) – Continuation

No.	Taxon	Organic matter amendment									
		Scots pine sawdust-spread		composted bark				coarse coniferous woody debris-spread		the control	
		2002	2012	2002	2012	2002	2012	2002	2012	2002	2012
Recedents (frequency 1–2% in at least one treatment)											
53	<i>Haptocillium sphaerosporum</i> (Goodey) Zare & W. Gams	–	–	–	1.7	–	–	–	0.3	–	–
54	<i>Penicillium ochrochloron</i> Biourge	–	1.7	–	–	–	–	–	–	–	–
55	<i>P. spinulosum</i> Thom	–	–	–	–	–	–	–	2.0	–	–
56	<i>Talaromyces pinophilus</i> (Hedgc.) Samson, Yilmaz, Frisvad & Seifert	–	0.4	–	–	–	–	1.3	–	–	2.8
57	<i>T. deliquescens</i> (Sopp) Jaklitsch	–	0.4	–	1.1	–	–	–	–	–	–
Subrecedents (frequency 0–1% in at least one treatment)											
Total abundance – number of colony forming units (cfu) in a sample		84 ^{afghi}	28 ^{amn^o}	117 ^{bfj}	176 ^{bmprs}	105 ^{gk}	379 ^{cnptu}	174 ^{dhjkl}	307 ^{dortw}	114 ^{eil}	36 ^{esuw}
Diversity – number of species		9 ^a	36 ^{ae}	11 ^b	27 ^b	12 ^c	31 ^c	8 ^d	29 ^d	12	16 ^e

Subrecedents (with frequency < 1%) included *Apiospora montagnei* Sacc., *A. fumigatus* Fresen., *A. versicolor* (Vuill.) Tirab. *C. pannicola* (Corda) Oorschot & Stalpers, *Chrysosporium* sp., *Fusarium oxysporum* Schldt., *Geomyces pannorum* (Link) Sigler & J.W. Carmich., *Humicola fuscoatra* Traaen, *H. grisea* Traaen, *Nectria inventa* Pethybr., *O. echinulatum* G.L. Barron, *P. variotii* Bainier, *P. dierckxii* Biourge, *P. expansum* Link, *P. megasporum* Orpurt & Fennell, *P. raistrickii* G. Sm., *Phialophora bubakii* (Laxa) Schol-Schwarz, *S. striatispora* (Onions & G.L. Barron) W. Gams, *T. flavus* (Klöcker) Stolk & Samson, *T. islandicus* (Sopp) Samson, Yilmaz, Frisvad & Seifert, *T. rugulosus* (Thom) Samson, Yilmaz, Frisvad & Seifert, *Trichothecium roseum* (Pers.) Link, and *U. isabellina* (Oudem.) W. Gams, non-sporulating – 13 species

The same letter indicates a statistically significant difference according to one-way ANOVA at $p \leq 0.001$ or at $p \leq 0.05$



- 1, 2 – Scots pine sawdust-spread (2002, 2012)
- 3, 4 – composted bark under roots (2002, 2012)
- 5, 6 – composted bark-spread (2002, 2012)
- 7, 8 – coarse post-harvest coniferous woody debris-spread (2002, 2012)
- 9, 10 – the control (2002, 2012)

Fig. 1. Frequencies of the most common taxa in forest barren soil (2002) and in 11-year-old Scots pine plantation (2012) after organic substrate amendment

application of coarse woody debris (Table 5). Comparable similarity among all treatments, 11 years after their application was also shown, however, with the CS index.

The fungal communities contained a few species known to be antagonists of *Armillaria* or *H. annosum*, i.e. *C. candelabrum*, *G. pannorum* (Link) Sigler & J.W. Carmich., *P. adametzii*, *P. commune*, *P. daleae*, *P. janczewskii*, and *Trichoderma* spp., or stimulants of *Armillaria*, i.e. *M. macrocystis* W. Gams, *P. adametzii*, *P. citrinum* Thom, *P. spinulo-*

sum Thom, *P. roseus* Raullo, and *T. opacum* (Table 6). Eleven years after treatment there was an increase in all treatments of only one antagonist of *H. annosum*, i.e. *P. commune*, whilst the abundance of *P. janczewskii* increased slightly after the sawdust treatment. Overall, there were small decreases in populations of four *Armillaria* stimulants, i.e. *M. macrocystis*, *P. adametzii*, *P. roseus*, and *T. opacum*. But the effect was significant only for *T. opacum* after the sawdust treatment.

Table 3. Abundances (*cfu* per sample) of the most common taxa in barren forest soil recently amended with organic substrates and planted with Scots pines (2002), and 11 years after amending (2012)

No.	Taxon	Organic matter amendment									
		Scots pine sawdust-spread		composted bark				coarse coniferous woody debris-spread		the control	
		2002	2012	under roots		spread		2002	2012	2002	2012
1	<i>Acronium + Sagenomella</i> spp.	6 ^a	47 ^{aefgh}	12 ^b	22 ^{beij}	0 ^c	17 ^{cfk}	0 ^c	62 ^{dgikl}	12 ^d	2 ^{dhl}
2	<i>Aspergillus</i> spp.	0	4 ^b	6	9 ^c	0 ^a	14 ^{abd}	0	2 ^c	0	0 ^d
3	<i>Clonostachys + Trichoderma</i> spp.	15 ^{aef}	1 ^a	9 ^{bgh}	2 ^b	21 ^{cgij}	0 ^c	63 ^{dehi}	3 ^d	6 ^{fi}	1
4	<i>Mortierella</i> spp.	0	0	5	0	0	0	5	0	0	0
5	<i>Oidiodendron</i> spp.	6 ^a	98 ^{aef}	0 ^b	36 ^{beghi}	0 ^c	89 ^{egi}	0 ^d	101 ^{dh}	0	2 ^{fij}
6	<i>Penicillium + Talaromyces</i> spp.	21 ^{aefghi}	97 ^{ajkl}	54 ^{bf}	83 ^{bmno}	42 ^{cg}	20 ^{cjmp}	90 ^{dh}	116 ^{dknp}	57 ^{ei}	11 ^{elo}
	Entomopathogenic and nematophagous <i>Haptocillium + Lecanicillium + Metacordyceps + M. anisopliae + Paecilomyces + Simplicillium</i> ¹	0 ^{aefg}	10 ^{akl}	9 ^{eh}	5 ^{mn}	12 ^{bfi}	26 ^{bkmop}	0 ^{chij}	13 ^{co}	12 ^{dgi}	1 ^{dnp}

Explanations – see table 2

¹according to Domsch *et al.* 1980; Watanabe 1980; Rombach *et al.* 1986; Gams *et al.* 2004; Inglis *et al.* 2006; Sun and Liu 2006; Quesada-Moraga *et al.* 2014

Table 4. Diversity indices for fungi in barren forest soil recently amended with organic substrates and planted with Scots pines (2002), and 11 years after amending (2012)

Index	Organic matter amendment									
	Scots pine sawdust-spread		composted bark				coarse coniferous woody debris-spread		the control	
	2002	2012	under roots		spread		2002	2012	2002	2012
Species richness indices										
Margalef's index (DMg)	2.40	6.19	2.72	5.02	3.09	5.05	1.72	4.88	3.02	4.18
Shannon's diversity index (H')	2.06	2.32	2.22	2.45	2.39	2.15	1.79	2.27	2.32	2.57
Simpson's diversity index (D)	0.14	0.16	0.13	0.11	0.10	0.18	0.19	0.15	0.11	0.09
Evenness or dominance indices										
Shannon's evenness index (E)	0.94	0.65	0.93	0.74	0.96	0.48	0.86	0.67	0.93	0.93
Berger-Parker's index (d)	0.25	0.33	0.28	0.23	0.17	0.31	0.29	0.27	0.24	0.19
Sorensen qualitative similarity index (CS)	0.08		0.10		0.09		0.21		0.07	

Table 5. Sorensen qualitative similarity index (CS) between treatments for fungal communities in barren forest soil recently amended with organic substrates and planted with Scots pines (2002), and 11 years after amending (2012)

Treatments compared		Sorensen qualitative similarity index (CS)	
1	2	2002	2012
Scots pine sawdust – spread	composted bark under roots	0.04	0.44
Scots pine sawdust – spread	composted bark – spread	0.19	0.44
Scots pine sawdust – spread	coarse coniferous woody debris – spread	0.23	0.40
Composted bark under roots	composted bark – spread	0.34	0.41
Composted bark under roots	coarse coniferous woody debris – spread	0.23	0.50
Composted bark –spread	coarse coniferous woody debris – spread	0.40	0.40

Table 6. Abundances (cfu per sample) of antagonists and stimulants of *Heterobasidium annosum* and of *Armillaria* in barren forest soil recently amended with organic substrates and planted with Scots pines (2002), and 11 years after amending (2012)

Taxon	Organic matter amendment									
	Scots pine sawdust-spread		composted bark				coarse coniferous woody debris-spread		the control	
			under roots		spread					
2002	2012	2002	2012	2002	2012	2002	2012	2002	2012	
<i>Armillaria</i> antagonists										
<i>Clonostachys candelabrum</i> ¹	0	0	0	0	0	0	45 ^{ab}	0 ^a	0 ^b	0
<i>Penicillium janczewskii</i> ¹	6 ^{acd}	19 ^{ajkl}	12 ^{bef}	1 ^{bjmn}	12 ^{gh}	11 ^{mop}	30 ^{cegi}	27 ^{knor}	0 ^{dghi}	2 ^{lpr}
Total	6 ^{ad}	19 ^{ajj}	12 ^{be}	1 ^{bik}	12 ^{fg}	11 ^l	75 ^{cefh}	27 ^{cklm}	0 ^{gh}	2 ^{jm}
<i>Armillaria</i> stimulants										
<i>Mortierella macrocystis</i> ¹	0	0	6	0	0	0	0	0	0	0
<i>P. adametzii</i> ¹	0	0	6	0	0	0	0	0	0	0
<i>P. citrinum</i> ¹	0 ^a	0	0	0	0	0	0	0	12 ^a	0
<i>P. spinulosum</i> ¹	0	0	0	0	0	0	0	6	0	0
<i>Pseudogymnoascus roseus</i> ¹	6 ^a	0 ^b	0	0	0	3	0	0	15 ^a	7 ^b
<i>Trichocladium opacum</i> ¹	12 ^a	0 ^a	0	0	0	0	0	0	6	0
Total	18 ^{ade}	0 ^{af}	12 ^{bd}	0 ^b	0 ^e	3	0 ^f	6	33 ^{ee}	7 ^{cf}
<i>Heterobasidium annosum</i> antagonists										
<i>Geomyces pannorum</i> ¹	0	1	0	0	0	0	0	0	0	0
<i>P. adametzii</i> ¹	0	0	6	0	0	0	0	0	0	0
<i>P. commune</i> ²	0 ^a	30 ^{aefg}	0 ^b	40 ^{bhij}	0 ^c	120 ^{cehkl}	0 ^d	56 ^{dikm}	0	3 ^{gilm}
<i>P. daleae</i> ¹	0 ^{cd}	0	9 ^{ace}	0 ^a	6 ^f	0	51 ^{bdefg}	3 ^b	0 ^g	0
<i>P. janczewskii</i> ²	6 ^{ac}	19 ^{ajj}	12 ^{bde}	1 ^{biklm}	12 ^{fg}	11 ^{knop}	30 ^{cdft}	27 ^{lmnp}	0 ^{egh}	2 ^{op}
<i>Trichoderma</i> spp. ^{1,2,3}	15 ^{ab}	1 ^a	9 ^{cd}	2	21 ^{bcef}	0 ^b	63 ^{bddeg}	3 ^c	6 ^{fg}	1
Total	21 ^{adefg}	50 ^{anopr}	36 ^{d hij}	43 ^{stu}	39 ^{bekln}	131 ^{bhosvw}	144 ^{efkm}	89 ^{ciptvz}	6 ^{glm}	6 ^{iruwx}

Explanations – see table 2

¹according to Kwaśna 1995, 1997a, b, c, 2001, 2002; Kwaśna *et al.* 2001; Szwajkowska-Michałek *et al.* 2012

²according to Nona *et al.* 1968; Werner and Zadworny 2002

³according to Nikolajeva *et al.* 2012

Discussion

The application of organic substrates to soil usually results in short- or long-term shifts in microbial populations. If such population changes are directed and stimulated, the outcome may be an enhanced biological control of soil-borne pathogens.

Natural, organic substrates applied in forestry usually include sawdust, composted bark or chipped post-harvest woody debris. These have been previously applied with good biological control effects. In forest soils, coniferous sawdust effectively increased the density of *Trichoderma harzianum* Rifai, a potential antagonist of *Armillaria* spp. and *H. annosum* (Sierota 1995, 2013; Sierota and Kwaśna 1999; Kwaśna *et al.* 2000; Nikolajeva *et al.* 2012). Incorporation of sawdust into the soil or sawdust with *Trichoderma* into the soil, increased the survival of Scots pine seedlings affected by damping-off fungi; the effect was comparable to protection with chemical fungicides (Duda and Sierota 1987). Composted tree bark suppresses many soil-borne pathogens, eliminating the need to sterilise the media, and reducing fungicide applications. Composted tree bark has been most widely adopted for commercial, containerised production of nursery stock, particularly

of ornamental plants and trees (Hoitink *et al.* 1975, 1976). Chipped post-harvest coniferous woody debris left on the soil surface or mixed with the topsoil had a long-term suppressive effect towards *Armillaria* and *Heterobasidium* on a Scots pine planted acreage (Kwaśna *et al.* 2014).

The abundance and diversity of fungi increased 11 years after organic matter amendment was used. These results are consistent with Werner *et al.* (2001) who observed that fungi were more abundant in arable soils richer in nutrients. However, the extent of the response of the resident fungi to cellulose inputs is currently unknown (Eichorst and Kusk 2012). The induction and production of cellulase by fungi, particularly in natural conditions, are not completely understood (Dashtban *et al.* 2011; Nayebyazdi *et al.* 2012). The abundance and diversity of cellulose-responsive fungi is generally unique to a certain soil type, suggesting a strong potential influence of multiple edaphic factors.

In the present study, total abundances of fungi were only weakly correlated (mainly because of low abundance in the control in 2012) with organic carbon and total nitrogen contents ($r = 0.57$ and 0.56 , $p < 0.0001$), and negatively, with extractable potassium and sodium

($r = -0.49$ and -0.28 , $p < 0.0001$). The increased abundances of the common taxa were mostly not correlated with soil nutrient contents (i.e. *Acremonium* + *Sagenomella*, *Oidiodendron*, *Penicillium* + *Talaromyces*) or moderately or strongly negatively correlated with organic carbon, total nitrogen, extractable phosphorus, potassium, magnesium, and calcium contents (i.e. *Clonostachys* + *Trichoderma* spp., $r = -0.53 - 0.97$, $p < 0.0001$). The response of *Trichoderma* can be attributed to the negative effect of carbon and nitrogen on sporulation of the fungus (Rajput *et al.* 2014). Considering the above-mentioned relationships, the resulting positive correlation of the total abundances of fungi to soil nutrient content seemed to be due to the positive correlation of a large number of infrequent species recorded.

The seasonal abundances and diversity of fungi seemed to result from the degradation rate of the incorporated substrate and the succession due to specific degrading abilities. Many recorded fungi were soft-rot fungi, with cellulolytic activity and cellulase production (Stursová *et al.* 2012). Others, however, i.e. *Acremonium* + *Sagenomella*, *Oidiodendron* and *Penicillium* + *Talaromyces* spp. mostly utilise starch and pectin, which are released after earlier decomposition of cellulose (Bååth and Söderström 1980; Deacon *et al.* 2006), and they appear in the later phase of degradation. They appear after an absence of cellulose causes the elimination of *Clonostachys* + *Trichoderma* spp. which are common decomposers of cellulose and strong antagonists of other fungi; Deacon *et al.* 2006). The very frequent appearance of *Oidiodendron* in the soil, 11 years after the treatments, could have resulted, not only from its preferences for woody substrates (Nilsson 1973; Domsch *et al.* 1980; Dalpé 1991) but also from its ability to form ericoid mycorrhizas with *Calluna* and *Vaccinium* species (Couture *et al.* 1983), which were present on the site.

Classification of fungi into eudominants, dominants, subdominants, recedents, and subrecedents helps to show the scale of the functional potential of individual species. The relatively small numbers of frequent species and larger numbers of infrequent species in fungal communities agree with Rubino and McCarthy (2003), and Richard *et al.* (2004). The less frequent species were particularly common 11 years after treatments. They are particularly relevant for decomposition processes and ecosystem functioning. The less frequent species are often more active and efficient decomposers than those occurring more frequently (Deacon *et al.* 2006).

The most frequent species of *Penicillium* (i.e., *P. commune*, *P. daleae*, *P. janczewskii*, *P. lilacinum*, *P. steckii*, and *P. vinaceum*) are common in Polish forest soils and may not occur elsewhere in the world (Kwaśna and Nirenberg 1994; Grantina-Ievina *et al.* 2013).

The behaviour of the fungal communities (abundance of single species and diversity) was similar among all four treatments. These findings may indicate that the degradation of sawdust, composted bark, and even coarse coniferous woody debris proceeds at the same rate and that similar physico-chemical and biological effects can be expected after their application.

The fungal communities contained a few species known to be antagonists of *Armillaria* or *H. annosum*, i.e.

C. candelabrum, *G. pannorum*, *P. adametzii*, *P. commune*, *P. daleae*, *P. janczewskii*, and *Trichoderma* spp., or stimulants of *Armillaria*, i.e. *M. macrocystis*, *P. adametzii*, *P. citrinum*, *P. spinulosum*, *P. roseus*, and *T. opacum* (Table 6). Mycelium and/or metabolites of the antagonists are known to: (i) inhibit growth of *Armillaria* and *Heterobasidion* in paired cultures *in vitro*; (ii) inhibit growth and ramification of *Armillaria* rhizomorphs *in vivo*; (iii) decrease the extent of root rot caused by *Armillaria* and decrease the extent of necrosis caused by *H. annosum* on *P. sylvestris* seedlings and other plants; (iv) stimulate growth of Scots pine seedlings inoculated with *Armillaria*; (v) effectively colonise the thin (< 5 mm diam) and thick (> 5 mm diam.) living roots and rhizosphere of younger and older coniferous and broad-leaved trees and their stumps, which helped in competition with pathogens (Kwaśna 1995, 1997a, b, c, 2001, 2002; Kwaśna *et al.* 2001; Werner and Zadworny 2002; Nikolaeva *et al.* 2012; Szwajkowska-Michałek *et al.* 2012).

Abundance of species that are entomopathogenic [*M. anisopliae* (Metschn.) Sorokin, *P. carneus* (Duché & R. Heim) A.H.S. Br. & G. Sm.] or nematophagous [*H. sphaerosporum* (Goodey) Zare & W. Gams] often increased after treatment and was weakly positively correlated with organic carbon and extractable calcium contents ($r = 0.70$ and 0.68 , $p < 0.0001$), and negatively correlated with extractable potassium content ($r = -0.82$, $p < 0.0001$) (Table 3). These fungi may help control insects or nematodes in forest soil (Watanabe 1980; Inglis and Tigano 2006).

Since *Armillaria* and *Heterobasidion* continuously infect trees over a long period, useful antagonists are required to have long persistence in the ecosystem without the addition of nutrients or other amendments. Only one *H. annosum* antagonist: *P. commune*, was found to have an increased population 11 years after all soil amendment treatments. This fungus was previously found to occur at a low frequency in barren and abandoned farmland soils 1–2 years after amendment with Scots pine sawdust (Sierota and Kwaśna 1988, 1998). In the present study, *P. commune* became one of the eudominants 11 years after treatment, which theoretically suggests a long-lived functional potential. It should be mentioned, though, that only one of two tested isolates of *P. commune* was found to be antagonistic to *H. annosum in vitro* (Werner and Zadworny 2002), indicating functional diversity in its activity. Functional diversity of fungi had been observed earlier (van der Heijden *et al.* 1998; Szwajkowska-Michałek *et al.* 2012). Intra-specific variation in function in different conditions, can be high. Some organisms, e.g. *Scytalidium lignicola* Pesante, have remarkable antagonistic activity towards *Heterobasidion* only *in vitro*, while others, e.g. *Resinicium bicolor* (Alb. & Schwein.) Parmasto, have activity only on/in wood (Highley 1990, 1994). So far, similar antagonistic activity towards *H. annosum in vitro* and *in vivo* has been observed in *P. adametzii* (Szwajkowska-Michałek *et al.* 2012).

Conclusion

Spreading Scots pine sawdust, composted bark or coarse, post-harvest coniferous woody debris on the soil surface of barren forest land increased soil nutrients and

the abundance and diversity of culturable fungi in the long term (after 11 years). The increased abundance of only two known antagonists of *Armillaria* and *Heterobasidion* (particularly where sawdust or composted bark was spread on the soil surface) does not guarantee the effective long-term suppressiveness of soil. The increased abundance of entomopathogenic and nematophagous species does suggest the possibility of long-term control of insects and nematodes.

Acknowledgements

The authors wish to thank the National Forest Holding "State Forests" in Poland for funding part of the study within the research project No BLP-214 (2001–2002) and the Ministry of Science and Higher Education of the Republic of Poland for funding another part of the study within the research project No IBL 204304 (2011–2012).

References

- Barron G.L. 1968. The Genera of Hyphomycetes from the Soil. The Williams & Wilkins Co., Baltimore, USA, 364 pp.
- Bååth E., Söderström B.E. 1980. Degradation of macromolecules by microfungi isolated from different podzolic soil horizons. *Canadian Journal of Botany* 58 (4): 422–425.
- Couture M., Fortin J.A., Dalpe Y. 1983. *Oidiendron griseum* Robak: an endophyte of ericoid mycorrhiza in *Vaccinium* spp. *New Phytologist* 95 (3): 375–380.
- Dai Y.-C., Vainio E.J., Hantula J., Niemelä T., Korhonen K. 2003. Investigations on *Heterobasidion annosum* s. lat. in central and eastern Asia with the aid of mating tests and DNA fingerprinting. *Forest Pathology* 33 (5): 269–286.
- Dai Y.-C., Yuan H.S., Wei Y.L., Korhonen K. 2006. New records of *Heterobasidion parviporum* from China. *Forest Pathology* 36 (4): 287–293.
- Dalpe Y. 1991. Statut endomycorrhizien du genre *Oidiendron*. *Canadian Journal of Botany* 69 (8): 1712–1714.
- Dashtban M., Buchkowski R., Qin W. 2011. Effect of different carbon sources on cellulase production by *Hypocrea jecorina* (*Trichoderma reesei*) strains. *International Journal of Biochemistry and Molecular Biology* 2 (3): 274–286.
- Deacon L.J., Pryce-Miller E.J., Frankland J.C., Bainbridge B.W., Moore P.D., Robinson C.H. 2006. Diversity and function of decomposer fungi from a grassland soil. *Soil Biology and Biochemistry* 38 (1): 7–20.
- Dettman J.R., van der Kamp B.J. 2001. The population structure of *Armillaria ostoyae* and *Armillaria sinapina* in the central interior of British Columbia. *Canadian Journal of Botany* 79 (5): 600–611.
- Domsch K.H., Gams W., Anderson T.H. 1980. Compendium of Soil Fungi. Academic Press, London, UK, 672 pp.
- Duda B., Sierota Z.H. 1987. Survival of Scots pine seedlings after biological and chemical control of damping-off in plastic greenhouses. *Forest Pathology* 17 (2): 110–117.
- Eichorst S.A., Kusk C.R. 2012. Identification of cellulose-responsive bacterial and fungal communities in geographically and edaphically different soils by using stable isotope probing. *Applied and Environmental Microbiology* 78 (8): 2316–2327.
- Gams W., Diederich P., Poldmaa K. 2004. Fungicolous fungi. p. 344–391. In: "Biodiversity of Fungi: Inventory and Monitoring Methods" (G.M. Mueller, G.F. Bills, M.S. Foster, eds.). Elsevier Academic Press, Burlington, MA, USA, 777 pp.
- Grantina-Ievina L., Kasparinskis R., Tabors G., Nikolajeva V. 2013. Features of saprophytic soil microorganism communities in conifer stands with or without *Heterobasidion annosum sensu lato* infection: a special emphasis on *Penicillium* spp. *Environmental and Experimental Biology* 11: 23–38.
- Gregory S.C., Rishbeth J., Shaw C.G. III. 1991. Pathogenicity and virulence. p. 76–87. In: "Armillaria Root Disease" (C.G. III Shaw, G.A. Kile, eds.). Agriculture Handbook No. 691, United States Department of Agriculture, Forest Service, Washington, DC, USA, 233 pp.
- Highley T.L. 1990. Laboratory studies on antagonism of *Scytalidium lignicola* to wood decay fungi. *Material und Organismen* 25 (3): 181–192.
- Highley T.L. 1994. Effect of *Scytalidium lignicola* on decay resistance and strength of wood. International Research Group on Wood Preservation. Document NoIRG/WP/94–10061.
- Hoitink H.A.J., Schmitthenner A.F., Herr L.J. 1975. Composted bark for control of root rot in ornamentals. *Ohio Report on Research and Development* 60: 25–26.
- Hoitink H.A.J., Herr L.J., Schmitthenner A.F. 1976. Survival of some plant pathogens during composting of hardwood tree bark. *Phytopathology* 66 (11): 1369–1372.
- Inglis P.W., Tigano M.S. 2006. Identification and taxonomy of some entomopathogenic *Paecilomyces* spp. (Ascomycota) isolates using rDNA-ITS sequences. *Genetics and Molecular Biology* 29 (1): 132–136.
- Kile G.A., McDonald G.I., Byler J.W. 2001. Ecology and disease in natural forests. p. 102–121. In: "Armillaria Root Disease" (C.G. III Shaw, G.A. Kile, eds.). Agriculture Handbook No. 691, United States Department of Agriculture, Forest Service, Washington, DC, USA, 233 pp.
- Klich M.A., Pitt J.I. 1992. A Laboratory Guide to the Common *Aspergillus* Species and their Teleomorphs. Commonwealth Scientific and Industrial Research Organisation, Division of Food Processing, North Ryde, New South Wales, Australia, 116 pp.
- Korhonen K., Capretti P., Karjalainen R., Stenlid J. 1998. Distribution of *Heterobasidion annosum* intersterility groups in Europe. p. 93–104. In: "Heterobasidion annosum: Biology, Ecology, Impact and Control" (S. Woodward, J. Stenlid, R. Karjalainen, A. Hüttermann, eds.). CABI, Wallingford, UK, 589 pp.
- Korhonen K., Stenlid J. 1998. Biology of *Heterobasidion annosum*. p. 43–70. In: "Heterobasidion annosum: Biology, Ecology, Impact and Control" (S. Woodward, J. Stenlid, R. Karjalainen, A. Hüttermann, eds.). CABI, Wallingford, UK, 589 pp.
- Kwaśna H. 1995. Fungal communities in soil beneath Scots pine and their stumps. Effect of fungi on *Heterobasidion annosum* and *Armillaria ostoyae* growth. *Acta Mycologica* 30 (2): 193–205.
- Kwaśna H. 1997a. Antagonistic effect of fungi communities from Scots pine fine roots on *Heterobasidion annosum* (Fr.) and *Armillaria ostoyae* (Romagn.) Herink growth. *Phytopathologia Polonica* 13: 133–146.
- Kwaśna H. 1997b. Antagonistic effect of fungi from Scots pine stump roots on *Heterobasidion annosum* and *Armillaria ostoyae*. *Acta Mycologica* 32 (2): 369–381.

- Kwaśna H. 1997c. Fungi on the surface of roots of Scots pine and its stumps and effect on *Heterobasidion annosum* (Fr.) Bref. and *Armillaria ostoyae* (Romagn.) Herink growth. Polish Agricultural Annual, ser. E, 26 (1/2): 109–123.
- Kwaśna H. 2001. Fungi in the rhizosphere of common oak and its stumps and their possible effect on infection by *Armillaria*. Applied Soil Ecology 17 (3): 215–227.
- Kwaśna H. 2002. Changes in microfungal communities in roots of *Quercus robur* stumps and their possible effect on colonization by *Armillaria*. Journal of Phytopathology 150 (7): 403–411.
- Kwaśna H., Bateman G.L., Ward E. 2008. Determining species diversity of microfungal communities in forest tree roots by pure-culture isolation and DNA sequencing. Applied Soil Ecology 40 (1): 44–56.
- Kwaśna H., Kotyńska U., Łakomy P., Mallett K. 2001. Stimulation of *Armillaria* rhizomorph formation by oak root fungi. Acta Mycologica 36 (2): 257–272.
- Kwaśna H., Sierota Z., Bateman G.L. 2000. Fungal communities in fallow soil before and after amending with pine sawdust. Applied Soil Ecology 14 (2): 177–182.
- Kwaśna H., Nirenberg H. 1994. The effectiveness of two methods used for isolating soil fungi. Acta Mycologica 29 (1): 13–22.
- Kwaśna H., Walkowiak U., Łakomy P., Gornowicz R., Mikiciński A., Gałązka S., Szewczyk W. 2014. An effect of different silviculture techniques on the functional diversity of microorganisms in soil of *Pinus sylvestris* plantation. Journal of Plant Protection Research 55 (3): 241–253.
- Magurran A.E. 1988. Ecological Diversity and its Measurement. Princeton University Press, Princeton, NJ, USA, 179 pp.
- Nayebyazdi N., Salary M., Ghanbary M.A.T., Ghorbany M., Bahmanyar M.A. 2012. Investigation of cellulase activity in some soil borne fungi isolated from agricultural soils. Annals of Biological Research 3: 5705–5713.
- Nikolajeva V., Petrina Z., Vulfa L., Alksne L., Eze D., Grantina L., Gaitnieks T., Lielpetere A. 2012. Growth and antagonism of *Trichoderma* spp. and conifer pathogen *Heterobasidion annosum* s.l. *in vitro* at different temperatures. Advances in Microbiology 2 (3): 295–302.
- Nilsson T. 1973. Studies on wood degradation and cellulolytic activity of microfungi. Studia Forestalia Suecica 104: 1–40.
- Nona D.A., Blake M.I., Crespi H.L., Katz J.J. 1968. Effect of deuterium oxide on the culturing of *Penicillium janczewskii* II. Isolation of fully deuterated griseofulvin. Journal of Pharmaceutical Sciences 57 (6): 975–979.
- Otrosina W.J., Garbelotto M. 2010. *Heterobasidion occidentale* sp. nov. and *Heterobasidion irregulare* nom. nov.: a disposition of North American *Heterobasidion* biological species. Fungal Biology 114 (1): 16–25.
- Pitt J. 1979. The Genus *Penicillium* and its Teliomorphic States *Eupenicillium* and *Talaromyces*. Academic Press, New York, USA, 634 pp.
- Quesada-Moraga E., Herrero N., Zabalgogezcoa I. 2014. Entomopathogenic and nematophagous fungal endophytes. Advances in Endophytic Research 1: 85–99.
- Rajput A.Q., Khanzada M.A., Shahzad S. 2014. Effect of different organic substrates and carbon and nitrogen sources on growth and shelf life of *Trichoderma harzianum*. Journal of Agricultural Science and Technology 16 (4): 731–745.
- Raper K.B., Thom C. 1949. A Manual of the Penicillia. Williams & Wilkins Co. Baltimore, USA, 875 pp.
- Richard F., Moreau P.A., Selosse M.A., Gardes M. 2004. Diversity and fruiting patterns of ectomycorrhizal and saprobic fungi in an old-growth Mediterranean forest dominated by *Quercus ilex* L. Canadian Journal of Botany 82 (12): 1711–29.
- Rombach M.C., Shepard B.M., Roberts D.W. 1986. Biological control: Insect pathogens. In: "Management of Rice Insects" (E.A. Henrichs, ed.). Wiley, New York, USA, 791 pp.
- Rubino D.L., McCarthy B.C. 2003. Composition and ecology of macrofungal and myxomycete communities on oak woody debris in a mixed-oak forest of Ohio. Canadian Journal of Forest Research 33 (11): 2151–2163.
- Sierota Z. 1995. Rola grzyba *Phlebiopsis gigantea* (Fr.:Fr.) Jülich w ograniczaniu huby korzeni w drzewostanach sosny zwyczajnej (*P. sylvestris* L.) na gruntach porolnych. [The Role of the Fungus *Phlebiopsis gigantea* (Fr.:Fr.) Jülich as a Limiting Factor of the *Heterobasidion annosum* (Fr.) Bref. in the Scots Pine (*Pinus sylvestris* L.) Stands in Post-agricultural Lands]. Prace Instytutu Badawczego Leśnictwa, Warszawa, 810 pp.
- Sierota Z. 2013. *Heterobasidion* root rot in forests on former agricultural lands in Poland: Scale of threat and prevention. Academic Journal 8 (47): 2298–2305.
- Sierota Z., Kwaśna H. 1988. Effect of pine sawdust on structure of soil fungi communities in the soils of post agricultural land. Acta Mycologica 33 (1): 77–90.
- Sierota Z., Kwaśna H. 1998. Changes in fungal communities in abandoned farmland soil enriched with pine sawdust. Folia Forestalia Polonica. Series A. Forestry 40: 85–94.
- Sierota Z., Kwaśna H. 1999. Ocena mikologiczna zmian zachodzących w glebie gruntu porolnego po dodaniu trocin iglastych. [A mycological assessment of changes occurring in the post-agricultural land soil after adding softwood sawdust]. Sylwan 4: 57–66.
- Stursová M., Zifčáková L., Leigh M.B., Burgess R., Baldrian P. 2012. Cellulose utilization in forest litter and soil: identification of bacterial and fungal decomposers. FEMS Microbiology Ecology 80 (3): 735–746.
- Sun M., Liu X. 2006. Carbon requirements of some nematophagous, entomopathogenic and mycoparasitic hyphomycetes as fungal biocontrol agents. Mycopathologia 161 (5): 295–305.
- Szwajkowska-Michałek L., Kwaśna H., Łakomy P., Perkowski J. 2012. Inhibition of *Armillaria* and *Heterobasidion* growth by *Penicillium adametzii* isolated from *Pinus sylvestris* forest soil. Forest Pathology 42 (6): 454–466.
- Tarwacki G. 2013. Krótkoterminowa prognoza występowania ważniejszych szkodników i chorób infekcyjnych drzew leśnych w Polsce w 2013 roku. [Short-term prognosis of occurrence of the important pests and diseases of trees in Poland in 2013]. Instytut Badawczy Leśnictwa: 1–173.
- Tischler W. 1949. Grundzüge der terrestrischen Tierökologie. Friedrich Vieweg und Sohn, Braunschweig, Germany, 219 pp.
- Trojanowski J., Heider K. 1975. Degradation of phenolic compounds by soft rot and white rot fungi. p. 417–418. In: "Biodegradation and Humification" (G. Kilbertus, O. Reisinger, A. Mourey, J.A. Consela da Fonseca, eds.). Pierron, Sarreguemines, France, 496 pp.
- van der Heijden M.G.A., Klironomos J.N., Ursic M., Moutoglou P., Streitwolf-Engel R., Boller T., Wiemken A., Sanders I.R. 1998. Mycorrhizal fungal diversity determines plant biodiversity, ecosystem variability and productivity. Nature 396: 69–72.

- van Elsas J.D., Boersma F.G.H. 2011. Review of molecular methods to study the microbiota of soil and the mycosphere. *European Journal of Soil Biology* 47 (2): 77–87.
- Warcup J.H. 1950. The soil-plate method for isolation of fungi from soil. *Nature* 166: 117–118.
- Warcup J.H. 1955. Isolation of fungi from hyphae present in soil. *Nature* 175: 953–954.
- Watanabe T. 1980. A new variety of *Verticillium sphaerosporum*, an endoparasite of nematodes and its antagonism to soil-borne plant pathogens. *Japanese Journal of Phytopathology* 46 (5): 598–605.
- Werner A., Werner M., Kwaśna H. 2001. Grzyby wybranych środowisk gleb użytkowanych rolniczo i ugorów. [Fungi in some agricultural soils or barren soils]. p. 287–295. In: „Drobnoustroje środowiska glebowego – aspekty fizjologiczne, biochemiczne, genetyczne”. [“Microorganisms in soil habitat – physiological, biochemical and genetical aspects”] (H. Dahm, A. Pokojska-Burdziej, eds.). Adam Marszałek, Toruń, Poland, 331 pp.
- Werner A., Zadworny M. 2002. Interaction between microfungi from arable and fallow land soils and *Heterobasidion annosum in vitro*. *Dendrobiology* 4: 51–58.
- Worrall J.J., Harrington T.C., Blodgett J.T., Conklin D.A., Fairweather M.L. 2010. *Heterobasidion annosum* and *H. parviporum* in the southern Rocky Mountains and adjoining states. *Plant Disease* 94 (1): 115–118.
- Zajac S., Kwiecień R. 2001. Niektóre aspekty modyfikacji krajowego programu zwiększania lesistości. [Some aspects of modification of the National Program for the Augmentation of Forest Cover]. *Sylvan* 10: 23–34.
- Żółciak A. 2007. *Armillaria* species in coniferous stands. *Acta Mycologica* 42 (2): 211–217.