

ORIGINAL ARTICLE

## Involvement of iturin and surfactin in inhibition of a post-harvest fungal pathogen on green bell pepper

Yuliar Yuliar<sup>1\*</sup> , Shun Tomita<sup>2</sup>, Hinako Hishinuma<sup>2</sup>, Kenji Yokota<sup>2</sup>

<sup>1</sup>Research Center for Applied Microbiology, National Research and Innovation Agency, Jl. Raya Jakarta -Bogor, KM 46, KST Soekarno, Cibinong, Bogor 16911, Indonesia

<sup>2</sup>Department of Agriculture Chemistry, Faculty of Agriculture, Tokyo University of Agriculture, 1-1-1 Sakuragaoka, Setagaya-ku, Tokyo 156-8502, Japan

---

**Vol. 65, No. 3: 411-412, 2025**

**DOI: 10.24425/jppr.2025.155786**

Received: September 30, 2024

Accepted: February 02, 2025

Online publication: September 23, 2025

\*Corresponding address:

[yrivaie@yahoo.com](mailto:yrivaie@yahoo.com); [yuli003@brin.go.id](mailto:yuli003@brin.go.id)

Responsible Editor:

Assunta Bertaccini

---

### SUPPLEMENTARY MATERIAL

The authors are fully responsible for both the content and the formal aspects of the supplementary material. No editorial adjustments were made.



**Fig. S1.** Suppressive ability of *Bacillus* sp. TS001 strain to reduce rot lesion diameter of green bell pepper caused by *Colletotrichum scovillei* 244830; (A) "no inoculum, (B) *C. scovillei* infection, (C) treatment group (green bell pepper with pathogen and TS001 strain inoculations)