

SHORT COMMUNICATION

Does the invasion intensity of *Erigeron annuus* (L.) Pers. and community invasibility increase as the number of plant species increases?

Yingsheng Liu¹, Yizhuo Du¹, Xiaoxuan Geng¹, Congyan Wang^{1*}, Daolin Du²

¹ Institute of Environment and Ecology, School of the Environment and Safety Engineering, Jiangsu University, Zhenjiang, China

² Jingjiang College, Jiangsu University, Zhenjiang, China

DOI: 10.24425/jppr.2026.3254

Received: October 10, 2025

Accepted: January 23, 2026

Online publication: September 07, 2026

*Corresponding address:

liuyuxue623@ujs.edu.cn

Responsible Editor:


Irena Suwara

SUPPLEMENTARY MATERIAL

The authors are fully responsible for both the content and the formal aspects of the supplementary material. No editorial adjustments were made.

Table S1. The determination methods and the corresponding references for the analyzed variables in this study

Measured indices	Determination methods	References
Relative Abundance	$P_i = \frac{n_i}{N}$ where n_i represents the number of individuals of plant species i and N represents the total number of individuals of all plant species in one particular quadrat, respectively	Wang <i>et al.</i> (2021); Wang <i>et al.</i> (2018); Wang <i>et al.</i> (2019)
Invasion intensity index (III)	$III = \frac{P_i}{MaxP_i}$ where P_i represents the relative abundance of invasive plant species i in one particular invaded quadrat and $MaxP_i$ is the maximum value of relative abundance of invasive plant species i among all invaded quadrats, respectively. In particular, invasive plant species in invaded quadrats with higher values of III possess greater degree of invasion intensity compared to those with lower values of III.	[Wang <i>et al.</i> (2020)]
Shannon's diversity index (H')	$H' = -\sum_{i=1}^S P_i \ln P_i$ where S represents the number of plant species and P_i represents the relative abundance of plant species i in one particular quadrat, respectively. P_i was calculated as follows: $P_i = n_i / N$, where n_i represents the number of individuals of plant species i and N represents the total number of individuals of all plant species in one particular quadrat, respectively.	[Shannon and Weaver (1949)]
Simpson's dominance index (D)	$D = 1 - \sum_{i=1}^S (n_i/N)^2$ where S represents the number of plant species in one particular quadrat. n_i represents the number of individuals of plant species i and N is the total number of individuals of all plant species in one particular quadrat, respectively.	[Simpson (1949)]
Pielou's evenness index (E_H)	$E_H = \frac{H'}{\ln S}$ where S represents the number of plant species and H' is Shannon's diversity index in one particular quadrat, respectively.	[Pielou (1966)]
Margalef's richness index (F)	$F = \frac{S - 1}{\ln N}$ where S represents the number of plant species and N is the total number of individuals of all plant species in one particular quadrat, respectively.	[Margalef (1951)]
Community invasibility index (CII)	$CII = 1 - (MaxP_i - P_i)$ where P_i represents the observed relative abundance of invasive plant species in one particular invaded quadrat and $MaxP_i$ is the maximum relative abundance of invasive plant species among all invaded quadrats, respectively. The relative abundance of invasive plant species was calculated as the ratio of the number of invasive plant species individuals to the total number of individuals of all plant species in one particular invaded quadrat. In particular, invaded quadrats with higher values of CII possess greater degree of invasibility compared to those with lower values of CII.	[Wang <i>et al.</i> (2020)]

**Fig. S1.** The monodominant communities of *Erigeron annuus* (L.) Pers

References

- Margalef R. 1951. Diversidad de especies en las comunidades naturales. Barcelona: Publicaciones del Instituto de Biología Aplicada 6: 59–72
- Pielou E.C. 1966. The measurement of diversity in different types of biological collections. *Journal of Theoretical Biology* 13: 131–144. DOI: 10.1016/0022-5193(66)90013-0
- Shannon C.E., Weaver W. 1949. *The Mathematical Theory of Communication*. University of Illinois Press, Urbana, Illinois, 117 pp. DOI: 10.1063/1.3067010
- Simpson E.H. 1949. Measurement of diversity. *Nature* 163: 688. DOI: 10.1136/thx.27.2.261
- Wang C.Y., Cheng H.Y., Wang S., Wei M., Du D.L. 2021. Plant community and the influence of plant taxonomic diversity on community stability and invasibility: A case study based on *Solidago canadensis* L. *Science of the Total Environment* 768: 144518. DOI: 10.1016/j.scitotenv.2020.144518
- Wang C.Y., Jiang K., Liu J., Zhou J.W., Wu B.D. 2018. Moderate and heavy *Solidago canadensis* L. invasion are associated with decreased taxonomic diversity but increased functional diversity of plant communities in East China. *Ecological Engineering* 112: 55–64. DOI: 10.1016/j.ecoeng.2017.12.025
- Wang C.Y., Wei M., Wang S., Wu B.D., Cheng H.Y. 2020. *Erigeron annuus* (L.) Pers. and *Solidago canadensis* L. antagonistically affect community stability and community invasibility under the co-invasion condition. *Science of the Total Environment* 716: 137128. DOI: 10.1016/j.scitotenv.2020.137128
- Wang S., Wei M., Wu B.D., Jiang K., Du D.L., Wang C.Y. 2019. Degree of invasion of Canada goldenrod (*Solidago canadensis* L.) plays an important role in the variation of plant taxonomic diversity and community stability in eastern China. *Ecological Research* 34: 782–789. DOI: 10.1111/1440-1703.12049