

## REVIEW

## Availability constraints of plant protection products in Polish winter cereal production – a post-EU accession perspective

Przemysław Strażyński<sup>1\*</sup>, Roman Kierzek<sup>2</sup>, Marek Mrówczyński<sup>1</sup>

<sup>1</sup> Department of Entomology and Agricultural Pests, Institute of Plant Protection – National Research Institute, Poznań, Poland

<sup>2</sup> Department of Weed Science and Plant Protection Techniques, Institute of Plant Protection – National Research Institute, Poznań, Poland

DOI: 10.24425/jppr.2025.156889

Received: June 04, 2025

Accepted: August 04, 2025

Online publication: December 04, 2025

\*Corresponding address:

p.strazynski@iorpib.poznan.pl

Responsible Editor:

Arkadiusz Artyszak

### Abstract

Due to their extensive cultivation and the significant impact of pests causing economically relevant losses, winter cereal crops require effective chemical protection. A major challenge in their protection is the steadily diminishing range of available plant protection products (PPPs), specifically the active substances (AS) they contain. Although the core principles of the European Green Deal have been temporarily suspended, the European Commission has continued to phase out several active substances. As a result, dozens of these substances have been withdrawn from the market in recent years, creating increasing difficulties in crop protection - particularly for winter cereals. This is partly due to the growing resistance of pests, stemming from reduced opportunities for rotating products with different mechanisms of action. Further reductions in active substances, in the absence of viable alternative methods for agrophage control, may potentially lead to a decrease in both the cultivated area and overall production volumes for economic reasons. Moreover, such a scenario increases the risk of illegal imports or off-label use of plant protection products (PPP).

**Keywords:** agronomics constraints, cereal sustainability, crop protection policy, EU pesticide regulation, integrated pest management,

## Introduction

In Poland, winter cereals have dominated the sown area for many years, currently covering approximately 4.5 million hectares (SP 2024, 2025). Agroclimatic conditions, including recurring spring droughts (IMWM-NRI 2025), provide more favorable conditions for the cultivation of winter cereal varieties, which typically have higher yields than their spring counterparts (IAFP-NRI 2024). In the autumn of 2024, winter cereal sowing was led by winter wheat, which occupied 2.2 million hectares, followed by winter triticale (1.2 million ha), winter rye (0.7 million ha), and winter barley (0.4 million ha) (SP 2024, 2025). The extensive cultivation of winter wheat and the intensity of its production contribute to significant pest threats, with as many as 100 economically important species

identified. A similar level of pest pressure – approximately 90 species – is observed in winter barley. In Poland, slightly lower pest incidence is reported for winter triticale (around 80 species), with the lowest pressure recorded in winter rye plantations, which are affected by about 70 pests (Strażyński *et al.* 2024).

In the European Union, since January 1, 2014, all professional users have been required to implement integrated plant protection systems, which has led to a reduction in the use of chemical agents in agricultural production, including winter cereals (OJEU 2009; IAFP-NRI 2024). The application of the general principles of integrated plant protection by professional users of plant protection products is regulated in Poland by the Act of March 8, 2013 on Plant Protection

Products (Journal of Laws 2013, item 455), as well as by the Regulation of the Minister of Agriculture and Rural Development of April 18, 2013 on the Requirements of Integrated Plant Protection (Journal of Laws 2013, item 505). Within integrated protection, non-chemical methods – agrotechnical, biological, and breeding-based practices – are prioritized for reducing pest pressure. Nevertheless, crops such as winter cereals still require chemical protection. Although the number of microbiological plant protection products is gradually increasing, their availability remains insufficient (MARD 2025). However, under the Integrated Plant Production eco-scheme, their use in cereals (as well as in other crops for which Integrated Production methodologies have been developed) is a mandatory requirement.

The withdrawal of active substances (ASs) from plant protection products is a legally regulated process designed to safeguard human and animal health, protect the environment, and ensure food quality (OJEU 2009; Alix and Lewis 2010; MacLeod *et al.* 2010; Cilia and Kandris 2023). If an active substance is deemed unsafe or fails to meet regulatory standards, it may be withdrawn from the market (Mie and Rudén 2022; Slunge *et al.* 2023; Solé *et al.* 2024). This process typically involves several stages, including scientific risk assessment and formal administrative decisions (MARD 2025):

**Risk and Safety Assessment:** Regulatory bodies such as the European Food Safety Authority (EFSA) and national authorities evaluate the risks associated with the active substance. The assessment considers potential health impacts on humans and animals, environmental consequences, and the risk of contamination in groundwater or agricultural products.

**Administrative Decisions:** If the active substance poses an unacceptable risk, regulatory authorities may decide to withdraw it from use. In the European Union, this process is coordinated by the European Commission, which can prohibit the use of the substance across all Member States. Such decisions are formalized through regulations or directives.

**Loss of Marketing Authorization:** A withdrawn active substance loses its marketing authorization, meaning that it can no longer be manufactured, sold, or applied. This withdrawal may affect the substance in all uses or be restricted to specific applications considered particularly hazardous.

The diversity of cropping systems across Europe, driven by significant geographic and climatic variation, presents substantial challenges for harmonized crop protection strategies. The economic viability of European agriculture is increasingly constrained by stringent regulatory frameworks, particularly the prohibition of numerous pesticides previously authorized within the EU but still in use globally. This regulatory

landscape has the potential to place EU agricultural production at a competitive disadvantage, prompting European farmers to seek scientific support to develop and implement Integrated Pest Management (IPM) strategies. Achieving stable yields and high crop quality while simultaneously reducing dependence on chemical plant protection products remains a key challenge for the sector. The substantial biophysical and socio-economic heterogeneity across European regions further complicates efforts to meaningfully reduce pesticide use (Lamichhane *et al.* 2016; Jørgensen *et al.* 2019).

Over the past decade, the use of major conventional pesticides in European agriculture has shown considerable variability, with no consistent downward trend despite multiple policy initiatives. This fluctuation is partly attributable to interannual climatic variability, which affects pest incidence and, consequently, pesticide application levels. Moreover, the reduced availability of certain active substances may have led to increased use of alternative compounds, often at higher application rates. For instance, in Denmark, pesticide use (measured in  $\text{kg} \cdot \text{ha}^{-1}$ ) declined between 1981 and 2000 but has subsequently increased. These trends have been influenced by factors such as changes in pesticide taxation and anticipated regulatory restrictions on specific active ingredients. Similar patterns have been observed in France, where pesticide use has risen in recent years. Data from the German Reference Farm Network also reveal significant year-to-year and crop-specific variability in pesticide application intensity (Freier *et al.* 2013).

The aim of this paper was to assess the current constraints on the availability of plant protection products in Polish winter cereal production, with particular attention to regulatory, market, and agronomic factors shaped by Poland's EU accession. The scope of the paper includes a comparative analysis of product registrations, and national-level consequences of EU policy changes.

## Protective treatments in winter cereals

Polish agriculture, in comparison to many other European Union countries, is characterized by moderate and stable consumption of plant protection products per 1 ha of crops (Zieliński *et al.* 2025a). In Poland, winter wheat receives the highest number of protective treatments among winter cereal crops, with an average of 8.76 applications of plant protection products (PPP) per growing season. This is followed by winter barley (5.89 treatments), winter triticale (4.75 treatments), and winter rye, which has the lowest number of applications at 3.59 treatments per season. The greatest

number of fungicide treatments is also observed in winter wheat, averaging 4.22 applications. This is followed by winter barley (3.00), winter triticale (1.90), and winter rye (1.33), which receives nearly three fewer fungicide treatments than winter wheat. Herbicide applications are most frequent in winter wheat (3.10 treatments), followed by winter triticale (2.53), winter barley (2.21), and winter rye (1.84), indicating lower sensitivity to weed pressure. Insecticide treatments are generally limited across all winter cereals. Winter wheat receives an average of 0.47 applications, winter barley 0.27, winter triticale 0.10, and winter rye the fewest at 0.06 treatments. The application of plant growth regulators is highest in winter wheat (0.98 treatments), followed by winter barley (0.53), winter rye (0.37), and winter triticale (0.21). For comparison, in the United Kingdom, winter wheat plantations typically receive a total of nine protective treatments, a number comparable to that in Poland. These include three fungicide applications, three herbicide treatments, two applications of growth regulators, and one treatment for pest control (IAFP–NRI 2024; Zieliński *et al.* 2025a).

## Consumption of active substances in winter cereals

Currently, the highest consumption of active substances among winter cereals occurs in the cultivation of winter wheat (*Triticum aestivum*), with an average

application rate of 1.24 kg · ha<sup>-1</sup> (Table 1). Slightly lower amounts are applied in winter triticale (*x Triticosecale*) (1.16 kg · ha<sup>-1</sup>) and winter barley (*T. vulgare*) (1.14 kg · ha<sup>-1</sup>). The lowest consumption is observed in winter rye (*Secale cereale*), at 0.31 kg · ha<sup>-1</sup>, which is 0.93 kg · ha<sup>-1</sup> less than in winter wheat. Fungicides are used most extensively in winter barley (0.45 kg · ha<sup>-1</sup>), followed by winter wheat (0.44 kg · ha<sup>-1</sup>), winter triticale (0.29 kg · ha<sup>-1</sup>), and winter rye (0.08 kg · ha<sup>-1</sup>), representing a 0.36 kg · ha<sup>-1</sup> difference between winter wheat and rye. In terms of herbicide use, the highest consumption is recorded in winter triticale (0.74 kg · ha<sup>-1</sup>), followed by winter barley (0.60 kg · ha<sup>-1</sup>) and winter rye (0.17 kg · ha<sup>-1</sup>), indicating a 0.57 kg · ha<sup>-1</sup> difference between triticale and rye. The highest insecticide use is found in winter wheat, at 0.024 kg · ha<sup>-1</sup>, while no insecticides are currently registered for use in winter rye. A similar pattern is seen in the application of molluscicides, with the highest usage in winter wheat and the lowest in winter barley (IAFP–NRI 2024).

## Number of registered plant protection products in winter cereals

The sheer number of available products does not equate to effectiveness. A large number of products is registered for winter cereals but the number of new and effective products with new modes of action have diminished over the past years. Presently, the Ministry of Agriculture and Rural Development in Poland,

**Table 1.** Consumption of active substances (kg · ha<sup>-1</sup>) in winter cereals in selected years

| Years            | Fungicides | Herbicides | Insecticides | Molluscicides | Total |
|------------------|------------|------------|--------------|---------------|-------|
| WINTER WHEAT     |            |            |              |               |       |
| 2020             | 0.444      | 0.444      | 0.024        | 0.256         | 1.239 |
| 2017             | 0.55       | 0.41       | 0.005        | 0.31          | 1.32  |
| 2011             | 0.50       | 0.62       | 0.03         | 0.35          | 1.49  |
| 2003             | 0.64       | 0.73       | 0.02         | 0.46          | 1.85  |
| WINTER TRITICALE |            |            |              |               |       |
| 2022             | 0.29       | 0.74       | 0.01         | 0.12          | 1.16  |
| 2016             | 0.24       | 0.44       | 0.01         | 0.08          | 0.77  |
| 2010             | 0.16       | 0.51       | 0.01         | 0.06          | 0.74  |
| WINTER BARLEY    |            |            |              |               |       |
| 2024             | 0.45       | 0.60       | 0.01         | 0.01          | 1.13  |
| 2005             | 0.42       | 0.81       | 0.01         | 0.13          | 1.37  |
| WINTER RYE       |            |            |              |               |       |
| 2024             | 0.08       | 0.17       | 0.00         | 0.06          | 0.31  |
| 2012             | 0.04       | 0.23       | 0.00         | 0.04          | 0.30  |
| 2003             | 0.13       | 0.33       | 0.00         | 0.06          | 0.53  |

Source: Own elaboration based on data from the Ministry of Agriculture and Rural Development

responsible for registering plant protection products, has approved the highest number of such products for winter wheat – 1,192 in total (Table 2). This is followed by winter triticale (793), winter barley (711), and winter rye, which has the lowest number at 662 registered products – 530 fewer than winter wheat (MARD 2025). The European Union has released a list of active substances that are slated for substitution or potential removal. The withdrawal of these substances is expected

to have the greatest impact on winter wheat, with up to 19 active substances at risk of being phased out. Similar reductions may occur in winter rye (18 substances), winter triticale (16), and to a slightly lesser extent in winter barley (15). The potential withdrawal of active substances could lead to significant reductions in available fungicidal seed dressings – by as much as 93% in winter barley, 86% in winter triticale, and 85% in both winter wheat and winter rye.

**Table 2.** Number of registered plant protection products for winter cereals in Poland (June 2025)

| Type of PPP           | Total Number of Preparations | Number of Preparations Remaining after Reduction | Reduction Level [%] |
|-----------------------|------------------------------|--|---------------------|
| WINTER WHEAT          |                              |  |                     |
| Herbicides            | 471                          | 352  | 25                  |
| Fungicides            | 422                          | 311  | 26                  |
| Insecticides          | 86                           | 61   | 29                  |
| Molluscicides         | 31                           | 31   | 0                   |
| Growth regulators     | 98                           | 97   | 0                   |
| Insecticide dressings | 0                            | 0  | 0                   |
| Fungicide dressings   | 84                           | 13   | 85                  |
| TOTAL                 | 1192                         | 865  | 27                  |
| WINTER TRITICALE      |                              |  |                     |
| Herbicides            | 334                          | 241  | 28                  |
| Fungicides            | 267                          | 207  | 23                  |
| Insecticides          | 24                           | 11   | 54                  |
| Molluscicides         | 28                           | 28   | 0                   |
| Growth regulators     | 71                           | 71   | 0                   |
| Insecticide dressings | 0                            | 0  | 0                   |
| Fungicide dressings   | 69                           | 10   | 86                  |
| TOTAL                 | 793                          | 568  | 28                  |
| WINTER BARLEY         |                              |  |                     |
| Herbicides            | 253                          | 182  | 28                  |
| Fungicides            | 272                          | 221  | 19                  |
| Insecticides          | 17                           | 6  | 65                  |
| Molluscicides         | 29                           | 29   | 0                   |
| Growth regulators     | 71                           | 71   | 0                   |
| Insecticide dressings | 0                            | 0  | 0                   |
| Fungicide dressings   | 69                           | 5  | 93                  |
| TOTAL                 | 711                          | 514  | 28                  |
| WINTER RYE            |                              |  |                     |
| Herbicides            | 280                          | 192  | 31                  |
| Fungicides            | 210                          | 151  | 28                  |
| Insecticides          | 19                           | 8  | 58                  |
| Molluscicides         | 29                           | 29   | 0                   |
| Growth regulators     | 58                           | 58   | 0                   |
| Insecticide dressings | 0                            | 0  | 0                   |
| Fungicide dressings   | 66                           | 10   | 85                  |
| TOTAL                 | 662                          | 448  | 32                  |

Source: Own elaboration based on data from the Ministry of Agriculture and Rural Development

## Weed control in winter cereal plantations

On a global scale, potential crop losses for the six major crops (wheat, rice, maize, potatoes, soybeans, and

cotton) amount to 69%. Weeds account for the largest share of these losses at 34%, followed by pests at 19% and diseases at 16% (Oerke 2006). Weeds pose a significant economic threat to winter cereals, particularly winter wheat, where they contribute most substantially to yield reduction. The highest number of herbicides

**Table 3.** Number of herbicides registered in Poland for weed control in winter cereals (June 2025)

| Weeds                             | Current Registration | After Possible Withdrawal |
|-----------------------------------|----------------------|---------------------------|
| <b>WINTER WHEAT</b>               |                      |                           |
| Geranium pusillum                 | 90                   | 56                        |
| Centaurea cyanus                  | 197                  | 121                       |
| Viola arvensis                    | 273                  | 167                       |
| Stellaria media                   | 375                  | 268                       |
| Papaver rhoeas                    | 275                  | 163                       |
| Matricaria perforata              | 282                  | 189                       |
| Apera spica-venti                 | 220                  | 161                       |
| Veronica persica                  | 207                  | 114                       |
| Galium aparine                    | 209                  | 193                       |
| <i>Brassica napus</i> (self-sown) | 233                  | 176                       |
| Capsella bursa-pastoris           | 338                  | 265                       |
| <b>WINTER TRITICALE</b>           |                      |                           |
| Geranium pusillum                 | 86                   | 47                        |
| Centaurea cyanus                  | 162                  | 121                       |
| Viola arvensis                    | 202                  | 160                       |
| Stellaria media                   | 271                  | 182                       |
| Papaver rhoeas                    | 234                  | 171                       |
| Matricaria perforata              | 230                  | 158                       |
| Apera spica-venti                 | 165                  | 111                       |
| Veronica persica                  | 162                  | 119                       |
| Galium aparine                    | 230                  | 182                       |
| <i>Brassica napus</i> (self-sown) | 185                  | 143                       |
| Capsella bursa-pastoris           | 250                  | 174                       |
| <b>WINTER BARLEY</b>              |                      |                           |
| Geranium pusillum                 | 51                   | 27                        |
| Centaurea cyanus                  | 109                  | 58                        |
| Viola arvensis                    | 134                  | 75                        |
| Stellaria media                   | 200                  | 141                       |
| Papaver rhoeas                    | 157                  | 108                       |
| Matricaria perforata              | 143                  | 101                       |
| Apera spica-venti                 | 107                  | 70                        |
| Veronica persica                  | 110                  | 71                        |
| Galium aparine                    | 158                  | 102                       |
| <i>Brassica napus</i> (self-sown) | 128                  | 83                        |
| Capsella bursa-pastoris           | 178                  | 127                       |

**Table 3.** Number of herbicides registered in Poland for weed control in winter cereals (June 2025) – continuation

| Weeds                             | Current Registration | After Possible Withdrawal |
|-----------------------------------|----------------------|---------------------------|
| WINTER RYE                        |                      |                           |
| <i>Geranium pusillum</i>          | 75                   | 38                        |
| <i>Centaurea cyanus</i>           | 138                  | 101                       |
| <i>Viola arvensis</i>             | 164                  | 122                       |
| <i>Stellaria media</i>            | 226                  | 133                       |
| <i>Papaver rhoeas</i>             | 198                  | 144                       |
| <i>Matricaria perforata</i>       | 191                  | 127                       |
| <i>Apera spica-venti</i>          | 143                  | 98                        |
| <i>Veronica persica</i>           | 150                  | 106                       |
| <i>Galium aparine</i>             | 197                  | 150                       |
| <i>Brassica napus</i> (self-sown) | 162                  | 124                       |
| <i>Capsella bursa-pastoris</i>    | 217                  | 132                       |

Source: Own elaboration based on data from the Ministry of Agriculture and Rural Development

registered for weed control is found in winter wheat. Among these, the most common are herbicides targeting *Stellaria media*, with as many as 375 registered products (Table 3). This is followed by 271 in winter triticale, 226 in winter rye, and 200 in winter barley (MARD 2025). Consequently, the potential reduction in available herbicides due to regulatory changes is likely to be greatest in winter wheat. Conversely, winter barley is expected to experience the smallest reduction, with a projected decrease of only 59 registered herbicides. An increasingly serious problem will arise with the control of herbicide-resistant species, such as *Apera spica-venti* (Mayerová *et al.* 2018; Adamczewski *et al.* 2019; Marcinkowska *et al.* 2023; Bhattacharya *et al.* 2025).

fungicides is registered for the control of powdery mildew in this crop, compared to 165 in winter barley, and the fewest in winter rye, with only 39 registered fungicides – 279 fewer than those available for winter wheat (MARD 2025) (Table 4). The most significant phytopathogens include fungi from the genera *Fusarium* and *Puccinia*, along with *Zymoseptoria tritici* and *Parastagonospora nodorum* (Duba *et al.* 2018). For example, yield losses of winter wheat caused by pathogen infection can exceed 20% (Savary *et al.* 2019). Effective control of these diseases often relies on triazole-based fungicides such as epoxiconazole and tebuconazole, both of which face regulatory restrictions or phase-out in the EU (Marchand 2023a, b; Solé *et al.* 2024).

## Control of winter cereal pathogens

In Poland, pathogens that affect winter wheat have the greatest economic significance. A total of 318

## Control of pests in winter cereals

Pests pose the greatest economic significance on winter wheat plantations. For aphid control in this crop,

**Table 4.** Number of foliar fungicides registered in Poland for disease control in winter cereals (June 2025)

| Disease                      | Pathogen(s)                         | Current Registration | After Possible Withdrawal |
|------------------------------|-------------------------------------|----------------------|---------------------------|
| WINTER WHEAT                 |                                     |                      |                           |
| Brown leaf spot              | <i>Pyrenophora tritici-repentis</i> | 220                  | 161                       |
| Fusarium head blight         | <i>Fusarium</i> spp.                | 206                  | 148                       |
| Powdery mildew of cereals    | <i>Blumeria graminis</i>            | 318                  | 252                       |
| Cereal stem breakage         | <i>Oculimacula</i> spp.             | 62                   | 49                        |
| Brown rust                   | <i>Puccinia recondita</i>           | 315                  | 258                       |
| Yellow rust                  | <i>Puccinia striiformis</i>         | 120                  | 90                        |
| Septoria nodorum blotch      | <i>Parastagonospora nodorum</i>     | 175                  | 126                       |
| Wheat leaf banded septoriosi | <i>Mycosphaerella graminicola</i>   | 371                  | 273                       |

**Table 4.** Number of foliar fungicides registered in Poland for disease control in winter cereals (June 2025) – continuation

| Disease                               | Pathogen(s)  | Current Registration | After Possible Withdrawal |
|---------------------------------------|--|----------------------|---------------------------|
| WINTER TRITICALE                      |  |                      |                           |
| Brown leaf spot                       | <i>Pyrenophora tritici-repentis</i>                      | 48                   | 29                        |
| Fusarium head blight                  | <i>Fusarium</i> spp.                                     | 32                   | 12                        |
| Fusarium stem base and root rot       | <i>Fusarium</i> spp.                                     | 7                    | 7                         |
| Cereal stem breakage                  | <i>Oculimacula</i> spp.                                  | 18                   | 17                        |
| Powdery mildew of cereals             | <i>Blumeria graminis</i>                                 | 141                  | 128                       |
| Brown rust                            | <i>Puccinia recondita</i>                                | 161                  | 112                       |
| Yellow rust                           | <i>Puccinia striiformis</i>                              | 48                   | 30                        |
| Cereal rhynchosporiosis               | <i>Rhynchosporium secalis</i>                            | 53                   | 30                        |
| Septoria nodorum blotch               | <i>Parastagonospora nodorum</i>                          | 68                   | 38                        |
| Septoria leaf blotch                  | <i>Zymoseptoria tritici</i>                              | 168                  | 122                       |
| WINTER BARLEY                         |  |                      |                           |
| Sooty mould black mould on wheat ears | <i>Cladosporium herbarum</i> ,<br><i>Alternaria</i> spp. | 23                   | 23                        |
| Fusarium head blight                  | <i>Fusarium</i> spp.                                     | 55                   | 25                        |
| Fusarium stem base and root rot       | <i>Fusarium</i> spp.                                     | 8                    | 8                         |
| Cereal stem breakage                  | <i>Oculimacula</i> spp.                                  | 25                   | 17                        |
| Powdery mildew of cereals             | <i>Blumeria graminis</i>                                 | 165                  | 128                       |
| Barley rust                           | <i>Puccinia hordei</i>                                   | 142                  | 102                       |
| Cereal rhynchosporiosis               | <i>Rhynchosporium secalis</i>                            | 194                  | 153                       |
| WINTER RYE                            |  |                      |                           |
| Brown leaf spot                       | <i>Pyrenophora tritici-repentis</i>                      | 25                   | 7                         |
| Fusarium head blight                  | <i>Fusarium</i> spp.                                     | 22                   | 1                         |
| Fusarium stem base and root rot       | <i>Fusarium</i> spp.                                     | 1                    | 1                         |
| Cereal stem breakage                  | <i>Oculimacula</i> spp.                                  | 10                   | 9                         |
| Powdery mildew of cereals             | <i>Blumeria graminis</i>                                 | 39                   | 26                        |
| Brown rust                            | <i>Puccinia recondita</i>                                | 182                  | 136                       |
| Yellow rust                           | <i>Puccinia striiformis</i>                              | 3                    | 2                         |
| Cereal rhynchosporiosis               | <i>Rhynchosporium secalis</i>                            | 161                  | 126                       |
| Septoria nodorum blotch               | <i>Parastagonospora nodorum</i>                          | 1                    | 1                         |

Source: Own elaboration based on data from the Ministry of Agriculture and Rural Development

44 insecticides are registered, followed by 20 in winter triticale, and the fewest in winter barley and winter rye, with 15 preparations registered for each (Table 5). For several decades there have been no repellents registered for use in winter cereal crops to prevent damage caused by animals (MARD 2025). Among the pests of greatest economic importance are aphids (primarily due to their role as virus vectors) and *Oulema* spp. The most prevalent aphid species in cereal crops is *Rhopalosiphum padi*, which contributes significantly to virus transmission (Finlay and Luck 2011; Strażyński and Ruszkowska 2015). Losses caused by aphids due to viral infections can reach up to 30% (Singh *et al.* 2020).

## Application of microbiological agents in winter cereals

Currently, only three microbiological agents are registered in Poland for the protection of winter wheat against selected pathogens and pests (Table 6). For other winter cereal species, such options are not available (MARD 2025). Microbiological agents, despite their potential, must be both effective and economically viable. The adoption of diverse biocontrol methodologies is environmentally benign, harmless, and has enough potential to significantly boost plant production (Pandit *et al.* 2022; Šunjka and Mechora 2022; Bakr *et al.* 2025). Increasingly, research is being conducted on

**Table 5.** Number of insecticides registered in Poland for pest control in winter cereals (June 2025)

| Pests                          | Current Registration | After Possible Withdrawal |
|--------------------------------|----------------------|---------------------------|
| WINTER WHEAT                   |                      |                           |
| <i>Zabrus tenebrioides</i>     | 1                    | 1                         |
| Aphididae                      | 44                   | 25                        |
| <i>Oulema</i> spp.             | 44                   | 32                        |
| Gastropoda                     | 28                   | 28                        |
| <i>Eurygaster testudinaria</i> | 9                    | 9                         |
| WINTER TRITICALE               |                      |                           |
| <i>Zabrus tenebrioides</i>     | 1                    | 1                         |
| Aphididae                      | 20                   | 7                         |
| <i>Oulema</i> spp.             | 11                   | 3                         |
| Gastropoda                     | 28                   | 28                        |
| WINTER BARLEY                  |                      |                           |
| <i>Zabrus tenebrioides</i>     | 1                    | 1                         |
| Aphididae                      | 15                   | 4                         |
| <i>Oulema</i> spp.             | 6                    | 0                         |
| Gastropoda                     | 29                   | 29                        |
| WINTER RYE                     |                      |                           |
| <i>Zabrus tenebrioides</i>     | 1                    | 1                         |
| Aphididae                      | 15                   | 4                         |
| <i>Eurygaster testudinaria</i> | 3                    | 3                         |
| Gastropoda                     | 29                   | 29                        |

Source: Own elaboration based on data from the Ministry of Agriculture and Rural Development

**Table 6.** Microbiological preparations currently registered for use in winter wheat in Poland (June 2025)

| Pathogen/Pest            | Active Substances (Microorganism)                | Number of Preparations |
|--------------------------|--|------------------------|
| <i>Blumeria graminis</i> | <i>Bacillus amyloliquefaciens</i> strain QST 713 | 1                      |
| <i>Fusarium</i> spp.     | <i>Pythium oligandrum</i>                        | 1                      |
| Aleyrodidae              |  |                        |
| Thripidae                |  |                        |
| Tetranychidae            | <i>Beauveria bassiana</i> strain ATTC 74040      | 1                      |
| Elateridae               |  |                        |

Source: Own elaboration based on data from the Ministry of Agriculture and Rural Development

the effective utilization of natural enemies of pests (Dainese *et al.* 2017; Liu and Chen 2024).

## Use of resistant and tolerant varieties against winter cereal pathogens

The withdrawal of active substances from plant protection products in winter cereal production by the European Commission necessitates the accelerated introduction of new varieties that are resistant or tolerant to pathogens. This approach aligns with the principles of integrated plant protection and sustainable crop production (Góral *et al.* 2015; Rudnicki and Piekarczyk 2019; Dyda *et al.* 2022; Spetsov 2022; Dracatos

*et al.* 2023; Zhang *et al.* 2023b; Wallace *et al.* 2024; Han *et al.* 2025; Madajska *et al.* 2025). In Poland, the resistance levels of winter cereal varieties to key pathogens are assessed by the Research Centre for Cultivar Testing (RCCT). Varieties deemed suitable for integrated production (IP) are included in the National Register maintained by RCCT and are also featured in the List of Recommended Varieties for individual voivodeships. This classification indirectly contributes to reducing the reliance on chemical plant protection products. All winter cereal species are susceptible to powdery mildew of cereals and grasses. According to the 9-point RCCT resistance scale, an average resistance level (i.e., score of 5) to powdery mildew is recorded in 86.8% of all winter rye varieties, followed by winter barley (67.4%), winter wheat (52.4%), and winter triticale

(46.5%). A resistance level above the average is most common in winter triticale varieties (37.2%), followed by winter wheat (27.0%), winter barley (5.2%), and winter rye (4.4%). The highest proportion of varieties with low resistance is found in winter wheat and winter triticale (both 4.6%), while the lowest is observed in winter barley (2.1%) (RCCT 2025).

## Certified integrated production of winter cereals

In Poland, starting in 2023, the Ministry of Agriculture and Rural Development (MARD) implemented funding for a voluntary, yet certified Integrated Production (IP) system as part of the eco-scheme framework (Journal of Laws 2023, item 412). Upon its introduction in 2023, the area under certified winter wheat cultivation was 3,352 hectares, which expanded significantly to 61,345 hectares in 2024. A similar upward trend was observed for winter barley, with certified IP cultivation rising from 1,014 hectares in 2023 to 6,028 hectares in 2024 (MARD 2025). Prior to the introduction of the IP eco-scheme (2019–2022), the certified cultivation area for winter wheat remained limited, ranging between only 431 and 536 hectares (SPHSIS 2025). A key requirement of the IP system includes the mandatory use of certified seed from varieties that exhibit resistance or tolerance to pests, thereby contributing to a reduced need for plant protection products (RCCT 2025). Furthermore, only plant protection products listed on an officially maintained and regularly updated registry may be used in certified IP systems (Strażyński *et al.* 2024). Zieliński *et al.* (2024, 2025a,b) emphasize the importance of the Integrated Production (IP) system as one of the key tools supporting sustainable agricultural practices. However, they point out differences in the implementation of eco-schemes depending on the size of farms and their location.

## Challenges and future outlook

Simplification of winter cereal production technologies, coupled with climate change – including the absence of prolonged winter conditions – and the ongoing withdrawal of active substances (AS) in plant protection products by the European Commission, may significantly increase economic challenges related to pest management and crop protection (Strażyński and Ruskowska 2015; Strażyński *et al.* 2016; Goulet *et al.* 2023; Marchand 2023a, b). Over the past several decades, an increasing resistance of pests to the prolonged and unilateral use of active substances in agricultural

crops has been observed (Peterson *et al.* 2019; Nakka *et al.* 2019; Gruner *et al.* 2020; Gong *et al.* 2021; Synowiec *et al.* 2021; Luo *et al.* 2022; Waclawowicz *et al.* 2022; Wenda-Piesik *et al.* 2022; Stankiewicz-Kosyl *et al.* 2023) or the occurrence of mycotoxins in products (Kosicki *et al.* 2020). Further reductions in the availability of active substances, in the absence of effective alternative solutions, are likely to exacerbate this issue (Metcalf *et al.* 2024; Wynn and Webb 2022). This situation may lead to an increase in the occurrence of counterfeit plant protection products or those originating from illegal imports (Streløke 2018). Additionally, there have been reported cases of the use of formulations containing withdrawn active substances, as well as the misuse of approved products – such as their application on crops or against pests not listed on the product label.

The aforementioned conditions necessitate, among other measures, the rapid introduction of new crop varieties that are resistant or tolerant to pests. A notable example of breeding progress is the development of winter barley varieties with resistance and tolerance to viral infections (Zhang *et al.* 2023a; RCCT 2025). Significant emphasis is also being placed on the development and application of novel biopreparations (Villaverde *et al.* 2013; Bänziger *et al.* 2022; Marchand 2023c, d). However, the current number of available products remains insufficient. Promising alternatives under investigation include ionic liquids (Kaczmarek *et al.* 2019; Pernak *et al.* 2022; Marcinkowska *et al.* 2023).

In the foreseeable future, chemical plant protection will likely remain the primary strategy for managing pests in winter cereals. Consequently, there is a pressing need for innovative formulations of plant protection products – characterized by prolonged efficacy, high selectivity, and rapid environmental degradation – to ensure both effectiveness and safety for human health and the environment. In particular, such innovations are especially desirable in the form of seed treatments (Dufour *et al.* 2021). In addition to the need to introduce new varieties or innovative plant protection products (safe for humans and the environment, and at the same time effective), the improvement of agrotechnical methods is considered to be of great importance (Brévault and Clouvel 2019; Turner *et al.* 2021; Yamini *et al.* 2025).

Sustainable biological control is achievable under a favorable environment. This means that sustained investments in scientific research and innovation, socio-economic and ecological studies, and influencing consumers' value systems are imperative (Harding and Raizada 2015; Šunjka and Mechora 2022; Nchu 2024).

In light of the overproduction of cereal grains and the insufficient share of leguminous crops in Poland, it is necessary to promote crop diversification as

a means to improve crop rotation. Instead of relying solely on the rotation of active substances, the integration of crop rotation, breeding resistant varieties, and the application of non-chemical plant protection methods must be prioritized. To counteract the narrowing spectrum of PPPs, national authorities should simplify the approval process for low-risk and biological agents, offer incentives for companies registering innovative active substances, and strengthen monitoring of counterfeit products. Industries, on the other hand, should invest in research on biopesticides and resistant cultivars, and engage in farmer education on IPM (Sawińska *et al.* 2020). The question remains whether, with limited possibilities of chemical protection of winter cereals, will it be possible to produce food profitably. Non-chemical methods are usually more labor-intensive and expensive and do not always guarantee the expected effect. Nevertheless, the current systems of subsidies for agricultural crops allow for obtaining a beneficial economic effect.

## References

- Adamczewski K., Matysiak K., Kierzek R., Kaczmarek S. 2019. Significant increase of weed resistance to herbicides in Poland. *Journal of Plant Protection Research* 59 (2): 139–150. DOI: <https://doi.org/10.24425/jppr.2019.129293>
- Alix A., Lewis G. 2010. Guidance for the assessment of risks to bees from the use of plant protection products under the framework of Council Directive 91/414 and Regulation 1107/2009. *EPPO Bulletin* 40 (2): 196–203. DOI: <https://doi.org/10.1111/j.1365-2338.2010.02376.x>
- Bakr R., Abdelmoteleb A., Mendez-Trujillo V., Gonzalez-Mendoza D., Hewedy O. 2025. The potential of beneficial microbes for sustainable alternative approaches to control phytopathogenic diseases. *Microbiology Research* (16) 5: 105. DOI: <https://doi.org/10.3390/microbiolres16050105>
- Bänziger L., Kägi A., Vogelgsang S., Klaus S., Hebeisen T., Büttner-Mainik A., Sullam K. 2022. Comparison of thermal seed treatments to control snow mold in wheat and loose smut of barley. *Frontiers in Agronomy* 3. DOI: [10.3389/fagro.2021.775243](https://doi.org/10.3389/fagro.2021.775243)
- Bhattacharya S., Kumar Sen M., Hamouzová K., Košnarová P., Bharati R., Menendez J., Soukup J. 2025. Pyrooxulam Resistance in *Apera spica-venti*: An Emerging Challenge in Crop Protection. *Plants* 14 (1): 74. DOI: <https://doi.org/10.3390/plants14010074>
- Brévault T., Clouvel P. 2019. Pest management: Reconciling farming practices and natural regulations. *Crop Protection* 115: 1–6. DOI: <https://doi.org/10.1016/j.cropro.2018.09.003>
- Cilia N., Kandris I. 2023. Training in the evaluation of pesticides (plant protection products and active substances) according to Regulation (EC) No 1107/2009. Special Issue: EU-FORA Series 6: 21 (S1). DOI: <https://doi.org/10.2903/j.efsa.2023.e211007>
- Dainese M., Schneider G., Krauss J., Steffan-Dewenter I. 2017. Complementarity among natural enemies enhances pest suppression. *Scientific Reports* 7 (1): 8172. DOI: [10.1038/s41598-017-08316-z](https://doi.org/10.1038/s41598-017-08316-z)
- Dracatos P.M., Lu J., Sánchez-Martín J., Wulff B.B.H. 2023. Resistance that stacks up: engineering rust and mildew disease control in the cereal crops wheat and barley. *Plant Biotechnology Journal* 21 (10): 1938–1951. DOI: <https://doi.org/10.1111/pbi.14106>
- Duba A., Goriewa-Duba K., Wachowska U. 2018. A Review of the Interactions between Wheat and Wheat Pathogens: *Zygomoseptoria tritici*, *Fusarium* spp. and *Parastagonospora nodorum*. *International Journal of Molecular Sciences* 19 (4): 1138. DOI: <https://doi.org/10.3390/ijms19041138>
- Dufour T., Gutierrez Q., Bailly C. 2021. Sustainable improvement of seeds vigor using dry atmospheric plasma priming: evidence through coating wettability, water uptake and plasma reactive. *Journal of Applied Physics* 129 (8): 084902. DOI: <https://doi.org/10.1063/5.0037247>
- Dyda M., Tyrka M., Gołębiowska G., Rapacz M., Wędzony M. 2022. Genetic mapping of adult-plant resistance genes to powdery mildew in triticale. *Journal of Applied Genetics* 63: 73–86. DOI: <https://doi.org/10.1007/s13353-021-00664-x>
- OJEU 2009. L. 309. Official Journal of the European Union. Regulation (EC) No 1107/2009 of the European Parliament and of the Council of 21 October 2009 concerning the placing of plant protection products on the market and repealing Council Directives 79/117/EEC and 91/414/EEC.
- Finlay K.J., Luck J.E. 2011. Response of the bird cherry-oat aphid (*Rhopalosiphum padi*) to climate change in relation to its pest status, vectoring potential and function in a crop-vector-virus pathosystem. *Agriculture, Ecosystems & Environment* 144 (1): 405–421. DOI: <https://doi.org/10.1016/j.agee.2011.08.011>
- Freier B., Sellmann J., Strassemeyer J., Schwarz J., Klocke B., Kehlenbeck H., Zornbach W. 2013. Network of reference farms for plant protection - Annual Report 2012 – Analysis of Results of 2007 to 2012. *Berichte aus dem Julius Kühn-Institut* 172.
- Gong P., Chen D., Wang C., Li M., Li X., Zhang Y., Li X., Zhu X. 2021. Susceptibility of four species of aphids in wheat to seven insecticides and its relationship to detoxifying enzymes. *Frontiers in Physiology* 11: 623612. DOI: [10.3389/fphys.2020.623612](https://doi.org/10.3389/fphys.2020.623612)
- Goulet F., Aulagnier A., Hubert M. 2023. Strong withdrawal or weak withdrawal? Problematization of pesticides and categorization of their alternatives in Argentina, Brazil and France. p. 153–167. In: "New Horizons for Innovation Studies" (F. Goulet, D. Vinck, eds.), 332 pp. DOI: <https://doi.org/10.4337/9781803925554>
- Gruner P., Schmitt A.-K., Flath K., Schmiedchen B., Eifler J., Gordillo A., Schmidt M., Korzun V., Fromme F.J., Siekmann D., Tratwal A., Danielewicz J., Korbas M., Marciniak K., Krysztofik R., Niewińska M., Koch S., Piepho H.P., Miedaner T. 2020. Mapping stem rust (*Puccinia graminis* f. sp. *secalis*). Resistance in self-fertile winter rye populations. *Frontiers in Plant Science* 11: 667. DOI: [10.3389/fpls.2020.00667](https://doi.org/10.3389/fpls.2020.00667)
- Góral T., Ochodzki P., Walentyn-Góral D., Belter J., Majka M., Kwiatek M., Wiśniewska H., Bogacki J., Drzazga T., Ługowska B., Matysik P., Witkowski E., Rubrycki K., Woźna-Pawlak U. 2015. Odporność genotypów pszenicy ozimej na fuzariozę kłosów i akumulację toksyn fuzaryjnych w ziarnie scharakteryzowana za pomocą różnych typów odporności. *Biuletyn Instytutu Hodowli i Aklimatyzacji Roślin* 276: 19–37. DOI: <https://doi.org/10.37317/biul-2015-0017> (in Polish)
- Han G., Yan H., Li L., An D. 2025. Advancing wheat breeding using rye: a key contribution to wheat breeding history. *Trends in Biotechnology*. DOI: <https://doi.org/10.1016/j.tibtech.2025.03.008> (in press)
- Harding D.P., Raizada M.N. 2015. Controlling weeds with fungi, bacteria and viruses: A review. *Frontiers in Plant Science* 6: 659. DOI: <https://doi.org/10.3389/fpls.2015.00659>
- IAFP–NRI 2024. Institute of Agricultural and Food Economics – National Research Institute. Market Analysis. [Available on: <https://ierigz.waw.pl/publikacje/analizy-rynkowe>] [Accessed 25 Mar. 2025]
- IMWM–NRI 2025. Institute of Meteorology and Water Management – National Research Institute. [Available on: <https://>]

- modele.imgw.pl/cmm/?page\_id=44688] [Accessed 25 Mar. 2025]
- Journal of Laws of the Republic of Poland, item 455. Ustawa z dnia 8 marca 2013 r. o środkach ochrony roślin.
- Journal of Laws of the Republic of Poland, item 505. Rozporządzenie Ministra Rolnictwa i Rozwoju Wsi z dnia 18 kwietnia 2013 r. w sprawie wymagań integrowanej ochrony roślin.
- Journal of Laws of the Republic of Poland, item 412. Ustawa z dnia 8 lutego 2023 r. o Planie Strategicznym dla Wspólnej Polityki Rolnej na lata 2023–2027.
- Jørgensen L.N., Kudsk P., Ørum J.E. 2019. Links between pesticide use pattern and crop production in Denmark with special reference to winter wheat. *Crop Protection* 119. DOI: 10.1016/j.cropro.2019.01.024
- Kaczmarek D., Rzemieniecki T., Marcinkowska K., Pernak J. 2019. Synthesis, properties and adjuvant activity of docosate-based ionic liquids in pesticide formulations. *Journal of Industrial and Engineering Chemistry*: 78: 440–447. DOI: <https://doi.org/10.1016/j.jiec.2019.05.023>
- Kosicki R., Twarużek M., Dopierała P., Rudzki B., Grajewski J. 2020. Occurrence of mycotoxins in winter rye varieties cultivated in Poland (2017–2019). *Toxins* 12: 423. DOI: <https://doi.org/10.3390/toxins12060423>
- Lamichhane J.R., Dachbrodt-Saaydeh S., Kudsk P., Messéan A. 2016. Toward a Reduced Reliance on Conventional Pesticides in European Agriculture. *Plant Disease* 100 (1): 10–24. DOI: <https://doi.org/10.1094/PDIS-05-15-0574-FE>
- Liu T.-X., Chen X.-X. 2024. Biological control of aphids in china: successes and prospects. *Annual Review of Entomology* 70 (1): 401–419. DOI: 10.1146/annurev-ento-121423-012130
- Luo K., Zhao H., Wang X., Kang Z. 2022. Prevalent pest management strategies for grain aphids: opportunities and challenges. *Frontiers in Plant Science, Section Plant Pathogen Interactions* 12. DOI: <https://doi.org/10.3389/fpls.2021.790919>
- MacLeod A., Pautasso M., Jeger M.J., Haynes-Young R. 2010. Evolution of the international regulation of plant pests and challenges for future plant health. *Food Security* 2: 49–70. DOI: <https://doi.org/10.1007/s12571-010-0054-7>
- Madajska K., Tratwal A., Bocianowski J. 2025. Przydatność pszenżyta ozimego w różnych warunkach gospodarowania w świetle wymogów integrowanej ochrony oraz Europejskiego Zielonego Ładu. [The suitability of winter triticale in different farming conditions in the light of the requirements of integrated pest management and the European Green Deal]. *Progress in Plant Protection* 65 (1): 40–52. DOI: 10.14199/ppp-2025-006 (in Polish with English abstract)
- Marchand P.A. 2023a. EU Chemical Plant Protection Products in 2023: Current State and Perspectives. *Agrochemicals* 2 (1): 106–117. DOI: <https://doi.org/10.3390/agrochemicals2010008>
- Marchand P.A. 2023b. EU Chemical Plant Protection Products in 2023: Current State and Perspectives. *Agrochemicals* 2: 106–117. DOI: <https://doi.org/10.3390/agrochemicals2010008>
- Marchand P.A. 2023c. Evolution of plant protection active substances in Europe: the disappearance of chemicals in favour of biocontrol agents. *Environmental Science and Pollution Research* 30: 1–17. DOI: <https://doi.org/10.1007/s11356-022-24057-7>
- Marchand P.A. 2023d. BioControl Agents in Europe: Substitution Plant Protection Active Substances or a New Paradigm? *Agrochemicals* 2 (4): 538–550. DOI: <https://doi.org/10.3390/agrochemicals2040030>
- Marcinkowska M., Praczyk T., Niemczak M., Rzemieniecki T., Kaczmarek D.K., Łacka A., Pernak J. 2023. Herbicidal ionic liquids containing double or triple anions as a new potential tool for weed control including herbicide-resistant biotypes. *Crop Protection* 169: 106238. DOI: <https://doi.org/10.1016/j.cropro.2023.106238>
- MARD 2025. Ministry of Agriculture and Rural Development. [Available on: <https://www.gov.pl/web/rolnictwo/rejestr-rodkow-ochrony-roslin>] [Accessed: 25 Mar. 2025]
- Mayerová M., Mikulka J., Soukup J. 2018. Effects of selective herbicide treatment on weed community in cereal crop rotation. *Plant, Soil and Environment* 64 (9): 413–420. DOI: 10.17221/289/2018-PSE
- Metcalfé H., Storkey J., Hull R., Bullock J.M., Whitmore A., Sharp R.T., Milne A.E. 2024. Trade-offs constrain the success of glyphosate-free farming. *Scientific Reports* 14 (1): 8001. DOI: 10.1038/s41598-024-58183-8.
- Mie A., Rudén C. 2022. What you don't know can still hurt you – underreporting in EU pesticide regulation. *Environmental Health* 21: 79. DOI: <https://doi.org/10.1186/s12940-022-00891-7>
- Nakka S., Jugulam M., Peterson D., Asif M. 2019. Herbicide resistance: Development of wheat production systems and current status of resistant weeds in wheat cropping systems. *The Crop Journal* 7 (6): 750–760. DOI: <https://doi.org/10.1016/j.cj.2019.09.004>
- Nchu F. 2024. Sustainable biological control of pests: the way forward. *Applied Sciences* 14 (7): 2669. DOI: <https://doi.org/10.3390/app14072669>
- Oerke E.C. 2006. Crop losses to pests. *Journal of Agricultural Science* 144 (1): 31–43. DOI: <https://doi.org/10.1017/S0021859605005708>
- Pandit M.A., Kumar J., Gulati S., Bhandari N., Mehta P., Katyal R., Rawat C.D., Mishra V., Kaur J. 2022. Major Biological Control Strategies for Plant Pathogens. *Pathogens* 19: 273. DOI: 10.3390/pathogens11020273
- Pernak J., Niemczak M., Rzemieniecki T., Marcinkowska K., Praczyk T. 2022. Dicationic herbicidal ionic liquids comprising two active ingredients exhibiting different modes of action. *Journal of Agricultural and Food Chemistry* 70: 2545–2553. DOI: <https://doi.org/10.1021/acs.jafc.1c07750>
- Peterson M.A., Collavo A., Ovejero R., Shivrain V., Walsh M.J. 2017. The challenge of herbicide resistance around the world: a current summary. *Pest Management Science* 7 (10): 2246–2259. Special Issue: Global Herbicide Resistance Challenge. DOI: <https://doi.org/10.1002/ps.4821>
- RCCT 2025. Research centre for cultivar testing. selection of varieties for integrated crop production. [Available on: <https://www.coboru.gov.pl/pdo/ipr>] [Accessed 27 Mar. 2025]
- Rudnicki F., Piekarczyk M. 2019. Postęp hodowlany w polskim rejestrze odmian pszenicy ozimej w latach 1969–2016. Część II. Odporność na czynniki biotyczne i abiotyczne [Breeding progress in the polish register of varieties of the winter wheat in years 1969–2016. Part II. Resistance to biotic and abiotic stresses]. *Fragmenta Agronomica* 36 (2): 65–85. DOI 10.26374/fa.2019.36.17 (in Polish with English abstract)
- Savary S., Willocquet L., Pethybridge S.J., Esker P., McRoberts N., Nelson A. 2019. The global burden of pathogens and pests on major food crops. *Nature Ecology & Evolution* 3 (3): 430–439. DOI: <https://doi.org/10.1038/s41559-018-0793-y>
- Sawińska Z., Świtek S., Głowicka-Wołoszyn R., Kowalczewski P.Ł. 2020. Agricultural Practice in Poland Before and after Mandatory IPM Implementation by the European Union. *Sustainability* 12 (3): 1107. DOI: 10.3390/su12031107
- Singh B., Simon A., Halsey K., Kurup S., Clark S., Aradottir G.I. 2020. Characterisation of bird cherry-oat aphid (*Rhopalosiphum padi* L.) behaviour and aphid host preference in relation to partially resistant and susceptible wheat landraces. *Annals of Applied Biology* 177 (2): 184–194. DOI: 10.1111/aab.12616
- Slunge D., Miguel M., Lindahl L., Backhaus T. 2023. The implementation of the substitution principle in European chemical legislation: a comparative analysis. *Environmental Sciences Europe* 35: 107. DOI: <https://doi.org/10.1186/s12302-023-00817-1>

- Solé M., Brendel S., Aldrich A., Dauber J., Ewald J., Duquesne S., Gottschalk E., Hoffmann J., Kuemmerlen M., Leake A., Matezki S., Meyer S., Nabel M., Natal-da-Luz T., Pieper S., Piselli D., Rigal S., Roß-Nickoll M., Schäffer A., Settele J., Sigmund G., Sotherton N., Wogram J., Messner D. 2024. Assessing in-field pesticide effects under European regulation and its implications for biodiversity: a workshop report. *Environmental Sciences Europe* 36: 153. DOI: <https://doi.org/10.1186/s12302-024-00977-8>
- SP 2024. Statistics Poland. Zużycie środków ochrony roślin wg zharmonizowanej klasyfikacji substancji. Rocznik Statystyczny Rolnictwa 2010–2024. Wyd. GUS, Warszawa. (in Polish)
- SP 2025. Statistics Poland. [Available on: <https://stat.gov.pl/obszary-tematyczne/rolnictwo-lesnictwo/uprawy-rolnicze-wynikowe-szacunek-glownych-ziemnioplodow-rolnych-i-ogrodnictwa-w-2024-roku,5,23.html>] [Accessed 25 Mar. 2025]. (in Polish)
- Spetsov P. 2022. Resistance to pathogens in wheat-rye and triticale genetic stocks. *Journal of Plant Pathology* 104 (8). DOI: [10.1007/s42161-021-01019-5](https://doi.org/10.1007/s42161-021-01019-5)
- SPHSIS 2025. State Plant Health and Seed Inspection Service. Available on: <https://www.gov.pl/web/piorin/integrowana-produkcja-roslin>] [Accessed 25 Mar. 2025]
- Stankiewicz-Kosyl M., Haliniarz M., Wrochna M., Obrępałska-Stęplowska A., Kuc P., Łukasz J., Wińska-Krysiak M., Wrzesińska-Krupa B., Puła J., Podsiadło C., Domaradzki K., Piekarczyk M., Bednarczyk M., Marcinkowska K. 2023. Occurrence and mechanism of *Papaver rhoas* als inhibitors resistance in Poland. *Agriculture* 13 (1): 82. DOI: <https://doi.org/10.3390/agriculture13010082>
- Strażyński P., Ruskowska M. 2015. The life cycle functional response of *Rhopalosiphum padi* (L.) to higher temperature: territorial expansion of permanent parthenogenetic development as a result of warmer weather conditions. *Journal of Plant Protection Research* 55 (2): 162–165. DOI: [10.1515/jppr-2015-0021](https://doi.org/10.1515/jppr-2015-0021)
- Strażyński P., Ruskowska M., Krówczyńska A. 2016. Aphids of the genus *Diuraphis* caught by Johnson suction trap in Poznań, Poland. *Journal of Plant Protection Research* 56 (4): 328–330. DOI: <https://doi.org/10.1515/jppr-2016-0053>
- Strażyński P., Tratwal A., Sosnowska D., Mrówczyński M. 2024. Integrowana ochrona roślin – stan i określenie zagrożeń w kontekście wyzwań związanych z wdrażaniem nowej perspektywy Unii Europejskiej na lata 2023–2027. [Integrated pest management – status and identification of risks in the context of challenges related to the implementation of the new European Union perspective for 2023–2027]. *Progress in Plant Protection* 64 (4). DOI: [10.14199/ppp-2024-016](https://doi.org/10.14199/ppp-2024-016)
- Strelake M. 2018. Illegal trade of plant protection products: a highly profitable way to smuggle chemicals. *Journal of Consumer Protection and Food Safety* 13: 255–256. DOI: <https://doi.org/10.1007/s00003-018-1177-6>
- Synowiec A., Jop B., Domaradzki K., Podsiadło C., Gawęda D., Wacławowicz R., Wenda-Piesik A., Nowakowski M.M., Bocianowski J., Marcinkowska K., Praczyk T. 2021. Environmental factors effects on winter wheat competition with herbicide-resistant or susceptible silky bentgrass (*Apera spica-venti* L.). *Agronomy* 11 (5): 871. DOI: <https://doi.org/10.3390/agronomy11050871>
- Šunjka D., Mechora Š. 2022. An alternative source of biopesticides and improvement in their formulation – recent advances. *Plants* 11: 3172. DOI: <https://doi.org/10.3390/plants11223172>
- Turner J.A., Chantry T., Taylor M.C., Kennedy M.C. 2019. Changes in agronomic practices and incidence and severity of diseases in winter wheat in England and Wales between 1999 and 2019. *Plant Pathology* 70 (8): 1759–1778. DOI: <https://doi.org/10.1111/ppa.13433>
- Wacławowicz R., Tendziagolska E., Synowiec A., Bocianowski J., Podsiadło C., Domaradzki K., Marcinkowska K., Kwiecińska-Poppe E., Piekarczyk M. 2022. Competition between winter wheat and cornflower (*Centaurea cyanus* L.) Resistant or susceptible to herbicides under varying environmental conditions in Poland. *Agronomy* 12 (11): 2751. DOI: <https://doi.org/10.3390/agronomy12112751>
- Wallace S., Chhabra B., Dong Y., Ma X., Coleman G., Tiwari V., Rawat N. 2024. Exploring Fusarium head blight resistance in a winter triticale germplasm collection. *Journal of Plant Registration* 18 (3): 457–465. DOI: <https://doi.org/10.1002/plr2.20392>
- Wenda-Piesik A., Synowiec A., Marcinkowska K., Wrzesińska B., Podsiadło C., Domaradzki K., Kuc P., Kwiecińska-Poppe E. 2022. Intra- and interspecies competition of blackgrass and wheat in the context of herbicidal resistance and environmental conditions in Poland. *Scientific Reports* 12 (8720): 1–14. DOI: <https://doi.org/10.1038/s41598-022-12777-2>
- Wynn S., Webb E. 2022. Impact assessment of the loss of glyphosate within the EU: a literature review. *Environmental Sciences Europe* 34: 91. DOI: <https://doi.org/10.1186/s12302-022-00667-3>
- Villaverde J.J., Sevilla-Morán B., Sandín-España P., López-Goti C., Alonso-Prados J.L. 2013. Biopesticides in the framework of the European Pesticide Regulation (EC) No. 1107/2009. *Pest Management Science* 70 (1): 2–5. DOI: <https://doi.org/10.1002/ps.3663>
- Yamini V., Singh K., Antar M., El Sabagh A. 2025. Sustainable cereal production through integrated crop management: a global review of current practices and future prospects. *Frontiers for Sustainable Food Systems*. Section: Crop Biology and Sustainability 9. DOI: <https://doi.org/10.3389/fsufs.2025.1428687>
- Zieliński M., Gołębiewska B., Adamski M., Sobierajewska J., Tyburski J. 2024. Adaptation of eco-schemes to Polish agriculture in the first year of the EU CAP 2023–2027. *Economics and Environment* 89 (2): 817. DOI: [10.34659/eis.2024.89.2.817](https://doi.org/10.34659/eis.2024.89.2.817)
- Zieliński M., Zalewski A., Łaba S. 2025a. Sustainability conditions of Polish agriculture in the context of the use of plant protection products, as compared to other European Union countries. Economic aspects. *Journal of Plant Protection Research* 65 (1): 45–60. DOI: [10.24425/jppr.2025.153818](https://doi.org/10.24425/jppr.2025.153818)
- Zieliński M., Józwiak W., Żak A., Rokicki T. 2025b. Development of Eco-Schemes as an Important Environmental Measure in Areas Facing Natural or Other Specific Constraints Under the Common Agriculture Policy 2023–2027: Evidence from Poland. *Sustainability* 17: 2781. DOI: <https://doi.org/10.3390/su17062781>
- Zhang M., Hong Y., Zhu J., Pan Y., Zhou H., Lv C., Guo B., Wang F., Xu R. 2023a. Molecular insights into the responses of barley to yellow mosaic disease through transcriptome analysis. *BMC Plant Biology*: 267. DOI: <https://doi.org/10.1186/s12870-023-04276-x>
- Zhang Z., Li Y., Li X., Zhu X., Zhang Y. 2023b. Efficacy of imidacloprid seed treatments against four wheat aphids under laboratory and field conditions. *Plants* 12: 238. DOI: <https://doi.org/10.3390/plants12020238>