

OCCURRENCE OF EYESPOT ON WINTER WHEAT IN THE CENTRAL-SOUTHERN REGION OF POLAND

Mariola Głazek*

Institute of Plant Protection – National Research Institute, Sońnicowice Branch
Gliwicka 29, 44-153 Sońnicowice, Poland

Received: April 23, 2009

Accepted: October 28, 2009

Abstract: Eyespot is one of the most important diseases of economical significance in wheat production in the regions of Śląskie and Opolskie voivodeships where it presents a permanent problem. The aim of performed experiment was to collect data on the occurrence and characteristics of *Oculimacula yallundae* and *O. acuformis* strains in these regions in the years 2003–2008, and to perform their taxonomic characterization. The studies originated as an attempt to find the explanation of processes related to winter wheat infection by casual agents of this disease in the Śląskie and Opolskie Provinces.

Key words: eyespot, *Oculimacula yallundae*, *O. acuformis*, infection index, PCR, laboratory analysis

INTRODUCTION

Eyespot is one of the most important diseases of economical significance in wheat production in the regions of moderate climate (Scott and Hollins 1974). It is a commonly occurring disease in winter wheat cultures in the regions of Śląskie and Opolskie voivodeships. Its intensity may be variable, however it presents a permanent problem in these regions (Głazek *et al.* 2008; Jaczewska-Kalicka 1998; Korbias 2008). It is caused by pathogenic fungi *Oculimacula yallundae* (type W) and *O. acuformis* (type R) (Nirenberg 1980; Nicholson *et al.* 1997). Epidemiological differences between these two species are not clear

and the disease development is difficult to predict (Fitt and Goulds 1988). The aim of performed experiment was to collect data on the occurrence and characteristics of *O. yallundae* and *O. acuformis* strains in the regions of Śląskie and Opolskie voivodeships in 2003–2008 and to perform their taxonomic characterization. Numerous plants with both initial and advanced symptoms of eyespot occurring on winter wheat cultivars at growth stage of late milk BBCH 73–77 were collected (Fig. 1) (Wan *et al.* 2005). The studies originated as an attempt to find the explanation of processes related to winter wheat infection by casual agents of the disease in the Śląskie and Opolskie provinces.



Fig. 1. Initial and advanced symptoms of eyespot on winter wheat at BBCH 73–75

*Corresponding address: m.glazek@ior.gliwice.pl

MATERIALS AND METHODS

Each year (2003–2008) winter wheat plants were taken for analysis from 5–8 chosen plantations. Those were experimental plantations, small area plantations and large commercial plantations as well (Table 1).

Table 1. Derivation of cultures used for mycological characterization of eyespot. A list of analyzed plantations

Year	Location	Number of plantations	Date of sampling
2003	śląskie	7	75–77 BBCH
2004	śląskie	5	75–77 BBCH
2005	śląskie	7	75–77 BBCH
	opolskie	1	75–77 BBCH
2006	śląskie	4	75–77 BBCH
	opolskie	1	75–77 BBCH
2007	śląskie	6	75–77 BBCH
	opolskie	2	75–77 BBCH
2008	śląskie	4	75–77 BBCH
	opolskie	1	75–77 BBCH

Visual analysis of 100 plants randomly taken from each plantation non – treated with fungicides was performed at 75–77 BBCH according to EPPO standards PP 1/28 (3).

From plant fragments of diseased tissues mycelium was taken in the laboratory for culturing the fungus on artificial medium (PDA) (Bojarczuk 1970). After obtaining pure cultures discs of 5 mm diameter were cut out and transferred to PDA plates. Fungus cultures showing typical for *O.yallundae* (type W) or *O.acuformis* (type R) morphology and growth rate were chosen (Scott *et al.*

1975; Creighton *et al.* 1988, 1991; Creighton 1989) for the further study.

Culturing the fungus obtained from excised culture discs was performed at uniform conditions of light and temperature of 18–19°C. At fixed time intervals measurements of linear growth of cultures were done after 5, 10 and 15 days from the transferring date. The morphology of cultures was compared with literature data (Nicholson *et al.* 1991; Nirenberg 1981; Dyer *et al.* 1994). Fungi from these cultures were grown on water agar to obtain a sporulation (Bojarczuk 1970). In the years 2002, 2003 and 2007, 2008 PCR Multiscreen Cereal Stem Base Disease Analysis was conducted to confirm visual diagnosis. For this purpose plants showing disease symptoms were chosen and send to a specialized laboratory NeoAdgen in Scotland (Smith 1990; Collet *et al.* 1992; Turner 1999). In 2008 a study based on PCR technique was also done in the Department of Mycology Institute of Plant Protection – National Research Institute in Poznań to confirm laboratory results (Turner *et al.* 1999).

RESULTS AND DISCUSSION

In the recent years, differences were observed with respect to the development and level of infection of wheat at the late milk growth stage of BBCH 75–77.

Table 2 shows examples of the level of infection in 2002–2003 and 2007–2008. The percentage of plants with disease symptoms on stems ranged from 26 to 100. Harmfulness of the disease, however, is assessed based on the rate of plants with medium and high level of infection (Scott *et al.* 1974). In order to compare the *Oculimacula* spp. infection on different plantations, infection pressure index was determined according to the EPPO standard

Table 2. Occurrence of eyespot in winter wheat selected plantations

Sośnicowice 200

No.	Localization/ cultivar		Date of sampling	<i>Oculimacula</i> spp.				
				% infected plants				infection index*, according to EPPO P1/28(3)
				infection severity			sum	
				slight (1)	moderate (2)	severe (3)		
1	śląskie	Mikon	73 BBCH	22	8	0	30	0,12
2	śląskie	Mikon	77 BBCH	26.7	38.9	5.5	71.1	0.41
3	śląskie	Zyta	75 BBCH	10	90	0	100	0.70

Sośnicowice 2002

Sośnicowice 200

No.	Localization/ cultivar		Date of sampling	<i>Oculimacula</i> spp .				
				% infected plants				infection index*, according to EPPO P1/28(3)
				infection severity			sum	
				slight (1)	moderate (2)	severe (3)		
1	śląskie	Mikon	77 BBCH	61	36	3	100	0.45
2	śląskie	Mikon	77 BBCH	91	9	0	100	0.30
3	śląskie	Kobra	77 BBCH	65	24	11	100	0.45
4	śląskie	Kobra	77 BBCH	35	13	6	54	0.25
5	śląskie	Mikon	77 BBCH	22	13	1	36	0.16

Sośnicowice 2003

continued table 2

Sośnicowice 2007

No.	Localization/ cultivar		Date of sampling	<i>Oculimacula</i> spp.				
				% infected plants			infection index*, according to EPPO P1/28(3)	
				infection severity				sum
				slight (1)	moderate (2)	severe (3)		
1	śląskie	Turnia	75–77 BBCH	11.25	57.75	9.75	78.75	0.56
2	śląskie	Nadobna	75–77 BBCH	13.25	62	8	83.25	0.58
3	śląskie	Finezja	75–77 BBCH	19.75	13	0.25	33	0.15
4	śląskie	Zyta	75–77 BBCH	28.50	30.50	1.75	53.00	0.32
5	opolskie	Buteo	75–77 BBCH	26	52	0	78	0.46
6	opolskie	Bogatka	75–77 BBCH	28	45	5	68	0.45
7	śląskie	Rubens	75–77 BBCH	19.00	18.50	2.25	39.75	0.21

Sośnicowice 2008

No.	Localization/ cultivar		Date of sampling	<i>Oculimacula</i> spp.				
				% infected plants			infection index*, according to EPPO P1 28(3)	
								infection severity
				slight (1)	moderate (2)	severe (3)		sum
1	śląskie	Rubens	75–77 BBCH	26.5	30.25	6.5	63.25	0.36
2	śląskie	Nadobna	75–77 BBCH	25.75	39.75	11.25	54.75	0.48
3	opolskie	Nadobna	75–77 BBCH	9	11	79	99	0.90
4	opolskie	Bogatka	75–77 BBCH	20	25	53	98	0.77
5	opolskie	Finezja	75–77 BBCH	15	33	48	96	0.77
6	śląskie	Turnia	75–77 BBCH	29.25	9.50	0.75	39.50	0.15
7	ślaskie	Zyta	75–77 BBCH	25	1	0	26	0.07

$$*) x = \frac{(n(II) \times 0.25) + (n(III) \times 0.75) + n(IV)}{n(I + II + III + IV)}$$

x – infection index

(I) – non-infected

(II) – slight

(III) – moderate

(IV) – severe

n – number of assessed plants

Table 3. Rate of linear growth of *O. yallundae* and *O. acuformis* cultures – means for 2003–2008

Year	Average growth of type W [mm]		
	5 days	10 days	15 days
2003	10.19	29.56	43.22
2004	12.00	26.81	46.91
2005	19.23	39.73	51.18
2006	12.03	24.15	35.73
2007	11.42	21.87	31.01
2008	9.71	22.32	32.99
Means for 2003–2008	12.43	27.41	40.17
Year	average growth of type R [mm]		
	5 days	10 days	15 days
2003	8.69	15.24	22.08
2004	7.41	14.06	22.06
2005	8.54	13.46	20.66
2006	7.75	12.38	23.25
2007	8.57	10.61	13.60
2008	6.75	10.68	13.27
Means for 2003–2008	7.95	12.74	19.15

P1/28 (3) (Table 2). The lowest infection index of 0.07 was found for Zyta cultivar grown in the Silesian Province and the highest of 0.90 for Nadobna in Opolskie in 2008. The average calculated infection pressure index was 0.41.

It indicates that the pathogen regularly infects about half of wheat plants to the extent that significantly weakens their condition.

The most important element of performed research was to identify fungal species of causal disease agents, as it could be of significance in the infection process, and in a delayed appearance of disease symptoms.

In each year of investigation from chosen plantations of winter wheat on which the occurrence of eyespot was observed, plants for laboratory analysis were taken. From fragments of diseased tissues isolations were made, and from obtained cultures were chosen those exhibiting typical characteristics for *O. yallundae* or *O. acuformis*. Measurements of linear growth were made. In case of *O. yallundae* after 15 days it amounted to 31–51 mm, (type W) and for *O. acuformis* (type R) it was 13–22 mm (Table 3).

The difference in linear growth means for 6 years for *O. yallundae* and *O. acuformis* was 20 mm (Fig. 2). In each year much faster growth of *O. yallundae* was evident. On figure 3 a, b are shown 15 days old cultures of types W and R. The next figure 4a–e shows differences in morphology of cultures of W and R types evident after about

4 weeks from transferring. Cultures of *O. yallundae* were characterized by darker colour and even colony margins. Cultures of *O. acuformis* are lighter, margins were uneven and latter they might be “fethery”. (Fig. 4f). Cultures of both strains had a button – like protrusion in their central part which is confirmed by other authors (Bojarczuk 1970; Nirenberg 1981; Bateman *et al.* 1990).

To confirm the correct classification of obtained fungi to the species further methods were applied. Among others, sporulation was observed on water agar and it was stated that sporulations of *O. yallundae* was more abundant, spores were shorter and slightly curved, and *O. acuformis* had longer spores and sometimes slightly curved at the end (Nirenberg 1981; Korbas 2008; Bojarczuk 1970) (Fig. 5a, b). In 2003–2008 it was stated that mean length of *O. yallundae* spores was 61,2 micrometers, and for *O. acuformis* it was 70 micrometers (Table 4). According to Nirenberg (1981) length of conidia of *yallundae* was from 35 to 80 micrometers and for *acuformis* 43–120 micrometers.




Fig. 2. Growth rate of *O. yallundae* and *O. acuformis* cultures on PDA medium. Average of 2003–2008

Table 4. Mean length of W and R types conidia from studied cultures of *O. yallundae* and *O. acuformis*

Year	Mean length of conidia [μm]	
	Type W <i>O. yallundae</i>	Type R <i>O. acuformis</i>
2003	57.2	64.1
2004	60.3	38.3
2005	59.0	85.5
2006	86.6	98.6
2007	53.9	60.4
2008	56.0	71.1
mean	61.2	70.0





Fig. 3. 15-days old cultures: a – *O. acuformis*, b – *O. yallundae*




Fig. 4. Morphological differences of *Oculimacula* cultures 4 weeks after inoculation:
a–c *O. yallundae*, d–f *O. acuformis*

Further studies were performed using PCR technique. The use of specific for W or R types primers enabled the confirmation laboratory studies and their correct classification (Korbas 2008; Nicholson 1997). On figure 6 spectrum showing the results of PCR analysis with specific primers for *O. yallundae* and *O. acuformis* is presented.

A highly significant element for elucidation of differentiated wheat infection at the milky ripening stage BBCH 71–77 were PCR studies in which the participation of types W and R were determined. At this stage the


species *O. acuformis* was clearly prevalent and comprised 70–100% as compared to *O. yallundae* occurring in 0–30% (Fig. 7). In investigated plants from 26 plantations during 6 years on 24 plantations type R was present. Type W was found on 4 plantations (Table 5).

Laboratory studies and PCR tests confirmed the participation of two species of *Oculimacula* in the infection process of winter wheat in the discussed regions.

The prevailing presence of *O. acuformis* can be explained by a late appearance of disease symptoms, as this



Fig. 5. Conidiospores of *O. yallundae* (a), *O. acuformis* (b)



M – ladder size marker, 1, 3, 5, 7, 10, 12, 13 – isolates of *O. yallundae* and 2, 4, 6, 8, 9, 11 – isolates of *O. acuformis*

Fig. 6. Spectrum showing the results of PCR analysis of *O.* cultures with specific primers for *O. yallundae* and *O. acuformis*




Fig. 7. Per cent share of *O. yallundae* and *O. acuformis* in fungal population of *Oculimacula* spp. in 2002, 2003 and 2007, 2008

Table 5. Results of PCR Multiscreen analysis of winter wheat plants from 22 different plantations

Year	Location	Type*		Development stage
		W <i>O. yallundae</i>	R <i>O. acuformis</i>	
2002	śląskie	0	2	71 BBCH
		0	5	77 BBCH
		5	5	75 BBCH
2003	śląskie	0	3	77 BBCH
		0	1	77 BBCH
		0	2	77 BBCH
		0	0	77 BBCH
		0	0	77 BBCH
2007	śląskie	0	5	75–77 BBCH
		0	5	75–77 BBCH
		0	2	75–77 BBCH
		0	4	75–77 BBCH
		0	2	75–77 BBCH
	opolskie	3	5	75–77 BBCH
		5	5	75–77 BBCH
2008	śląskie	0	1	75–77 BBCH
		2	1	75–77 BBCH
		0	1	75–77 BBCH
	opolskie	0	5	75–77 BBCH
		2	5	75–77 BBCH
	śląskie	0	0	75–77 BBCH
	opolskie	0	5	75–77 BBCH

* 0–5 scale: 0 – disease not detected; 1–5 – increasing level of occurrence

species has a longer period of latent infection and develops more slowly than the species *O. yallundae* (Ray *et al.* 2004).

A prevalence of *O. acuformis* and restricted occurrence *O. yallundae* may be related to the common use of carben-dazim for controlling eyespot (Bateman 2002; Bierman *et al.* 2002; Birchmore *et al.* 1992; Blein *et al.* 2009; Dayer *et al.* 1995; Głazek *et al.* 1993; Korbas *et al.* 1999; Smith *et al.* 1994).

Stating a regular occurrence in 2003–2008 of *O. acuformis* in the studied regions also permits to elucidate the problem of late appearance of disease symptoms at the beginning of ripening when the intensity of disease symptoms by this species considerably increases and endangers yielding of winter wheat.

Results of the presented above research point at the need of their continuation because this information presents a valuable indications for agricultural practice, permitting a proper application of wheat protection against eyespot.

REFERENCES

- Bateman G.L. 2002. Long-term performance of fungicides applied to control eyespot in winter wheat. *Ann. Appl. Biol.* 141: 29–33.
- Bateman G.L., Gould A., Ainsley A. 1990. comparison of infection by W-type and R-type isolates of the eyespot pathogen, *Pseudocercospora herpotrichoides*, in relation to host species and cultivar, temperature and watering regime in controlled environments. *J. Phytopathol.* 129: 271–282.
- Bierman S.M., Fitt B.D.L., van den Bosch F., Bateman G.L., Jenkyn J. F., Welham S. J. 2002. Changes in populations of the eyespot fungi *Tapesia yallundae* and *T. acuformis* under different fungicide regimes in successive crops of winter wheat, 1984–2000. *Plant Pathol.* 51: 191–201.
- Birchmore R.J., Ashman P.I., Stanley S., Russell P.E. 1992. Effects of crop history on sensitivity to prochloraz of *Pseudocercospora herpotrichoides* isolates from cereals in western Europe. Brighton Crop Protection Conference – Pests and Diseases: 183–189.
- Blein M., Levrel A., Lemoine J., Gautier V., Chevalier M., Barloy D. 2009. *Oculimacula yallundae* lifestyle revisited: relationships between the timing of eyespot symptom appearance, the development of the pathogen and the responses of infected partially resistant wheat plants. *Plant Pathol.* 58: 1–11.
- Bojarczuk J. 1970. Badania nad odpornością pszenicy ozimej na łamliwość źdźbła (*Cercospora herpotrichoides* Fron). cz. I. Znaczenie gospodarcze choroby. *Biologia patogena. Warunki infekcji. Hodowla Roślin, Aklimatyzacja i Nasiennictwo* 14, 4: 327–341.
- Collett M., Clark J.S.C., Kendall S.J., Hollomon D.W. 1992. Field evaluation of an immunodiagnostic assay for cereal eyespot. Brighton Crop Protection Conference – Pests and Diseases 7A-2: 697–703.
- Creighton N.F., Bateman G.L. 1988. Improved diagnosis of eyespot pathotypes: Application of a new method. Brighton Crop Protection Conference – Pests and Diseases 8C-27: 1023–1027.
- Creighton N.F., Bateman G.L. 1991. Problems in identifying pathogenicity types of *Pseudocercospora herpotrichoides* by colony morphology arising from the source and concentration of the nutrient agar. *Mycol. Res.* 95: 253–254.
- Creighton N.F. 1989. Identification of W-type and R-type isolates of *Pseudocercospora herpotrichoides*. *Plant Pathol.* 38: 484–493.
- Dyer P.S., Lucas J.A. 1995. Incidence of apothecia of *Tapesia yallundae* at set-aside sites in England and sensitivity of the

- ascospore offspring to the fungicides benomyl and prochloraz. *Plant Pathol.* 44: 796–804.
- Dyer P.S., Papaikonomou M., Lucas J.A., Peberdy J. F. 1994. Isolation of R-type progeny of *Tapesia yallundae* from apothecia on wheat stubble in England. *Plant Pathol.* 43: 1039–1044.
- Efficacy Evaluation of Plant Protection Products 2006. EPPO Standards PP 1/28 (3): 1–3.
- Fitt B.D.L., Goulds A. 1988. Eyespot (*Pseudocercospora herpotrichoides*) epidemiology in relation to prediction of disease severity and yield loss in winter wheat – a review. *Plant Pathol.* 37: 311–328.
- Głazek M., Pietryga J. 1993. Chemiczne zwalczanie łamliwości żdźbła na pszenicy ozimej i ocena szybkiego testu membranowego do określenia nasilenia choroby. *Materiały 33. Sesji Nauk. Inst. Ochr. Roślin, cz. II*: 124–129.
- Głazek M., Kościelniak W., Pyziak K. 2008. Choroby podstawy żdźbła pszenicy ozimej na terenie województwa opolskiego. *Materiały Szkoleniowe Opolskiego Ośrodka Doradztwa Rolniczego 2008*, 21 pp.
- Goulds A., Fitt B.D. 1991. Prediction of eyespot severity on winter wheat or winter barley inoculated with W-type or R-type isolates of *Pseudocercospora herpotrichoides*. *J. Phytopathol.* 132: 105–115.
- Jaczewska-Kalicka A. 1998. Risk of damage to winter wheat by *Pseudocercospora herpotrichoides* (Fron) Deighton, the causal agent of eyespot. II. Results of plot experiments. *J. Plant Protection Res.* 38 (2): 122–138.
- Korbas M., Remlein-Starosta D. 1999. Odporność sprawcy łamliwości żdźbła zbóż (*Pseudocercospora herpotrichoides* (Fron.) Deighton) na wybrane fungicydy benzimidazolowe. *Prog. Plant Protection/Post. Ochr. Roślin* 39 (1): 360–366.
- Korbas M. 2008. Epidemiologia łamliwości żdźbła pszenicy ozimej w Polsce. *Rozpr. Nauk. Inst. Ochr. Roślin* 18, 68 pp.
- Nicholson P., Rezanoor H.N. 1991. Occurrence of *Tapesia yallundae* apothecia on field and laboratory-inoculated material and evidence for recombination between isolates. *Plant Pathol.* 40: 626–634.
- Nicholson P., Rezanoor H.N., Simpson D.R., Joyce D. 1997. Differentiation and quantification of the cereal eyespot fungi *Tapesia yallundae* and *Tapesia acuformis* using a PCR assay. *Plant Pathol.* 46: 842–856.
- Nirenberg H.I. 1981. Differenzierung der Erreger der Halmbruchkrankheit I. Morphologie. *Z. Pflanzenkrankh. Pflanzensch.* 88 (5): 241–248.
- Ray R.V., Jenkinson P., Edwards S.G. 2004. Effects of fungicides on eyespot, caused predominantly by *Occulimacula acuformis*, and yield of early-drilled winter wheat. *Crop Protect.* 23: 1199–1207.
- Scott P.R., Hollins T.W., Muir P. 1975. Pathogenicity of *Cercospora herpotrichoides* to wheat, barley, oats and rye. *Trans. Br. Mycol. Soc.* 65 (3): 529–538.
- Scott P.R., Hollins T.W. 1974. Effects of eyespot on the yield of winter wheat. *Ann. Appl. Biol.* 78: 269–279.
- Smith C.M., Saunders D.W., Allison D.A., Johnson C.E.B., Labit B., Kendall S.J., Hollomon D.W. 1990. Immunodiagnostic assay for cereal eyespot; Novel technology for disease detection. *Proceedings Brighton Crop Protection Conference – Pests and Diseases*: 763–770.
- Smith C.M., Jańczak C., Korbas M., Pokacka Z., Głazek M., Weber Z., Kuś J., Mróz A., Zawadzki T. 1994. Occurrence, development and control by fungicides of *Septoria nodorum*, *S. tritici*, and *Pseudocercospora herpotrichoides* on wheat in Poland in 1993. *Brighton Crop Protection Conference – Pests and Diseases 3D-14*: 426–429.
- Turner A.S., O'Hara R.B., Rezanoor H.N., Nuttall M., Smith J. N., Nicholson P. 1999. Visual disease and PCR assessment of stem base diseases in winter wheat. *Plant Pathol.* 48: 742–748.
- Wan A.M., Bock C.H., Fitt B.D.L., Harvey J.L., Jenkyn J.F. 2005. Development of *Oculimacula yallundae* and *O. acuformis* (eyespot) on leaf sheaths of winter wheat in the UK in relation to thermal time. *Plant Pathol.* 54: 144–155.

POLISH SUMMARY

WYSTĘPOWANIE ŁAMLIWOŚCI ŻDŻBŁA NA PSZENICY OZIMEJ W REJONIE POLSKI POŁUDNIOWO-ŚRODKOWEJ

Celem pracy było zebranie danych o występowaniu i cechach gatunków *Oculimacula yallundae* i *O. acuformis* w rejonie Śląska i województwa opolskiego w latach 2003–2008 oraz wykonanie charakterystyki taksonomicznej. Badania podjęto w celu wyjaśnienia równoczesnego występowania początkowych i zaawansowanych objawów łamliwości żdźbła w fazach dojrzewania ziarniaków pszenicy, BBCH 73–77. Miały one także na celu wyjaśnienie zagadnień związanych z procesem infekcji pszenicy ozimej przez patogeny powodujące łamliwość żdźbła w omawianym rejonie. Corocznie na 5–8 plantacjach prowadzono ocenę wzrokową porażenia, a wyliczony średni wskaźnik porażenia wynosił 0,41. Świadczy to o występowaniu choroby, gdy średnio połowa roślin jest zainfekowana w stopniu znacząco obniżającym ich kondycję. Pobierano porażone rośliny do badań laboratoryjnych. Izolowano patogeny. Uzyskiwano kultury wykazujące cechy morfologiczne charakterystyczne dla badanych gatunków. Pomiary wzrostu liniowego kultur badanych grzybów wykazały szybszy rozwój *O. yallundae*. Na wodnym agarze obserwowano ich zarodnikowanie i stwierdzono, że średnia długość zarodników *O. yallundae* wynosiła 61,2 µm, a *O. acuformis* 70 µm.

Analizy PCR kultur grzybów potwierdziły ich przynależność gatunkową. Analiza roślin przy użyciu metody PCR Multiscreen Stem Base Analysis wykazała, że na 19 plantacjach występował gatunek *O. acuformis*, a na pięciu – *O. yallundae*. Gatunek *O. acuformis* był w wyraźnej przewadze stanowiąc 70–100% w stosunku do *O. yallundae*, który występował w 0–30%. Stwierdzone w latach 2003–2008 regularne występowanie *O. acuformis* w badanym rejonie pozwala na wyjaśnienie zagadnienia późno pojawiających się symptomów chorobowych, a także daje wskazówki dla praktyki rolniczej, pozwalające na właściwy dobór fungicydów dla ochrony pszenicy ozimej.